

The Immediate Effect of Chest Mobilization Technique on Dyspnea in Patients of COPD with Restrictive Impairment

Parmar Dharmesh

ARTICLE AUTHORSHIP & AFFILIATION DETAILS

COMMUNICATION DATE: JULY, 15, 2015

ACCEPTANCE DATE: JULY. 05, 2015

DOI:

**PARMAR DHARMESH,
GOVT. PHYSIOTHERAPY COLLEGE,
CIVIL HOSPITAL, AHMEDABAD,
GUJARAT, INDIA**

EMAIL:

DHARMESH_PARMAR62@YAHOO.IN

**KEY WORDS: CHEST MOBILIZATION,
COPD, DYSPNEA, MODIFIED BORG
SCORE.**

TO CITE THIS ARTICLE: Parmar Dharmesh. The immediate effect of chest mobilization technique on dyspnea in patients of COPD with restrictive impairment [Online]. *Journal Of Exercise Science And Physiotherapy*, Vol. 11, No. 2, June 2015: 134-141.

Abstract

Objective of study: to relieve dyspnea in patients of copd with restrictive impairment by chest mobilization technique. **Background:** copd is a primary lung disease but as it advances, there is restriction in chest wall mobility which decreases pulmonary functions and vital capacity of lung. So purpose of this study is to assess the immediate effect of chest mobilization on relieving dyspnea by improving the oxygen saturation. **Materials and methods:** an experimental study was conducted on 30 copd patients having vital capacity <80%, to assess the pre and post differences in modified borg scale by applying chest mobilization technique: rib rotation; lateral flexion, extension, rotation of chest wall and pectoralis major stretching. **Result:** for within group analysis, comparison of data for modified borg score was done using wilcoxon sign rank test. And for between groups analysis was done using mann whitney u test. Statistical analysis showed significant change in modified borg score after application of chest mobilization technique. **Conclusion:** it can be concluded from this study that chest wall mobilization has significant effect on dyspnea in copd patient who is having restrictive impairment of chest wall in later stage of disease.

Introduction

The term Chronic Obstructive Pulmonary Disease (COPD) refers to chronic disorder that disturbs airflow. COPD is a major cause of morbidity and mortality in INDIA (Singh V. et al. 2003). COPD is a preventable and treatable disease with

some significant extra-pulmonary effects that may contribute to the severity in individual patients. Its pulmonary component is characterized by airflow limitation that is not fully reversible. The airflow limitation is usually progressive

and associated with an abnormal inflammatory response of the lung to noxious particles or gases (Goldcopd, 2014). The common task force statement of The American Thoracic Society (ATS) and The European Respiratory Society (ERS) adds to this definition that: "COPD is....primarily caused by cigarette smoking."(Celli BR, 2004) Both GOLD and ATS/ERS agree that COPD is to be suspected when there is a history of exposure to risk factors for the disease, chronic cough, sputum production and/or dyspnoea and that diagnosis must be confirmed by spirometry. When forced expiratory volume in one second (FEV1) divided by forced ventilator capacity (FVC) is <70%, even after administration of a bronchodilator, the diagnosis is confirmed (Singh V. et al. 2003). In patients older than 70 years a somewhat lower ratio (< 65%) has been suggested. (Rabe KF, 2007) Some guidelines claim that besides FEV1/FVC < 70%, the FEV1 should be < 80% of predicted value for diagnosis of COPD (BTS guidelines, 2003).

Normally, people take deep breaths or sigh regularly. These actions stretch the respiratory structures. Patients of COPD with chronic respiratory muscle weakness have reductions in lung volumes and vital capacity (VC) and they may have decreases in lung distensibility with lung volume restriction (De Troyer. et al, 1980). As shown by Mizuri et al, failure to fully expand the lungs causes increases in lung tissue and chest wall elastance and decreases in compliance.

The total mechanical work of breathing (WOB) is the sum of the work of overcoming both the elastic and frictional forces opposing inflation. In healthy adults, about two thirds of the WOB can be attributed to elastic forces opposing ventilation. The remaining third is due to frictional resistance to gas and tissue movement. In diseased states, the WOB can dramatically increase. In patients with restrictive lung disease, work is the integration of the volume-pressure breathing curve. The increase in the WOB is a function of tissue elastance and an inverse function of pulmonary compliance (Slonim NB.et al, 1987).

Failure to take periodic deep breaths can change alveolar surface forces and increase the tendency for alveolar collapse. Gross muscle weakness alters the passive recoil of the thoracic cage, modifying the neutral position at which lung and cage recoil pressures are balanced. (Estenne M. et al, 1977) This results in altered inspiratory muscle length-tension relationships. The lungs and chest walls are susceptible to the effects of incomplete regular mobilization. The tendons and ligaments of the rib cage and the costovertebral and costosternal articulations stiffen, and the latter ankylose, as the intercostal and other respiratory muscles become fibrotic and contracted (Estenne M. et al, 1977).

Expiratory airflow is limited because of the obstruction, leading to air trapping and hyperinflation. This accentuates when the minute ventilation or respiration rate is increased, for example during exercise.

(O'Donnell DE, 2006) The hyperinflation induces increased strain on the respiratory muscles, which are forced to work in a limited range of movement with negative pressure/effort relationship, leading to fatigue and increased dyspnea (Bellemare F. et al, 1983). To avoid the distressing feeling of dyspnoea, the patients with COPD tend to avoid physical exertion and adapt a more sedated lifestyle than healthy elderly subjects (Pitta F. et al, 2005). This, in turn, leads to a vicious cycle of reduced exercise capacity inducing increased dyspnoea during exercise which leads to a further avoidance of exercise and so on. Exercise capacity is impaired in COPD, both peak exercise capacity and functional exercise capacity. Besides lung hyperinflation and physical inactivity, ventilation-perfusion mismatch, hypoxemia, cardio-vascular problems and muscular changes contribute to the reduced exercise capacity. Functional exercise capacity is one of the key prognostic factors of morbidity and mortality in COPD (Calverley PM. Et al, 2005) and correlates strongly with physical activities in daily life (Pitta F. et al, 2005).

Mobilization of rib cage joints appears as a specific aim for physiotherapy, as rib cage mobility seems to be reduced with obstructive lung disease. Chest wall mobilization improves mobility of chest wall, reduces respiratory rate, increases tidal volume, improves ventilation gas exchange, reduces dyspnea, decreases work of breathing and facilitate relaxation (Tabira Kelzuyuki. et al, 2007), (Kozo

Ryo. et al, 1998), (ACCP/AACVPR Pulmonary Rehabilitation Guidelines, 1997), (T.Shioya. et al, 2007), (Kriel, 2005), ([Leelarungrayub D.](#) et al, 2009), (Putt MT. et al, 2008).

Materials & Methods

Study Design: Experimental study (Before and after with control), one time study
 Dependent Variable: Modified Borg score values

Independent Variable: Chest Mobilization tech.

Sample design: Random sampling

Sample size: 30 Patients

Study Setting: General Hospital, Ahmedabad

Selection Criteria

Inclusion Criteria:

- Patients diagnosed as having COPD by the physician. The diagnosis was confirmed by COPD questionnaire,
- Patients with COPD with restrictive impairment (VC<80%),
- Age: >40yrs,
- Sex: male,
- Patients who are able to comprehend commands.

Exclusion Criteria:

- Patients with unstable vital parameters,
- Those who have active lung infection,
- Patients with congenital heart disease, ischemic heart disease, rheumatic heart disease,
- Patients who have recently taken bronchodilator drugs,

- Patients with continuous Oxygen therapy,
- Patients with artificial ventilation,
- Sex: Female Patients

TABLE 1: Difference in means of AGE

GROUP A	55.47±5.06
Group B	57.47±8.22

OUTCOME MEASURE: Modified Borg Score ([Mador MJ, 1590](#)):

0	None
0,5	Very, very light
1	Very light
2	Light
3	Moderate
4	A little intense
5	Intense
6	
7	Very intense
8	
9	Very, very intense
10	Maximum

Figure 1 Modified Borg Score

Patient Instructions for Borg Dyspnoea Scale: “This is a scale that asks you to rate the difficulty of your breathing. It starts at number 0 where your breathing is causing you no difficulty at all and progresses through to number 10 where your breathing difficulty is maximal. How much difficulty is your breathing causing you right now?”

Procedure: 30 patients were randomly selected according to inclusion criteria. PFT of all these patients were done. These patients were divided randomly into two groups (15 in each group), one group was experimental and other was control group. Breathlessness score was measured by Modified Borg Scale before and after giving chest mobilization technique. 3 Repetitions of each maneuver was done. Group A: Chest mobilization and Breathing exercises
Group B: Breathing exercises only

Chest Mobilization tech. are (Tabira Kelzuyuki. et al, 2007): 1.Rib rotation
2.Chest wall rotation 3.Lateral flexion of chest wall 4.Chest wall extension
5.Pectoralis major muscle stretching



Figure 2: Rib rotation



Figure 3: Chest wall rotation



Figure 4: Lateral flexion of chest wall



Figure 5: Chest wall extension



Figure 6: Pectoralis major muscle stretching

RESULT

For within group analysis, comparison of data for Modified Borg Score was done using Wilcoxon Sign Rank Test and for between groups analysis, it was done using Mann Whitney U test.

TABLE 2: Difference in means of MODIFIED BORG SCALE for Dyspnea:

	Before	After	W-value	p value
Group A	2.27±0.88	1.70±0.90	55	0.0020
Group B	2.20±0.86	1.77±0.96	41	0.0371

TABLE 3: The mean of differences of outcome measures between the groups

	MODIFIED BORG SCALE
Group A	1.13±0.58
Group B	0.43±0.59

The MODIFIED BORG SCORE mean of differences shows significant difference between both the groups. (U=174, P=0.0077)

DISCUSSION

Though COPD is obstructive type of pulmonary disease, as disease progresses, there is stiffening of chest wall which gives restrictive pattern to the diseased lung. If this little but important thing is missed in the rehabilitation of COPD patient then it

can hamper the progress of rehabilitation as this restriction alters other physiology of lungs and chest wall and it doesn't allow other rehabilitation protocols to work in improving the condition of the patient.

Kriel & Achmat (2005) had done an investigation into the immediate effect of rib mobilization and sham laser application on chest wall expansion and lung function in healthy asymptomatic males. There was improvement in dyspnea and chest expansion values

Leelarungrayub D, Pothongsunun P, et al. (2009) in their study they found Acute clinical benefits of chest wall-stretching exercise on expired tidal volume, dyspnea and chest expansion in a patient with chronic obstructive pulmonary disease.

Putt MT, Watson M et al. (2008), in their study on "Muscle stretching tech. increases VC and range of motion in patients with COPD" had concluded that the hold and relax technique to the pectoralis major compared with the sham technique produced significant effects on VC ($P < .01$), and rit ($P < .01$) and left ($P < .05$) upper-limb range of motion. There was no significant effect on ACE, XCE, perceived dyspnea, or respiratory rate. There was no order effect for either technique.²⁰

So, in present study, stretching of chest wall muscles like intercostals and pectoralis major muscle is also emphasized by chest mobilization technique.

T.Shioya, M.Satake, et al. (2007), in their study "Combination of chest wall mobilization and respiratory muscle

training in comprehensive outpatient pulmonary rehabilitation improves pulmonary function in patients with COPD" had concluded that combination of chest wall mobilization by squeezing technique, RMT and RMSG in outpatient PR improve pulmonary function, exercise capacity and HRQOL in patients with stable COPD.¹⁶

Above study suggest that chest mobilization can even affects the patient's Quality of life so using of this technique can give a better life to the patient.

CONCLUSION

It can be concluded from the present study that Chest Wall mobilization has significant effect on Dyspnea relief in COPD patient who is having restrictive impairment of chest wall in later stage of disease.

Chest Mobilization is the definite tool for the improving condition of the patient of COPD with restrictive impairment of chest wall. So it should be included as a part of management in the patient of COPD with other exercise treatment program.

LIMITATION OF STUDY:

This study was done on male patients only so future study can be done with taking female patients also in the study so result of this study generalized.

The major limiting factor in present study was smaller sample size. So future study can be done by taking a larger sample.

This was a one time study and no further follow up was taken so could not

assess the long term effect of aerobic exercise on hypertension.

Acknowledgements

I am thankful to the Govt. Physiotherapy College, Civil Hospital, Ahmedabad for support and guidance during the study. Sincere thanks to patients who participated in the study.

Source of funding: By Institute

Conflict of Interest: There is no conflict of interest.

References:

- ACCP/AACVPR Pulmonary Rehabilitation Guidelines Panel: Pulmonary rehabilitation Joint ACCP/AACVPR Evidence based guideline. CHEST:112(5): 1363-1396, 1997.
- BTS guidelines for the management of chronic obstructive pulmonary disease. The COPD Guidelines Group of the Standards of Care Committee of the BTS. Thorax 1997;52 Suppl 5:S1-28.
- Bellemare F, Grassino A. Force reserve of the diaphragm in patients with chronic obstructive pulmonary disease. J Appl Physiol. 1983;55:8-15.
- Celli BR, MacNee W. Standards for the diagnosis and treatment of patients with COPD: a summary of the ATS/ERS position paper. Eur Respir J 2004;23(6):932-46.
- Calverley PM, Rennard SI, Wouters EF, Agusti A, Anthonisen N, et al. Proposal for a multidimensional staging system for chronic obstructive pulmonary disease. Respir Med 2005;99(12):1546-54.
- De Troyer, A, Borenstein, S, Cordier, R Ankylosis of lung volume restriction in patients with respiratory muscle weakness. Thorax 1980;35:603-610
- Estenne, M, Heilporn, A, Delhez, L, et al Chest wall stiffness in patients with chronic respiratory muscle weakness. Am Rev Respir Dis. 1977;115:389-395
- Global Initiative for Chronic Obstructive Lung Disease. Strategy for the Diagnosis, Management and Prevention of Chronic Obstructive Pulmonary Disease. <http://www.goldcopd.com>; January 14,2007.
- Kozu Ryo, Yanase Kenji et. al.: Influence of chest expansion on pulmonary function and Dyspnoea in patients with chronic obstructive pulmonary disease. The Journal of Japanese Physical Therapy Association (JPTA) Vol. 25, No. 6(19980930)
- Kriel, Achmat. An investigation into the immediate effect of rib mobilization and sham laser application on chest wall expansion and lung function in healthy asymptomatic males : a pilot study. Dept. of Chiropractic, Durban Institute of Technology.2005.
- Leclarungrayub D, Pothongsunun P, et al. Acute clinical benefits of chest wall-stretching exercise on expired tidal volume, dyspnea and chest expansion in a patient with chronic obstructive pulmonary disease: a single case study. J Bodyw Mov Ther. 2009 Oct;13(4):338-43.
- Mador MJ, Rodis A, Magalang UJ. Reproducibility of Borg scale measurements of dyspnea during exercise in patients with COPD. Chest. 1995 Jun;107(6):1590-7.
- O'Donnell DE (2006). "Hyperinflation, Dyspnea, and Exercise Intolerance in Chronic Obstructive Pulmonary Disease. American Thoracic Society 3: 180–184.
- Pitta F, Troosters T, Spruit MA, Probst VS, Decramer M, Gosselink R. Characteristics of physical activities in daily life in chronic obstructive pulmonary disease. Am J Respir Crit Care Med 2005;171(9):972-7.
- Putt MT, Watson M. et al.: Muscle stretching technique increase vital capacity and range of motion in patients with chronic obstructive pulmonary disease. Arch Phys Med Rehabil. 2008. Jun: 89(6):1103-7.
- Rabe KF, Hurd S, Anzueto A, et al. (2007). "Global Strategy for the Diagnosis, Management, and Prevention of Chronic Obstructive Pulmonary Disease: GOLD Executive Summary". Am. J. Respir. Crit. Care Med. 176 (6): 532–55

Singh V, Khandelwal DC, Khandelwal R, Abusaria S. Pulmonary rehabilitation in patients with chronic obstructive pulmonary disease. *Indian J Chest Dis Allied Sci.* 2003 Jan-Mar;45(1):13-7

Slonim, NB, Hamilton, LH *Respiratory physiology.* 5th ed. 1987,26-38 Mosby. St. Louis.

Tabira Kelzuyuki, Sekikawa Noriko. et. al.: The immediate effect of chest mobilization tech. in patients of COPD. *The Journal of Japanese*

Physical Therapy Association.(JPTA) Vol. 34, No. 2(20070420) pp. 59-64

T.Shioya, M.Satake, H.Takahashi, K.Sugawara, C.Kasai, N.Kiyokawa, T.Watanabe, S.Fujii, M.Honma. Combination of chest wall mobilization and respiratory muscle training in comprehensive outpatient pulmonary rehabilitation improves pulmonary function in patients with COPD. Department of Rehabilitation, Akita City General Hospital, Akita, Japan. 2007



Conflict of Interest None Declared