

HOSTED BY



ELSEVIER

Contents lists available at ScienceDirect

Asian Pacific Journal of Tropical Biomedicine

journal homepage: www.elsevier.com/locate/apjtbOriginal article <http://dx.doi.org/10.1016/j.apjtb.2015.09.012>

Screening for anti-pancreatic lipase properties of 28 traditional Thai medicinal herbs

Ananya Dechakhamphu^{1*}, Nattapong Wongchum²¹Thai Traditional Medicine Program, Faculty of Thai Traditional and Alternative Medicine, Ubon Ratchathani Rajabhat University, Ubonratchathani 34000, Thailand²Biology Program, Faculty of Science, Ubon Ratchathani Rajabhat University, Ubonratchathani 34000, Thailand

ARTICLE INFO

Article history:

Received 7 Jul 2015

Received in revised form 20 Jul 2015

Accepted 15 Aug 2015

Available online 3 Oct 2015

Keywords:

Obesity

Medicinal plant

Herb

Anti-pancreatic lipase

Phenolic

Alkaloid

Flavonoid

ABSTRACT

Objective: To screen the effect of 28 medicinal plants on inhibition of pancreatic lipase and evaluate the phytochemical contents of extracts.**Methods:** The ethanolic extracts of 28 traditional Thai herbal medicines were assayed for their *in vitro* activities against porcine pancreatic lipase using *p*-nitrophenyl butyrate as a substrate. Quantitative estimation of flavonoids, phenolics, and alkaloids was done.**Results:** Extracts from four herbs, *Memecylon edule* Roxb., *Garcinia vilersiana* Pierre, *Cryptolepis elegans* Wall. and *Phyllanthus chamaepeuce* Ridl., at a concentration of 100 µg/mL, strongly inhibited porcine pancreatic lipase by 90.97%, 92.04%, 94.64% and 95.38%, respectively. There was a significant positive correlation between phenolic content and inhibition activity. Inhibition activity was significantly correlated with flavonoid and with alkaloid contents.**Conclusions:** From this result, it could be concluded that herbs represent a rich of anti-pancreatic lipase compounds, in particular, *Cryptolepis elegans* Wall. and *Phyllanthus chamaepeuce* Ridl. It is suggested that the phytochemical compounds from these plants may be applied for the prevention and treatment of obesity or hyperlipidemia.

1. Introduction

Obesity is becoming a worldwide epidemic, resulting in a major risk factor for coronary heart diseases including diabetes mellitus, metabolic syndrome, stroke, and some cancers [1]. Globally, around 39% of adults aged 18 and over were overweight in 2014 and 13% of them were clinically obese [2]. Therefore, prevention and treatment of obesity become an important factor for a healthy condition. The reduction of nutrient digestion and absorption by developing of enzyme inhibitors without altering major mechanism in gastrointestinal system became the most important strategies in the treatment of obesity [3,4]. The major source of unwanted calories is dietary lipids, therefore, lipid metabolism play a major role in maintaining energy homeostasis [5]. The identification and

characterization of several enzymes involved in lipid metabolism have yielded a rich pool of potential targets for drugs to treat obesity and other metabolic disorders [6]. Pancreatic lipase is the key enzyme for dietary fat digestion and absorption. Therefore, inhibition of this enzyme would be in effect to reduce lipid absorption from intestine and lead to a consequence suppress of weight gain. Orlistat, a specific drug for inhibiting pancreatic lipase that reduces dietary fat absorption by 30%, has been approved for clinical use [4,7,8]. However, Orlistat can result in adverse side effects, such as fecal incontinence, flatulence, and steatorrhea [9,10]. Therefore, the investigation to find new safety medication for anti-obesity is still needed. The significant progress of the development of anti-obesity from medicinal plants has provided potential therapeutic targets for obesity [11,12]. It has been recently reported that natural compounds from plants and other organisms have been approved as anti-pancreatic lipase activities. For example, ethanolic extract from *Terminalia paniculata* bark [13], polyphenols from Oolong tea [14], *Abroma augusta* extract [15], pomegranate leaves ethanol extract [16], and other components from other kinds of herbs. However, it remains necessary to search for more effective lipase inhibitors from traditional herbs. In this study, we investigated the ethanolic

*Corresponding author: Ananya Dechakhamphu, Thai Traditional Medicine Program, Faculty of Thai Traditional and Alternative Medicine, Ubon Ratchathani Rajabhat University, Ubonratchathani 34000, Thailand.

Tel: +66 942818585

E-mail: ananya.d@ubru.ac.th

Peer review under responsibility of Hainan Medical University.

Foundation Project: Supported by Ubon Ratchathani Rajabhat University through Plant Genetic Conservation Project Under The Royal Initiative of Princess Maha Chakri Sirindhorn Program (Grant No.UBRU_RSPG_2556).

extracts of 28 traditional Thai herbal medicines for their *in vitro* activities against porcine pancreatic lipase using *p*-nitrophenyl butyrate (*p*-NPB) as a substrate.

2. Materials and methods

2.1. Materials

Porcine pancreatic lipase, *p*-NPB, morpholinepropanesulphonic acid, quercetin, colchicine and gallic acid were purchased from Sigma Aldrich. The fresh leaves of 28 plants were collected from Plant Genetic Conservation Forest, Ubon Ratchathani Rajabhat University, Ubonratchathani Province, Thailand. All plant species were identified and authenticated by Mr. Prakorb Boonma, Senior Plant Taxonomist, Ubon Ratchathani Rajabhat University, Thailand.

2.2. Preparation of plant extracts

The leaves were dried in hot air oven at 50 °C for 48 h and grounded into fine powder. A total of 50 g powder was extracted in 95% ethanol and concentrated at 55 °C in a rotary vacuum evaporator (Heidorf, Germany). The obtained extracts were stored at –20 °C until use.

2.3. Porcine pancreatic lipase inhibition assay

Lipase activity was measured using *p*-NPB as a substrate. The method was modified from the previously described by Kim *et al.* [17]. Briefly, an enzyme buffer was prepared by the addition 30 µL of solution of porcine pancreatic lipase (2.5 mg/mL in 10 mmol/L morpholinepropanesulphonic acid and 1 mmol/L ethylenediamine tetraacetic acid, pH 6.8) to 850 µL of Tris buffer (100 mmol/L Tris–HCl and 5 mmol/L CaCl₂, pH 7.0). Then, either 100 µL of the plant extracts (100 µg/mL) or Orlistat was added and incubated for 15 min at 37 °C. Ten microliter of substrate (10 mmol/L *p*-NPB in dimethyl formamide) was then added and incubated for 30 min at 37 °C. Lipase activity was determined by measuring the hydrolysis of *p*-NPB to *p*-nitrophenol at 405 nm using an ELISA reader (Biochrome, England). The inhibitory activity (I) was calculated according to the following formula:

$$I\% = \left(1 - \frac{B-b}{A-a}\right) \times 100$$

where A is the activity of the enzyme without inhibitor, and a is the negative control without inhibitor; B is the activity of the enzyme with inhibitor, and b is the negative control with inhibitor.

2.4. The half maximal inhibitory concentration (IC₅₀) determination

The IC₅₀ value of extracts was determined at a concentration of 500.0, 250.0, 125.0, 100.0, 25.0, 12.5 and 5.0 µg/mL. Orlistat was used as a positive control. IC₅₀ value was calculated by the following formula:

$$IC_{50} = (50\% - Low_{Inh\%}) / (High_{Inh\%} - Low_{Inh\%}) \times (High_{Conc} - Low_{Conc}) + Low_{Conc}$$

formula: where Low_{Inh}%/High_{Inh}% signify % inhibition directly below/above 50% inhibition, and Low_{Conc}/High_{Conc} are the corresponding concentrations of extract.

2.5. Determination of total phenolic content

According to a previously described protocol [18], Folin–Ciocalteu reagent was used to determine the total phenolic content of extracts. Absorbance was measured at 725 nm. All tests were performed 6 times. The phenolic content was calculated based on a gallic acid standard curve.

2.6. Determination of total flavonoid content

Total flavonoid content was determined according to a previously discussed method [18] using quercetin as a standard. The absorbance was measured at 510 nm. The flavonoid content was calculated based on a quercetin standard curve.

2.7. Quantification of alkaloid content

Quantification of alkaloid content for extracts was carried out using a method described earlier [19]. The absorbance was taken at 500 nm and all tests were performed 6 times. The alkaloid content was evaluated based on the colchicine standard curve.

2.8. Statistical analysis

Statistical analysis of the data was performed using the SPSS 16.0 program. The comparison between Orlistat control and extract group was conducted using the Mann–Whitney *U* test, and the correlations between parameters were determined using the Spearman's rank test.

3. Results

A total of 28 extracts were prepared from leaf part of the traditional Thai medicinal herbal medicines and were tested at a concentration of 100 µg/mL for porcine pancreatic lipase inhibition (Table 1).

Table 1

Lipase inhibitory effects of 28 selected traditional Thai medicinal herbs. %.

Scientific name	Family	Inhibition
<i>Artocarpus lakoocha</i> Roxb.	Moraceae	12.68 ± 1.10*
<i>Azadirachta indica</i> A. Juss.	Meliaceae	34.85 ± 2.70*
<i>Belamcanda chinensis</i> (L.) DC.	Iridaceae	6.58 ± 1.30*
<i>Brucea javanica</i> (L.) Merr.	Simaroubaceae	22.83 ± 3.20*
<i>Canarium subulatum</i> Guill.	Burseraceae	–26.84 ± 1.70*
<i>Canthium berberidifolium</i> Geddes	Rubiaceae	–5.76 ± 0.70*
<i>Congea siamensis</i> Fletcher	Verbenaceae	70.00 ± 1.30
<i>Cratoxylum formosum</i> (Jack) Dyer T.	Guttiferae	67.47 ± 1.30
<i>C. elegans</i>	Asclepiadaceae	94.64 ± 1.10
<i>Dillenia ovata</i> Wall.	Dilleniaceae	5.97 ± 0.60*
<i>Diospyros filipendula</i> Pierre ex Lecomte	Ebenaceae	59.15 ± 0.90*
<i>Eurycoma longifolia</i> Jack	Simaroubaceae	1.04 ± 1.20*
<i>Ficus foveolata</i> Wall.	Moraceae	6.99 ± 1.30*
<i>Garcinia cowa</i> Roxb. ex DC.	Guttiferae	14.91 ± 2.40*

(continued on next page)

Table 1 (continued)

Scientific name	Family	Inhibition
<i>G. vilersiana</i>	Guttiferae	92.04 ± 1.10
<i>Gomphia serrata</i> (Gaertn.) Kanis	Ochnaceae	30.18 ± 2.70*
<i>Litchi chinensis</i> Sonn.	Sapindaceae	12.28 ± 0.40*
<i>Mecycylon edule</i> Roxb.	Melastomataceae	90.97 ± 0.30
<i>Naringi crenulata</i> (Roxb) Nicolson ST.	Rutaceae	-0.67 ± 1.20*
<i>P. chamaepeuce</i>	Euphorbiaceae	95.38 ± 2.40
<i>Phyllanthus gomphocarpus</i> Hook. f.	Euphorbiaceae	-5.19 ± 2.10*
<i>Prismatomeris sessiliflora</i> Pierre ex Pitard	Rubiaceae	80.11 ± 0.20
<i>Quercus kingiana</i> Craib	Fagaceae	-12.70 ± 1.10*
<i>Salacia verrucosa</i> Wight	Celastraceae	-6.09 ± 0.80*
<i>Spatholobus harmandii</i> Gagnep.	Fabaceae	24.28 ± 0.60*
<i>Syzygium gratum</i> (Wight) S.N. Mitra var. <i>gratum</i>	Myrtaceae	-34.93 ± 2.40*
<i>Tiliacora triandra</i> (Colebr.) Diels	Menispermaceae	-19.47 ± 2.20*
<i>Xantolis cambodiana</i> (Pierre ex Dubarb) P. Royen	Sapotaceae	87.55 ± 2.20
Orlistat		96.56 ± 0.20

* $P < 0.05$ compared to Orlistat, data were presented as mean ± SD ($n = 3$). The final concentration of the extracts used in this experiment was 100 µg/mL. *G. vilersiana*: *Garcinia vilersiana* Pierre; *C. elegans*: *Cryptolepis elegans* Wall.; *P. chamaepeuce*: *Phyllanthus chamaepeuce* Ridl.

Among 28 herbs, 9 plants were found to have strong inhibitory activity of >50% against porcine pancreatic lipase: 59.15% with *Diospyros filipendula*, 67.47% with *Cratoxylum formosum*, 70.00% with *Congea siamensis*, 80.11% with *Prismatomeris sessiliflora*, 87.55% with *Xantolis cambodiana*, 90.97% with *Mecycylon edule*, 92.04% with *G. vilersiana*, 94.64% with *C. elegans*, and 95.38% with *P. chamaepeuce*. The extracts of *Gomphia serrata* and *Azadirachta indica* showed >30% inhibition of pancreatic lipase. There were several herbs showed slightly effect on inhibition against pancreatic lipase. Besides, some of herbs increased in the activity of pancreatic lipase, such as *Syzygium gratum*, which promoted the activity of enzyme by 34.93%.

The different concentration of crude extracts of *G. vilersiana*, *C. elegans*, and *P. chamaepeuce* were measured for IC_{50} at a concentration of 500.0, 250.0, 125.0, 100.0, 25.0, 12.5 and 5.0 µg/mL. The extracts of *G. vilersiana*, *C. elegans* and *P. chamaepeuce* had IC_{50} values of 0.11, 0.30 and 0.45 µg/mL, respectively. Whereas, Orlistat had IC_{50} value of 0.08 µg/mL (Figure 1).

The correlations between phytochemical content measured in extracts and inhibition activity were shown in Figure 2. There

was a significant positive correlation between phenolic content and inhibition activity. Inhibition activity was significantly correlated with flavonoid and alkaloid contents.




Figure 1. The half maximal inhibitory concentration (IC_{50}) of selected crude extracts.

* $P < 0.05$ compared to Orlistat, data were presented as mean ± SD ($n = 3$).

4. Discussion

Pancreatic lipase is a key enzyme for lipid absorption by hydrolysis of total dietary fats. Therefore, inhibition of pancreatic lipase is suggested to be an effective therapy in the regulation of obesity. Although Orlistat has anti-obesity effects by inhibiting pancreatic lipase activity, however, it can cause adverse side effects, such as fecal incontinence, flatulence, and steatorrhea [9,10]. Therefore, the investigation of new agent for pancreatic lipase inhibitor is still needed. Our finding is the first time to show that crude extracts of *G. vilersiana*, *C. elegans* and *P. chamaepeuce* exhibited strong anti-lipase activity. This suggests that these herbs seem to be the potential candidates as the inhibitor of pancreatic lipase. However, further *in vivo* studies on animal model must be conducted in order to confirm this hypothesis.

Our results showing a significant positive correlation between phenolic, flavonoid, alkaloid contents and inhibition activity, which provide strong support that these phytochemical compounds are key agents for pancreatic lipase inhibition. This assumption is further supported by the previous results showing that phenolic compounds exhibit the ability to inhibit pancreatic




Figure 2. Correlation of phenolic, flavonoid, and alkaloid contents with porcine pancreatic lipase inhibition activity.

A: Phenolic content; B: Flavonoid content; C: Alkaloid content. Data from all plant extracts were pooled. Spearman correlation coefficients and P -value from a total sample analyses were listed.

lipase activity [14,20–23]. Published research also reported that flavonoids and alkaloid be able to inhibit pancreatic lipase [24,25]. The study *in vivo* model indicated that polyphenols and flavonoid glycoside derived from *Salix matsudana* leaf showed decreased in body weight gain in Wistar rats [21]. According to these reports, we hypothesized that these three compounds may be the main contributors to the inhibition of pancreatic lipase. Experimental proof for this assumption is now required.

Although high activities on pancreatic lipase inhibition were detected in crude extracts of *G. vilersiana*, *C. elegans* and *P. chamaepeuce*, the further investigation for both *in vitro* and *in vivo* should be performed to elucidate the bioactive compounds, to clarify the molecular mechanism and to verify the main effective phytochemicals in these three candidates which are responsible for the inhibition of pancreatic lipase activity. For example, caffeine, chlorogenic acid, feruloylquinic acids derived from *Coffea canephora* showed to inhibit pancreatic lipase *in vitro* and decrease body weight gain of mice by 157% [26]. It has been reported that crocetin derived from *Gardenia jasminoides* showed inhibitory activity of pancreatic lipase *in vitro* and 25% decreased in body weight gain of Triton WR-1339-induced hyperlipidemia mice [27,28].

We concluded that phenolic, flavonoid and alkaloid compounds in *G. vilersiana*, *C. elegans* and *P. chamaepeuce* are key agents for pancreatic lipase inhibition *in vitro*. These three plants should be explored as dietary supplements or nutraceutical foods with anti-obesity properties. To the best of our knowledge, the purification of active compounds of certain pancreatic lipase inhibitor is under investigation.

Conflict of interest statement

We declare that we have no conflict of interest.

Acknowledgments

This research was supported by grants funded by Ubon Ratchathani Rajabhat University through Plant Genetic Conservation Project Under The Royal Initiative of Princess Maha Chakri Sirindhorn Program (Grant No.UBRU_RSPG_2556). The authors thank the Faculty of Thai Traditional and Alternative Medicine, Ubon Ratchathani Rajabhat University for providing instruments. We thank Dr. Chayada Danuwong, English Program, Ubon Ratchathani Rajabhat University, for proofreading the manuscript.

References

- [1] Barnes A. Overweight versus obese: different risk and different management. *Tex Heart Inst J* 2015; **42**(3): 237-8.
- [2] World Health Organization. Global Health Observatory (GHO) data. Overweight and obesity. Geneva: World Health Organization; 2014. [Online] Available from: http://www.who.int/gho/ncd/risk_factors/overweight/en/ [Accessed on 1st June, 2015]
- [3] Yun JW. Possible anti-obesity therapeutics from nature—a review. *Phytochemistry* 2010; **71**: 1625-41.
- [4] Martin KA, Mani MV, Mani A. New targets to treat obesity and the metabolic syndrome. *Eur J Pharmacol* 2015; <http://dx.doi.org/10.1016/j.ejphar.2015.03.093>.
- [5] Loli H, Narwal SK, Saun NK, Gupta R. Lipases in medicine: an overview. *Mini Rev Med Chem* 2015; **15**(14): 1209-16.
- [6] Sukhdev S, Singh KS. Therapeutic role of phytomedicines on obesity: importance of herbal pancreatic lipase inhibitors. *Int Res J Med Sci* 2013; **1**(9): 15-26.
- [7] Cheung BMY, Cheung TT, Samaranyake NR. Safety of anti-obesity drugs. *Ther Adv Drug Saf* 2013; **4**(4): 171-81.
- [8] Hill JO, Hauptman J, Anderson JW, Fujioka K, O'Neil PM, Smith DK, et al. Orlistat, a lipase inhibitor for weight maintenance after conventional dieting: a 1-y study. *Am J Clin Nutr* 1999; **69**: 1108-16.
- [9] Birari RB, Bhutani KK. Pancreatic lipase inhibitors from natural sources: unexplored potential. *Drug Discov Today* 2007; **12**: 879-89.
- [10] Weigle DS. Pharmacological therapy of obesity: past, present, and future. *J Clin Endocrinol Metab* 2003; **88**: 2462-9.
- [11] Hasani-Ranjbar S, Jouyandeh Z, Abdollahi M. A systematic review of anti-obesity medicinal plants—an update. *J Diabetes Metab Disord* 2013; **12**: 28.
- [12] Ado MA, Abas F, Mohammed AS, Ghazali HM. Anti- and pro-lipase activity of selected medicinal, herbal and aquatic plants, and structure elucidation of an anti-lipase compound. *Molecules* 2013; **18**: 14651-69.
- [13] Mopuri R, Meriga B. Anti-lipase and anti-obesity activities of *Terminalia paniculata* bark in high calorie diet-induced obese rats. *Glob J Pharmacol* 2014; **8**(1): 114-9.
- [14] Nakai M, Fukui Y, Asami S, Toyoda-Ono Y, Iwashita T, Shibata H, et al. Inhibitory effects of oolong tea polyphenols on pancreatic lipase *in vitro*. *J Agric Food Chem* 2005; **53**(11): 4593-8.
- [15] Gupta N, Ganeshpurkar A, Jatav N, Bansal D, Dubey N. *In vitro* prevention of chick pancreatic lipase activity by *Abroma augusta* extract. *Asian Pac J Trop Biomed* 2012; **2**(Suppl 2): S712-5.
- [16] Adnyana IK, Sukandar EY, Yuniarto A, Finna S. Anti-obesity effect of the pomegranate leaves ethanol extract (*Punica granatum L.*) in high-fat diet induced mice. *Int J Pharm Pharm Sci* 2014; **6**(4): 626-31.
- [17] Kim YS, Lee YM, Kim H, Kim J, Jang DS, Kim JH, et al. Anti-obesity effect of *Morus bombycis* root extract: anti-lipase activity and lipolytic effect. *J Ethnopharmacol* 2010; **130**: 621-4.
- [18] Das A, Chaudhuri D, Mandal N, Chatterjee A. Study of antioxidant and reactive oxygen species scavenging activity of the edible tuber of “greater yam” (*Dioscorea alata L.*) from North-east India. *Asian J Pharm Clin Res* 2012; **5**(3): 74-84.
- [19] Ghate N, Chaudhuri D, Mandal N. *In vitro* antioxidant and free radical scavenging assessment of *Tinospora cordifolia* stem with DNA protective potential. *Int J Pharm Bio Sci* 2013; **4**(1): 373-88.
- [20] Yuda N, Tanaka M, Suzuki M, Asano Y, Ochi H, Iwatsuki K. Polyphenols extracted from black tea (*Camellia sinensis*) residue by hot-compressed water and their inhibitory effect on pancreatic lipase *in vitro*. *J Food Sci* 2012; **77**(12): H254-61.
- [21] Han LK, Sumiyoshi M, Zhang J, Liu MX, Zhang XF, Zheng YN, et al. Anti-obesity action of *Salix matsudana* leaves (part 1). Anti-obesity action by polyphenols of *Salix matsudana* in high fat-diet treated rodent animals. *Phytother Res* 2003; **17**: 1188-94.
- [22] Jeong JY, Jo YH, Kim SB, Liu Q, Lee JW, Mo EJ, et al. Pancreatic lipase inhibitory constituents from *Morus alba* leaves and optimization for extraction conditions. *Bioorg Med Chem Lett* 2015; **25**(11): 2269-74.
- [23] Uchiyama S, Taniguchi Y, Saka A, Yoshida A, Yajima H. Prevention of diet-induced obesity by dietary black tea polyphenols extract *in vitro* and *in vivo*. *Nutrition* 2011; **27**(3): 287-92.
- [24] Lee EM, Lee SS, Chung BY, Cho JY, Lee IC, Ahn SR, et al. Pancreatic lipase inhibition by C-glycosidic flavones isolated from *Eremochloa ophiuroides*. *Molecules* 2010; **15**: 8251-9.
- [25] Lunagariya NA, Patel NK, Jagtap SC, Bhutani KH. Inhibitors of pancreatic lipase: state of the art and clinical perspectives. *EXCLI J* 2014; **13**: 897-921.
- [26] Shimoda H, Seki E, Aitani M. Inhibitory effect of green coffee bean extract on fat accumulation and body weight gain in mice. *BMC Complement Altern Med* 2006; **6**: 9.
- [27] Lee IA, Lee JH, Baek NI, Kim DH. Antihyperlipidemic effect of crocin isolated from the fructus of *Gardenia jasminoides* and its metabolite crocetin. *Biol Pharm Bull* 2005; **28**: 2106-10.
- [28] Sheng L, Qian Z, Zheng S, Xi L. Mechanism of hypolipidemic effect of crocin in rats: crocin inhibits pancreatic lipase. *Eur J Pharmacol* 2006; **543**: 116-22.