

Association between Serum Ferritin and Thyroid profile in Radiologically proven patients with Goitre with Thyroiditis

Monu Sarin¹, Poonam Arora^{2*}, SC Mohapatra³

¹Associate Professor, Dept. of Radiodiagnosis & Imaging, ²Assistant Professor, Dept. of Biochemistry, ³Dean Academic Affairs & Senior Professor, Dept. of Community Medicine, SGT Medical College & Hospital, Gurugram, Haryana

***Corresponding Author:**

Email: drpoonamarora2017@gmail.com

Abstract

Introduction: Thyroid peroxidase (TPO) is a membrane-bound glycosylated hemoprotein that has a key role in the biosynthesis of thyroid hormones by organification. Ferritin is an iron storage protein found in almost all of the body tissues. Serum ferritin levels also have been reported to be altered in patients with thyroid disease.

Objective: The aim of this study is to establish a correlation between serum ferritin and thyroid hormones in thyroid disorder patients.

Materials and Method: This was a cross sectional study of 80 subjects, visiting medical outpatient department of SGT Medical College and Hospital, Budhera, Gurugram, Haryana. Patients were subjected to ultrasound. Radiologically proven Thyroid disorder patients sample were analysed for the measurement of T₃, T₄, TSH (by CLIA) and Serum Ferritin (by ELISA).

Results: The mean T₃, T₄, TSH and serum ferritin levels showed highly significant difference with $p < 0.001$ in hypothyroid group compared to controls

Conclusion: Low serum ferritin levels are associated with hypothyroidism. The estimation of iron storage protein ferritin may help in understanding the pathophysiology, diagnosis and treatment of thyroid disorders.

Keywords: Serum Ferritin, Thyroid Profile, Hypothyroidism.

Introduction

Several minerals and trace elements such as iodine, iron, selenium, and zinc are essential for normal thyroid hormone metabolism. Iodine has an important role in the synthesis of thyroid hormones; selenium is a component of the deiodinase enzymes that converts T₄ to T₃. It also protects the thyroid gland from damage by excessive iodide exposure. Zinc appears to be involved in thyroid conversion. Low iron, or more specifically, low ferritin, is one of the most overlooked causes of thyroid dysfunction.⁽¹⁾

Thyroid hormones plays a central role in differentiation, development, and maintenance of body homeostasis.⁽²⁾ Thyroid hormone performs a role in hemoglobin synthesis in adults and maturation of hemoglobin in fetus; and hypothyroidism leads to anemia via reducing oxygenation process through disturbing hematopoietic process.⁽³⁾ It is considered that thyroid hormones influence hematopoiesis through an increase in erythropoietin generation or hematopoietic factors by non-erythroid cells. The actions of T₃, like steroid hormones, are mediated through intracellular T₃-receptor proteins (TRs), which act predominantly to modulate transcription by binding to specific T₃ response elements in target genes. T₃ also exerts important effects at the post-transcriptional level to regulate the expression of several genes.⁽⁴⁾

Ferritin is an iron storage protein found in almost all of the body tissues. Serum ferritin levels also have been reported to be altered in patients with thyroid disease. Thus, changes in the serum concentrations of ferritin reflect thyroid function. Thyroid peroxidase

(TPO) is a membrane-bound glycosylated hemoprotein that has a key role in the biosynthesis of thyroid hormones by organification. Iron deficiency has been reported to impair the body's ability to make its own thyroid hormone, which could increase need for thyroid medication.⁽²⁾

Several studies have documented an association between T₃ levels and ferritin expression. Furthermore, administration of T₃ to hypothyroid individuals produced a significant increase in the serum ferritin level.⁽¹⁾ Although the cause of T₃-induced increase in the serum ferritin levels in human is unknown; increased synthesis of ferritin in the liver may well be an important contributor. These links between T₃ and the regulation of ferritin expression suggest that a positive correlation exists between the levels of T₄/T₃ and ferritin in serum. Thus, it has been suggested that serum ferritin measurement could be useful for the evaluation of thyroid hormone action on peripheral tissues.⁽⁵⁾ Therefore the present study was designed to assess serum ferritin levels in hypothyroid patients.

Materials and Method

The study was carried out in the Department of Radiodiagnosis and Department of Biochemistry, SGT Medical College, Budhera, Gurugram, Haryana. A total of 40 newly diagnosed thyroid disorder patients were compared with 40 age and sex matched normal healthy controls. A written informed consent was also taken from the cases with detailed history.

Inclusion Criteria: Clinically and radiologically diagnosed patients with Goitre with Thyroiditis in the age group of 20-60 years.

Controls: Healthy controls in the age group of 20-60 years.

Exclusion criteria:

- Pregnancy
- Hepatic disorder
- Renal diseases
- Polycystic ovarian syndrome

On Ultrasound, diffuse or multifocal decrease in echogenicity was the hallmark of many types of Thyroiditis. On color Doppler study increased blood flow was noted throughout the gland Enlargement of the gland was also noted. From the radiologically proven thyroid disorder patients, 4 ml of fasting blood sample was collected and centrifuged for serum separation. Thyroid function was assessed by quantitative estimation of T₃, T₄ and TSH levels in serum performed using SEIMENS-Centaur CP analyzer based on chemiluminescent immuno assay (CLIA). Serum ferritin was assessed by ELISA method by commercially available kit provided by Calbiotech.

Statistical Analysis: Statistical analysis was done using the SPSS software version 24. The data was represented by counts, percentage and mean± standard deviation. Statistical analysis of the biochemical parameters, T₃, T₄, TSH and serum ferritin were done by t-test to compare these parameters in cases and controls. A *p*-value of <0.05 was considered significant and *p*-value >0.05 as non-significant.

Results

Of the 40 cases 27 were females and 17 were males. Mean ±SD of age among cases and controls were 33.65± 9.46 years and 35.78± 11.28 years respectively. There was no significant difference with respect to age distribution in cases and controls (*p*>0.05).

Table 1: Ferritin levels in newly diagnosed hypothyroid patients

Parameters	Controls	Cases	p-Value
T ₃ (ng/ml)	1.23± 0.78	0.51±0.87	<0.001
T ₄ (µg/dl)	9.87±1.68	4.23±0.86	<0.001
TSH (mIU/ml)	2.68±0.34	8.95±1.14	<0.001
Serum Ferritin (ng/ml)	72.43±9.20	19.10±3.40	<0.001

The mean T₃, T₄, TSH and serum ferritin levels showed highly significant difference with *p*<0.001 in hypothyroid group compared to controls (Table 1).

Discussion

Thyroid gland has a critical effect on erythropoiesis by induction of erythropoietin production and proliferation of erythroid progenitors.^(6,7) Several studies

showed a relationship between body iron status (indulging iron metabolism, ferritin, and serum iron) and thyroid hormones function.^(8,9)

Our study shows that there is a state of low ferritin concentration in patients with hypothyroidism. It is observed that serum T₃ and T₄ levels were significantly lower in cases as compared to healthy controls (*p*<0.001), suggesting that depletion of iron stores may decrease these hormones. Our results are consistent with various studies.^(2,10,11) Iron deficiency impairs thyroid hormone synthesis by reducing activities of heme-dependent thyroid peroxidase (TPO). TPO is a membrane-bound glycosylated hemoprotein that plays a key role in the biosynthesis of thyroid hormones. This enzyme is responsible for the oxidation of iodide and binding of iodine to tyrosyl residue of thyroglobulin (organification).⁽¹²⁾

It has been suggested in various studies that thyroid hormones regulate ferritin expression. The iron regulatory protein (IRP, previously known as the iron-responsive element-binding protein, IRE-BP, and iron-responsive factor, IRF) is a transacting RNA-binding protein that binds with high affinity to conserved stem-loop structures, iron-responsive elements (IREs), present in the ferritin, and transferrin receptor (TfR). The IRP has a key a role in the regulation of iron homeostasis.⁽¹³⁾ In the absence of iron, the IRP binds to the IRE in the 5'-untranslated region (5'-UTR) of ferritin and represses translation.⁽¹⁴⁾ Binding of the IRP to IREs in the 3'-untranslated region (3'-UTR) of TfR mRNA stabilizes the mRNA and prevents its degradation.⁽¹⁵⁾ In iron-replete states, the reverse holds, which results in increased ferritin translation and decreased TfR mRNA stability. This reciprocal regulation is achieved at the post-translational level and is independent of new protein synthesis.⁽¹⁶⁾

Conclusion

Our study suggest that a significant difference in ferritin levels in hypothyroid patients could be a reflection of disturbed activities of iron dependent thyroid peroxidase (TPO) enzyme which impairs the metabolism of thyroid hormones but the mechanism by which thyroid hormone status alters ferritin levels is not well known. There is a need for further studies to evaluate the affect of thyroid hormones on ferritin metabolism for the diagnosis and treatment of thyroid illness.

Acknowledgement

Author acknowledges the immense help received from the scholars whose articles are cited and included in references of this manuscript. The author is also grateful to authors/editors/ publishers of all those articles, journals and books from where the literature for this article has been reviewed and discussed.

References

1. Takamatsu J, Majima M, Milki K, Kuma K, Mozai T. Serum ferritin as a marker of thyroid hormone action on peripheral tissues. *J Clin Endocrinol Metabol* 1985; 61 (4): 672-6.
2. Sachdeva A, Singh V, Malik I, Roy PS, Madaan H, Nair R. Association between serum ferritin and thyroid hormone profile in hypothyroidism. *Int J Med Sci Public Health* 2015; 4: 863-65.
3. Lippi G, Montagnana M, Salvagno GL, Guidi GC. Should women with abnormal serum thyroid stimulating hormone undergo screening for anaemia? *Arch Pathol Lab Med* 2008; 132(3): 321-22.
4. Pitt-Rivers RV, Trotter WR, Wolff EC, Wolff J. *The Thyroid Gland*. London: Butter Worths, 1964. pp.1. 234-82.
5. Akhter S, Nahar ZU, Parvin S, Alam A, Sharmin S, Arslan MI. Thyroid status in patients with low serum ferritin level. *Bangladesh J Med Biochem* 2012; 5(1): 5-11.
6. Golde DW, Bersch N, Choprea IJ, Cline MJ. Thyroid hormone stimulate erythropoiesis I vitro. *Br J Haematol* 1977; 37(2):173-77.
7. Das KC, Mukherjee M, Sarkar TK, Dash RJ, rastogi GK. Erythropoiesis and erythropoietin in hypo- and hyperthyroidism. *J Clin Endocrinol Metab* 1975; 40(2):211-20.
8. Dorgalaleh A, Mahmoodi M, Varmaghani B, Kiani Node F, Saeedi kia O, Alizadeh Sh et al. Effect of thyroid dysfunctions in blood cell count and red blood cell indice. *Iranian J Pedia Hematol Oncol* 2012; 3(2): 73-77.
9. Kubota K, Kurabayashi H, Tamura J, Kawada E, Tamura K, Nagashima K. Change in the thyroid function by use of deferoxamine in a patient with hemo-chromatosis complicated by hyper-thyroidism. *J Med* 1992; 23(1): 75-77.
10. Rad NR, Vakili M, Zaver-reza J, Rezaie S, Shirvani AR. The relationship between thyroid hormone levels and body iron stores in Iranian hypothyroidism patients. *Int J Med Lab* 2016; 3(3): 176-84.
11. Eftekhari MH, Keshvaraz SA, Jalali M, Saadat N, Eshiraghan MR. Thyroid hormone status in iron deficient adolescent girls. *Iran J Med Sc* 2003; 28(4): 161-65.
12. Granner DK. The diversity of endocrine system In: *Harper's Illustrated Biochemistry*, RK Murray, DK Granner, PA Mayes, VW Rodwell 26th edn. USA: McGraw Hill, 2003, 434-55.
13. Klausner RD, Roualt TA, Harford JB. Regulating the fate of mRNA: the control of cellular iron metabolism. *Cell* 1993; 72(1):19-28.
14. Goossen B, Hentze MW. Position is the critical determinant for function of iron- responsive elements as translational regulators. *Mol Cell Biol* 1992; 12:1959-66.
15. Mullner EW, Kuhn LC. A stem loop in the 3' untranslated region mediates iron dependent regulation of transferrin receptor mRNA stability in the cytoplasm. *Cell* 1988; 53(5): 815-25.
16. Tang CK, Chin J, Harford JB, Klausner RD, Roualt TA. Iron regulates the activity of the iron responsive element binding protein without changing its rate of synthesis or degradation. *J Biol Chem* 1992; 267 (34): 24466-70.