



CODEN [USA]: IAJPBB

ISSN: 2349-7750

**INDO AMERICAN JOURNAL OF  
PHARMACEUTICAL SCIENCES**<http://doi.org/10.5281/zenodo.1048471>Available online at: <http://www.iajps.com>

Research Article

**DETERMINATION OF PAZOPANIB HYDROCHLORIDE IN  
SOLID DOSAGE FORM BY RP-HPLC METHOD:  
DEVELOPMENT AND VALIDATION**Amareshwar Shabada<sup>1</sup>, Nikhat Fatima<sup>2</sup>, Humera Badar<sup>3</sup>, Syeda Saba Sultana<sup>4</sup><sup>1</sup>Assistant Professor, Department of Pharmaceutics, St.Mary's Pharmacy College,  
Deshmukhi (Village-Pochampally (Mandal), Yadadri (District), Telangana-508284<sup>2</sup>Department of Pharmaceutical Analysis & Quality Assurance, St.Mary's Pharmacy College,  
Deshmukhi (Village-508284), Pochampally (Mandal), Yadadri (District), Telangana.<sup>3</sup> Department of Pharmaceutical Analysis & Quality Assurance, St.Mary's Pharmacy College,  
Deshmukhi (Village-508284), Pochampally (Mandal), Yadadri (District), Telangana.<sup>4</sup> Department of Pharmaceutical Analysis & Quality Assurance, St.Mary's Pharmacy College,  
Deshmukhi (Village-508284), Pochampally (Mandal), Yadadri (District), Telangana.**Abstract:**

A simple, accurate, precise, sensitive and stability indicating RP-HPLC method has been developed for the determination of Pazopanib hydrochloride in bulk drug and pharmaceutical dosage form, in which separations are done using develosil C18, 5 $\mu$ m, 150  $\times$  4.6mm i.d. column at a flow rate of 1.0mL/min with an injection volume of 20 $\mu$ L. The beer's law was obeyed over the concentration range of 5 - 35 $\mu$ g/mL. The correlation coefficient was found to be 0.996 and it showed good linearity, reproducibility, precision in this concentration range. The aim of this paper was to develop and validate the stability indicating RP-HPLC method for the determination of Pazopanib hydrochloride in bulk and pharmaceutical dosage forms. The % recovery values were found to be within the limits, which showed that the method was accurate. The LOD and LOQ were calculated using statistical methods. The % RSD values were less than 2. The developed method was successfully applied for determination of Pazopanib hydrochloride in pharmaceutical dosage form. The results obtained are in good agreement with those obtained by using the standard method.

**Keywords:** Pazopanib hydrochloride, Develosil, Stability indicating, Method Development, Validation.**Corresponding Author:****Amareshwar Shabada,**

Assistant Professor,

Department of Pharmaceutics,

St.Mary's Pharmacy College, Deshmukhi (Village-508284),

Pochampally (Mandal), Yadadri (District), Telangana

Mobile No:- 9573262146

Email Id: [amar.pharma99@gmail.com](mailto:amar.pharma99@gmail.com)

QR code



Please cite this article in press as Amareshwar Shabada et al, *Determination of Pazopanib Hydrochloride in Solid Dosage Form by RP-HPLC Method: Development and Validation*, Indo Am. J. P. Sci, 2017; 4(11).

**INTRODUCTION:**

Pazopanib is presented as the hydrochloride salt, with the chemical name 5-[[4-[(2,3-dimethyl-2H-indazol-6-yl) methylamino] - 2 pyrimidinyl] amino] -2-methyl benzene sulfonamide mono hydrochloride. It has the molecular formula C<sub>21</sub>H<sub>23</sub>N<sub>7</sub>O<sub>2</sub>S•HCl and a molecular weight of 473.99. Pazopanib is a multi-tyrosine kinase inhibitor of vascular endothelial growth factor receptor (VEGFR)-1, VEGFR-2, VEGFR-3, platelet-derived growth factor receptor (PDGFR)- $\alpha$  and - $\beta$ , fibroblast growth factor receptor (FGFR) -1 and -3, cytokine receptor (Kit), interleukin-2 receptor inducible T-cell kinase (Itk), leukocyte-specific protein tyrosine kinase (Lck), and transmembrane glycoprotein receptor tyrosine kinase (c-Fms). In vitro, pazopanib inhibited ligand-induced autophosphorylation of VEGFR-2, Kit and PDGFR- $\alpha$  receptors 1-3. According to the literature survey 4-7 it was found that few analytical methods on HPLC, HPTLC, UPLC and UV Spectrophotometer were reported for the estimation of Pazopanib hydrochloride in bulk drug and formulations. Hence there is need to develop and validate an analytical method to estimate pazopanib hydrochloride the drug. The objective of the proposed method is to develop simple, accurate, precise, stability indicating RP-HPLC method for the estimation of pazopanib hydrochloride in

of Pazopanib hydrochloride in pharmaceutical dosage form.

**MATERIAL AND METHODS:***Reagents and Chemicals*

Pazopanib hydrochloride working standard was procured from Mylan Laboratories, Hyderabad, and Telangana. Pazopanib hydrochloride 400 mg tablets (Votrient®) were purchased from local pharmacy. Purified water was obtained from Millipore system. Acetonitrile (HPLC grade) and potassium dihydrogen orthophosphate were obtained from Loba Chem, Mumbai, and Sd fine-Chem Ltd, Mumbai, respectively. All other chemicals used in the analysis were AR grade.

*Instrumental and analytical conditions*

The HPLC analysis was performed using a HITACHI L2130 with D Elite 2000 Software with Isocratic with UV-Visible Detector (L-2400). Column used was Develosil C18, 5 $\mu$ m, 150 x 4.6 mm i.d. UV detection was performed at 268nm. The injection volume of sample was 20 $\mu$ L. An isocratic mobile phase containing Acetonitrile and 0.05M potassium dihydrogen orthophosphate buffer (30:70% v/v), at the pH 2.5 with O-phosphoric acid was carried out with the flow rate of 1.0mL/min. Column was maintained at room temperature.




Figure 1: Chemical structure of Pazopanib hydrochloride.




Figure 2: Standard Chromatogram.

### Experimental

#### Mobile Phase Preparation

Mixture of Acetonitrile and Phosphate buffer, pH 2.5 in the ratio of 30:70(% v/v) respectively was used.

#### Preparation of Standard Solution

Accurately weighed 50mg of Pazopanib hydrochlorideworking standard was transferred into 50mL volumetric flask. About 40mL of diluent (mobile phase) was added and sonicated to dissolve. The volume was made up with diluent and mixed. 0.1mL of this solution was diluted to 10mL with mobile phase and mixed.

#### Preparation of Sample Solution

Four capsules were weighed accurately and the average weight was calculated.

Capsules were opened, fine powder was collected and equivalent to 50mg of Pazopanib hydrochloridesample was weighed and transferred into 50mL volumetric flask. 40mL of diluent was added and sonicated for 30min with intermediate shaking. Volume was made up with diluent. The above solution was centrifuged for 10min at 8000rpm. 0.1mL of this solution was diluted to 10mL with mobile phase and mixed. The solution was filtered through 0.45  $\mu$ m filter.




Figure 3: Sample Chromatogram.




Figure 4: UV spectrum of Pazopanib hydrochloride.

**Method Development**

A develosil C18, 5 $\mu$ m, 150 x 4.6 mm i.d as a stationary phase with a mobile phase of Acetonitrile and Phosphate buffer, pH 3.1 (25:75) at a flow rate 1.0mL/min and a detection wavelength of 268nm afforded the best separation of drug. The standard solution and sample solution prepared as above were injected into the 20 $\mu$ L loop and the chromatograms were recorded as shown in the "Fig. 2" and "Fig. 3" respectively. The retention time of drug, Pazopanib hydrochloride was found to be 2.67mins. The amount of drug present in sample was calculated.

**Method Validation**

The proposed stability indicating method has been developed and validated for the determination of Pazopanib hydrochloride in pharmaceutical dosage forms. According to International Conference on Harmonization (ICH) guidelines<sup>13-14</sup>, validation of the method was carried out by using specificity, accuracy, linearity, suitability, LOD, LOQ, precision and stability studies.

**System suitability**

A standard solution was prepared by using Pazopanib hydrochloride working standard as per test method and was injected 5 times into the HPLC system. The system suitability parameters were evaluated from the Resolution, USP tailing and USP plate count values obtained from standard chromatograms as shown in Table 1.

**Specificity**

Specificity was evaluated by injecting standard solution and placebo solution individually into HPLC system.

**Linearity of test solution**

A series of solutions are prepared from standard stock solution at concentration levels from 05-35 $\mu$ g/ml for Pazopanib hydrochloride.

**Accuracy**

Drug assay was performed in triplicate as per test method with equivalent amount of drugs into each volumetric flask for each spike level to get the concentration of drugs equivalent to 80%, 100% and 120% of the labeled amount as the test method.

**Precision****Repeatability**

Repeatability of method was evaluated by calculating the %RSD of peak areas of five replicate injections for the standard concentration (10ppm) of drug.

**Ruggedness**

The ruggedness was also evaluated by analyzing five samples of drug by two analysts in the same laboratory using different HPLC systems.

**Limit of Detection and Limit of Quantitation**

The parameters LOD and LOQ were determined on the basis of standard deviation and slope of the regression equation.

**Forced degradation studies**

The forced degradation study was performed to determine the specificity and stability indicating property of developed method. The drug was deliberately subjected to stress conditions such as acidic condition, alkaline condition, oxidation condition and thermal condition. All the solutions for degradation were prepared by dissolving drug in diluent to get an initial concentration of 1mg/mL and filtered. Acid decomposition was carried out in 1N hydrochloric acid and alkaline degradation was conducted using 1N sodium hydroxide and kept aside for 24 hours. Solutions for oxidative degradation were prepared using 3% hydrogen peroxide at a concentration of 1mg/mL of Pazopanib hydrochloride and kept aside for 24 hours. For thermal degradation study, the drug solution 1mg/mL was heated in calibrated oven at 80°C for 8 hours, cooled and used. These solutions are injected into the HPLC system and values were noted.




Figure 5: Linearity plot of Pazopanib hydrochloride.

Table 1: Data of System Suitability Parameter.

S.No.	Parameter	Limit	Result
1	Resolution	$R_s > 2$	9.15
2	USP Tailing	$T \leq 2$	0.12
3	USP plate count	$N > 2000$	3246

Table 2: Result of different parameters.

Parameters	Pazopanib hydrochloride
Specificity	Specific
Linearity	Regression equation, $y = 77391x$
	Correlation coefficient (r)
	0.999
Accuracy (recovery) n=3	Level I (80%)
	99.76%
	Level II (100%)
	100.54%
	Level III (120%)
	98.91%
Precision (%RSD) n=5	0.76
Ruggedness (%RSD) n=5	0.56
Limit of Detection (LOD)	0.341 $\mu\text{g/mL}$
Limit of Quantitation (LOQ)	1.023 $\mu\text{g/mL}$
System Suitability	USP Plate Count
	3246
	USP Tailing
	0.12

## RESULTS AND DISCUSSION:

### *Selection of detection wavelength*

From the UV spectrum, suitable wavelength considered for monitoring the drug was 268nm as shown in "Fig. 4". A stability indicating RP-HPLC method was developed by using an Develosil C18 (150mm  $\times$  4.6mm, 5 $\mu\text{m}$  particle size) as a stationary phase with a mobile phase of Acetonitrile and Phosphate buffer, pH 2.5 (30:70) at a flow rate 1.0mL/min and a detection wavelength of 268nm afforded the best separation of drug. The injection volume is 20 $\mu\text{L}$  and retention time for Pazopanib hydrochloride is 2.67mins. The method was validated according to ICH guidelines for various parameters like accuracy, precision, linearity, specificity, ruggedness, LOD, LOQ and stability studies. Linearity was obtained in the concentration range of 05-35 $\mu\text{g/mL}$  with correlation coefficient (r) of 0.999. The %recovery was found to be 99.74%. The %RSD for precision was found to be 0.76 and for ruggedness it was found to be 0.53.

## CONCLUSION:

A simple, precise, accurate, rapid, economical, stability indicating RP-HPLC method for estimation

of Pazopanib hydrochloride has been developed and validated as per ICH guidelines. The proposed method shows good agreement with all validation parameters. The optimized method is precise, accurate, specific, rugged, and a linear relation is observed between the concentration and the result. The developed method can be used for the analysis of routine quality control sample.

## REFERENCES:

- Lacy, Charles F, Armstrong, Lora L, Goldman, Morton P, Lance, Leonard L. Lexi-Comp's Drug Information Handbook (12th Edition). Lexi-Comp Inc. ISBN 1-59195-083-X, 2004.
- [http://www.drugbank.ca/system/fda\\_labels/DB00619.pdf?1265922812](http://www.drugbank.ca/system/fda_labels/DB00619.pdf?1265922812)
- [http://en.wikipedia.org/wiki/Pazopanib\\_hydrochlorideinib](http://en.wikipedia.org/wiki/Pazopanib_hydrochlorideinib)
- Gorja Ashok, Sumanta Mondal, S. Ganapaty and Jahnavi Bandla. Development and validation of stability indicating method for the Estimation of pazopanib hydrochloride in pharmaceutical dosage forms By RP-HPLC. Der Pharmacia Lettre, 2015; 7 (12): 234-241.

5. K. Vinodh Kumar, K. Puspha and D. Gowri Sankar. Development and validation of stability indicating RP-HPLC method for the determination of pazopanib hydrochloride in bulk drug and its Pharmaceutical dosage form. *Journal of Chemical and Pharmaceutical Research*, 2015; 7(8):114-120.
6. Kishore Kumar Reddy Y., J. Raveendra Reddy and N. Devanna. Development and validation of a stability indicating RP-HPLC method for the determination of Pazopanib in bulk drug and its pharmaceutical dosage form. *Journal of Chemical and Pharmaceutical Research*, 2015; 7(9):127-132.
7. Ravi Pratap Pulla, Asadulla Khan, J. Venkateswara Rao, Suresh Kumar Sudam; K. Sujana. *International Journal of Advances in Pharmaceutical Analysis*. 2013; 3, 1, 24-29.
8. ICH, Q1A(R2) Stability Testing of New Drug Substances and Products (Geneva, Feb. 2003).
9. ICH, Q1B Stability Testing: Photostability Testing of New Drug Substances and Products (Geneva, Nov. 1996).
10. ICH, Q6A: Specifications: Test Procedures and Acceptance Criteria for New Drug Substances and New Drug Products: Chemical Substances (Geneva, Oct. 1999).
11. ICH, Q3A(R2) Impurities in New Drug Substances (Geneva, Oct. 2006).
12. ICH, Q3B(R2) Impurities in New Drug Products (Geneva, June 2006).
13. ICH, Validation of Analytical Procedures: Text and methodology Q2 (R1): International Conference on Harmonization, ICH, Geneva (2005).
14. United States Pharmacopoeia (USP), XXVI. Validation of compendial methods. United States Pharmacopoeial Convention Inc.; Rockville, MD, USA: 2003.

