

# Antimicrobial activity of the rhizome essential oil of *Zingiber officinale* Roscoe

Hassan AM, Abutalib AA, Almagboul AZ and Kabbashi AS\*

Medicinal and Aromatic Plants and Traditional Medicine Research Institute (MAPTMRI), P.O. Box 2404, National Center for Research, Khartoum, Sudan.

Accepted 6 March, 2017

---

## ABSTRACT

The antimicrobial activity of the rhizome essential oil of *Zingiber officinale* (family: Zingiberaceae) was tested against six standard bacteria: *Bacillus subtilis*, *Staphylococcus aureus*, *Escherichia coli*, *Klebsiella pneumoniae*, *Proteus vulgaris*, *Pseudomonas aeruginosa* and against two standard fungi namely *Aspergillus niger* and *Candida albicans* using the cup plate agar diffusion method. The rhizome essential oil of *Z. officinale* dissolved in methanol (1:10), showed high activity against the Gram positive *S. aureus* (23 mm), *B. subtilis* (20 mm), Gram negative *E. coli* (21 mm), *P. vulgaris* (20 mm) and *P. aeruginosa* (27 mm). It also showed moderate activity against the Gram negative *K. pneumoniae* (15 mm), *C. albicans* (16 mm) and no activity against *Aspergillus niger*. The rhizome oil was also tested against sixty clinical isolates, collected randomly from Khartoum and Soba Hospitals. The minimum inhibitory concentrations (MICs) of the essential oil against standard bacteria were determined using the agar diffusion method. The antimicrobial activity of the reference drugs were determined against the standard organisms and compared with the antimicrobial activity of the tested oil. The obtained results showed variable sensitivity against the organisms tested and confirmed its folkloric uses.

**Keywords:** *Zingiber officinale* (Rhizome), essential oil, antimicrobial activity.

---

\*Corresponding author. e-mail: ahmedsak88@gmail.com.

---

## INTRODUCTION

Ginger (*Zingiber officinale*), a member of the Zingiberaceae family, is a well-known spice used in the daily diet in many Asian countries (Demin and Yingying, 2010). It is a rhizomatous plant grown throughout Southeastern Asia, China in parts of Japan, Austria, Latin America, Jamaica and Africa. It has been used as a spice and medicine in India and China since ancient times. It was known in Germany and France in the ninth century and in England in 10th century for its medicinal properties (Sasidharan and Nirmala, 2010). Over three quarters of the world population still rely on plants and plant extracts for health care (Sasidharan and Nirmala, 2010). Ginger compounds are active against specific type of diarrhea which is leading to cause death in infant in developing countries. Moreover, it has been found that ginger is effective in treating nausea caused by sea sickness, morning sickness and chemotherapy, though it was found superior over a place for post operative nausea (Demin

and Yingying, 2010; Sebiomo et al., 2011).

In addition, it has been reported that the main ingredients of ginger like volatile oil, gingerol, shogaol and diarylheptanoids work as antioxidant, anti-inflammatory, anti-lipid, anti-diabetic, analgesic, antipyretic and anti-tumor (Demin and Yingying, 2010; Sasidharan and Nirmala, 2010; Sebiomo et al., 2011; Lee et al., 1986; Penna et al., 2003; Islam and Choi, 2008; Isa et al., 2008; Wang et al., 2009; Shim et al., 2011).

In the Tibb and Ayurvedic systems of medicine preventive and ameliorative effects of ginger have been described in the treatment of catarrh, rheumatism, nervous diseases, gingivitis, toothache, asthma, stroke, constipation and diabetes (Chrubasik et al., 2005; Ali et al., 2008). Extracts were reported to have effects as an anti-inflammatory (Chang et al., 1995). It is considered safe herbal medicine with only few and insignificant adverse side effects (Ali et al., 2008). The aim of this

study was to determine the antimicrobial activity of the essential oil of ginger against standard bacteria, fungi and to determine its minimum inhibitory concentrations.

## MATERIALS AND METHODS

### Plant materials

The rhizome of *Zingiber officinale* were purchased from Alyhya Mole in October 2013 and it was identified and authenticated by the taxonomist Dr. Haider Abd alGader, Medicinal and Aromatic Plants and Traditional Medicine Research Institute (MAPTMRI), Khartoum, Sudan.

### Method of extraction

The oil of the tested *Z. officinale* leaves was obtained by hydrodistillation technique using Clevenger's apparatus. Hundred grams from plant materials were placed in a two liters round bottom flask and distilled water was added and mixed thoroughly. The contents of the flask were boiled gently for four hours until the volatile oil has been distilled. The crude volatile oil of plant was transferred by means of a pipette into a separate brown glass bottle. Anhydrous sodium sulphate was added agitated gently to absorb the water and the clear oil was decanted into brown glass bottle and kept in the refrigerator until needed for analysis.

### Test microorganisms

The oily solution of *Z. officinale* was tested against two Gram positive bacteria *Bacillus subtilis* (NCTC 8236) and *Staphylococcus aureus* (ATCC 25923). Four Gram negative organisms, *Escherichia coli* (ATCC 25922), *Klebsiella pneumoniae* (ATCC 53657), *Proteus vulgaris* (ATCC 6380), *Pseudomonas aeruginosa* (ATCC 27853) and two standard fungi, *Aspergillus niger* (ATCC 9763) and *Candida albicans* (ATCC7596). The tested organisms were obtained from the Department of Microbiology, MAP TMRI and National Health Laboratory, Khartoum, Sudan.

60 clinical isolates of *Bacillus subtilis*, *Staphylococcus aureus*, *Escherichia coli*, *Klebsiella pneumoniae*, *Proteus vulgaris* and *Pseudomonas aeruginosa* were collected from Khartoum and Soba Hospitals.

The bacterial cultures were maintained on nutrient agar and incubated at 37°C for 18 h and then used for tests.

### *In vitro* testing the oil *Zingiber officinale* for antimicrobial activity

The cup-plate agar diffusion method (Kavanagh, 1972) was adopted with some minor modifications to assess the antibacterial activity of the prepared extracts. One ml of the standardized bacterial stock suspension  $10^8$  to  $10^9$  C.F.U/ml were thoroughly mixed with 100 ml of molten sterile nutrient agar which was maintained at 45°C. 20 ml aliquots of the inoculated nutrient agar were distributed into sterile Petri-dishes. The agars were left to set and in each of these plates 4 cups (10 mm in diameter) were cut using a sterile cork borer (No. 4) and agar discs were removed. Alternate cups were filled with 0.1 ml sample of the oil using automatic microlitre pipette, and allowed to diffuse at room temperature for two hours. The plates were then incubated in the upright position at 37°C for 18 h. Two replicates were carried out for the oil against each of the tested organisms. After incubation the diameters of the resultant growth inhibition zones were measured, averaged and the mean values were

tabulated. Positive control involving the addition of the solvent (methanol) instead of the oil was carried out.

### Testing for antifungal activity

The same method as for bacteria was adopted. Instead of Nutrient agar, Sabouraud dextrose agar was used the inoculated medium was incubated at 25°C for two days for *Candida albicans* and three days for *Aspergillus niger*.

## RESULTS AND DISCUSSION

The rhizome of ginger (*Z. officinale*) a member of the Zingiberaceae family, was screened for its *in vitro* antimicrobial activity against two standard Gram positive bacteria *Bacillus subtilis*, *Staphylococcus aureus*, four standard Gram negative bacteria *Escherichia coli*, *Klebsiella pneumoniae*, *Proteus vulgaris* and *Pseudomonas aeruginosa* and two standard fungi *Aspergillus niger* and *Candida albicans* using a cup plate agar diffusion method. Table 1 and Figure 1 represents the results of antimicrobial activity of the methanolic solution prepared from the rhizome of ginger which showed high activity against *B. subtilis* (20 mm), *S. aureus* (23 mm), *E. coli* (21 mm), *P. vulgaris* (20 mm), and *P. aeruginosa* (27 mm) and moderate activity against *K. pneumoniae* (15 mm) and *C. albicans* (16 mm). It showed no activity against *A. niger*. Similarly Yin and Cheng (1988), Belik (2014) showed that ginger had no significant action on some fungi (*A. niger* and *A. flavus*). Jagetia et al. (2003) and Badreldein et al. (2008) found similar to our result that the ginger extract exhibited antimicrobial activity against *P. aeruginosa*, *E. coli* and *C. albicans*. The methanol extract of ginger showed the highest activity against *P. aeruginosa*, followed by *S. aureus*, *E. coli*, *B. subtilis*, *P. vulgaris* and *K. pneumoniae* was the least. Therefore methanol extract was more effective against the Gram negative organism *P. aeruginosa* than the Gram positive bacteria. This result was different from that reported by Hasan et al. (2012); Kaushik and Goyal (2011) who found that the methanol extract was more effective against the Gram positive bacteria compared to Gram negative bacteria.

Furthermore the Gram positive bacteria *S. aureus* was found more susceptible than the Gram negative bacteria *E. coli*, *K. pneumoniae* and *P. vulgaris* which is in agreement with the study of Chen et al. (1985) and Belik (2014). This is probably due to the difference in chemical composition and structure of cell wall of both types of microorganisms.


Therefore ginger inhibits the growth of all Gram positive, Gram negative and fungus *C. albicans*. This result is similar to that reported by Nielsen and Rios (2000) and Belik (2014). Gupta and Sharma (2014) found similar to our results that ginger inhibited the growth of *E. coli*, *Pr. Species*, *S. species*, *Streptococci* and *Salmonella*. Shahnaz and Mohammed, (2015) found

**Table 1.** *In vitro* antimicrobial activity of the essential oil of *Zingiber officinale* rhizome.

Family/plant name/vernacular name	Part used	Solvent system	Yield (%)	Tested organisms used*/ MIZD (mm)							
				B.s	S.a	E.c	K.p	Pr.v	Ps.a	A.n	C.a
Zingiberaceae (Zingiber officinale) Genzabil	Rhizome	MeOH	0.6%	20	23	21	15	20	27	-	16

Key: Bacteria\*: *B.s* = *Bacillus subtilis*, *S.a* = *Staphylococcus aureus*, *E.c* = *Escherichia coli*, *K.p* = *Klebsiella pneumoniae*, *Pr.v* = *Proteus vulgaris*, *Ps.a* = *Pseudomonas aeruginosa*, Fungi\*\*: *A.n* = *Aspergillus niger* and *C.a* = *Candida albicans*. Concentration of oil dissolved in methanol (1:10) at 0.1 ml/cup. MIZD\*(mm): Mean of Inhibition Zone Diameter in mm.

Interpretation of result: MIZD\* (mm): >14 mm: Resistant.<18 mm: Sensitive. , 14 -18 mm: Moderate.

**Figure 1.** *In vitro* antimicrobial activity of the essential oil of *Zingiber officinale* rhizome.**Table 2.** The activity of *Zingiber officinale* oil against different clinical isolates.


Microorganisms	No. of isolate	No. of clinical isolates		
		Sensitive	Moderate	Resistant
<i>B. subtilis</i>	10	7	3	-
<i>S. aureus</i>	10	10	-	-
<i>E. coli</i>	10	6	4	-
<i>K. pneumoniae</i>	10	-	9	-
<i>P. vulgaris</i>	10	3	7	-
<i>P. aeruginosa</i>	10	-	2	8

similar to our results that the oil showed significant antimicrobial activity against *E. coli*, *S. aureus*, *B. subtilis*, *C. albicans* but contrary to our result the oil was also active against *A. niger*. There are several reports on the inhibitory effect of ginger on the growth of *A. niger* (Singh et al., 2008; Sasidharan and Nirmala, 2010; Tagoe et al., 2011).

Comparison of observation given in Tables 1, 2 and 3, and Figure 1, 2 and 3 showed that the methanol extract of the rhizome of ginger inhibited *B. subtilis* and *P. aeruginosa* similar its 10 µg/ml Gentamicin, *S. aureus* more than 20 µg/ml and less than 40 µg/ml Ampicillin. It also inhibited *E. coli* more than 10 µg/ml Gentamicin, *K. pneumoniae* and *P. aeruginosa* less than 10 µg/ml

**Table 3.** Antibacterial and antifungal activity of reference drugs against standard microorganisms.


Drugs	Concentrations (µg/ml)	Standard microorganisms used MDIZ* (mm)					
		Tested bacteria used					
		<i>B.s</i>	<i>S.a</i>	<i>E.c</i>	<i>K.pn</i>	<i>Pr.v</i>	<i>Ps.a</i>
Ampicillin	40	15	25	-	35	16	16
	20	14	20	-	26	-	13
	10	13	18	-	25	-	12
	5	12	15	-	21	-	-
Gentamicin	40	29	35	32	26	25	23
	20	22	33	30	24	24	22
	10	20	30	17	21	23	21
	5	17	28	-	20	22	19
Clotrimazole			<i>A.n</i>			<i>C.a</i>	
	40		30			42	
	20		22			40	
	10		19			33	
Nystatin	5		16			30	
	50		28			17	
	25		26			14	
	12.5		23			-	



**Figure 2.** Minimum inhibitory concentrations of *Zingiber officinale* essential oil against standard microorganisms.

Gentamicin. It also inhibited *C. albicans* less than 50 µg/ml Nystatin. The methanolic extract of rhizome of

ginger was subjected to MIC by the cup plate agar diffusion method against bacterial species (Table 4 and



**Figure 3.** Antibacterial and antifungal activity of reference drugs against standard microorganisms.

**Table 4.** Minimum inhibitory concentrations of *Zingiber officinale* essential oil against standard microorganisms.

Concentration (mg/ml)	Bacteria						Fungi	
	B.s	S.a	E.c	K.p	Pr.v	Ps.a	A.n	C.a
100	20	23	21	15	20	27	-	16
50	18	21	19	14	18	23	-	15
25	15	20	15	13	17	20	-	14
12.5	14	18	14	12	16	18	-	13

Figure 2). The results indicated that the methanol extract inhibited the organisms less than 12.5 µg/ml. Control experiments using methanol for extract preparation (that is, negative control) showed no inhibition of any bacteria indicating that ginger itself and not the solvent inhibited the growth of Gram positive, Gram negative. Ampicillin, Gentamicin, Clotrimazole and Nystatin (positive controls) showed variable inhibition diameters ranging from 12 to 35 mm against Gram positive Gram negative bacteria and from 14 to 42 mm against Clotrimazole and Nystatin.

The results of this study showed that *Z. officinale* possessed antimicrobial properties as antibiotics and antifungal and may be used to manage some infectious disease caused by the tested organisms. The high activity of ginger essential oil may be due to the presence of phenolic compounds as it is well known that ginger essential oil contains considerable amount of phenolic compounds (eugenol, shogaols, zingerone, gingerol etc (Pawar et al., 2011, Din et al., 2012).

## CONCLUSION

The essential oil of *Z. officinale* showed various degree of inhibitory activity against the microorganisms tested. The obtained result may justify the use of the Sudanese *Z. officinale* as antimicrobial therapy in folkloric medicine in Sudan and the neighboring countries.

## ACKNOWLEDGEMENTS

The authors are grateful to Medicinal and Aromatic Plants and Traditional Medicine Research Institute (MAPTMRI), National Center for Research, Khartoum, Sudan.

## REFERENCES

Ali BH, Blunden G, Tanira MO, Nemmar A, 2008. Some phytochemical, pharmacological and toxicological properties of ginger (*Zingiber*

- officinale* Roscoe): a review of recent research. *Food. Chem Toxicol*, 46(2): 409-420.
- Belik Y, 2014.** Total antioxidant and antimicrobial potency of essential oil and oleoresin of *Zingiber officinale* Roscoe. *Asian Pac J Trop Dis*, 4(1): 40-44.
- Chang CP, Chang JY, Wang FY, Chang JG, 1995.** The effect of Chinese medicinal herb *Zingiberis* rhizome extract on cytokine secretion by human peripheral blood mononuclear cells. *J Ethnopharmacol*, 48(1): 13-19.
- Chen HC, Chang MD, Chang TJ, 1985.** Antibacterial properties of some spice plants before and after heat treatment. *Chin J Microbiol Immunol*, 18: 190-195.
- Chrubasik S, Pittler MH, Roufogalis BD, 2005.** *Zingiberis rhizoma* a comprehensive review on the effect and efficacy profiles. *Phytomedicine*, 12(19): 684-701.
- Demin, G., Yingying, Z. 2010.** Comparative antibacterial activities of crudopolysaccharides and flavonoids from *Zingiber officinale* and their extraction. *Am J Trop Med*, 5: 235-238.
- Din SH, An KJ, Zhao CP, Li Y, Guo YH, Wang ZF, 2012.** Effect of drying methods on volatiles of Chinese ginger (*Zingiber officinale* Roscoe). *Food Bioprod Process*, 3: 515-524.
- Gupta S, Sharma A, 2014.** Medicinal properties of *Zingiber officinale* Roscoe: A review. *IOSR J Pharm Biol Sci*, 9: 124-129.
- Hasan HA, Rasheed RAM, Abd Razik BM, Rasool HBA, 2012.** Chemical composition and antimicrobial activity of the crude extracts isolated from *Zingiber officinale* by different solvents. *Pharmaceut Anal Acta*, 3: 184.
- Isa Y, Miyakawa Y, Yanagisawa M, Goto T, Kang MS, Kawada T, Morimitsu Y, Kubota K, Tsuda T, 2008.** 6-Shogaol and 6-gingerol, the pungent of ginger, inhibit TNF- $\alpha$  mediated down regulation of adiponectin expression via different mechanisms in 3T3-L1 adipocytes. *Biochem Biophys Res Commun*, 373: 429-434.
- Islam MS, Choi H, 2008.** Comparative effects of dietary ginger (*Zingiber officinale*) and garlic (*Allium sativum*) investigated in a type 2 diabetes model of rats. *J Med Food*, 11: 152-159.
- Jagetia GC, Baliga MS, Venkatesh P, Ulloor JN, 2003.** Influence of ginger rhizome (*Zingiber officinale* Rosc.) on survival, glutathione and lipid peroxidation in mice after whole-body exposure to gamma radiation. *Radiat Res*, 160: 584-592.
- Kaushik P, Goyal P, 2011.** Evaluation of various crude extracts of *Zingiber officinale* rhizome for potential antibacterial activity: A study *in vitro*. *Adv Microbiol*, 1: 7-12.
- Kavanagh F, 1972.** Analytical Microbiology. Vol II Academic press (Pub) New York and London, pp 11.
- Lee YB, Kim YS, Ahmore CR, 1986.** Antioxidant property in ginger rhizoma and its application to mate products. *J Food Sci*, 51: 20-23.
- Nielsen PV, Rios R, 2000.** Inhibition of fungal growth on bread by volatile compounds from species herbs and mustard essential oil. *Int J Food Microbiol*, 60(2-3): 219-229.
- Pawar N, Pai S, Nimbalkar M, Dixit G, 2011.** RP-HPLC analysis of the phenolic antioxidant compound 6-gingerol from different ginger cultivars. *Food Chem*, 3: 281-287.
- Penna SC, Medeiros MV, Aimbire FS, Faria-Neto HC, Sertié JA, Lopes-Martins RA, 2003.** Anti-inflammatory effect of the hydroalcoholic extract of *Zingiber officinale* rhizomes on rat paws and skin edema. *Phytomedicine*, 10: 381-385.
- Sasidharan I, Nirmala MA, 2010.** Comparative chemical composition and antimicrobial activity fresh and dry ginger oils (*Zigiber Officinale* Roscoe). *Int J Curr Pharmaceut Res*, 2: 40-43.
- Sebiomo A, Awofodu AD, Awosanya AO, Awotona FE, Ajayi AJ, 2011.** Comparative studies of antibacterial effect of some antibiotics and ginger (*Zingiber officinale*) on two pathogenic bacteria. *J Microbiol Antimicrob*, 3: 18-22.
- Shahnaz S, Mohammed A, 2015.** Chemical composition of volatile oil of rhizome of *Zingiber officinale* Roscoe and its antimicrobial activity. *World J Pharmaceut Sci*, 4: 741-752.
- Shim S, Kim S, Choi DS, Kwon YB, Kwon J, 2011.** Anti-inflammatory effects of [6]-shogaol: Potential roles of HDAC inhibition and HSP70 induction. *Food Chem Toxicol*, 49: 2734-2740.
- Singh G, Kapoor IP, Singh P, de Heluani CS, de Lampasona MP, Catalan CAN, 2008.** Chemistry, antioxidant and antimicrobial investigations on essential oil and oleoresins of *Zingiber officinale*. *Food Chem Toxicol*, 46: 3295-3302.
- Tagoe DN, Nyarko HD, Akpaka RA, 2011.** A comparison of antifungal properties of onion (*Allium sativum*) against *Aspergillus flavus*, *Aspergillus niger* and *Cladosporium herbarum*. *Res J Med Plant*, 5(3): 281-287.
- Wang W, Li CY, Wen XD, Li P, Qi LW, 2009.** Simultaneous determination of 6-gingerol, 8-gingerol, 10-gingerol and 6-shogaol in rat plasma by liquid chromatography-mass spectrometry: Application to pharmacokinetics. *J Chromatogr B Analyt Technol Biomed Life Sci B*, 877: 671-679.
- Yin MC, Cheng WS, 1988.** Inhibition of *Aspergillus niger* and *Aspergillus flavus* by some herbs and spices. *J Food Protect*, 61: 123-125.

---

**Citation:** Hassan AM, Abutalib AA, Almagboul AZ, Kabbashi AS, 2017. Antimicrobial activity of the rhizome essential oil of *Zingiber officinale* Roscoe. *Adv Med Plant Res*, 5(1): 5-10.

---