

Superior intestinal integrity and limited microbial translocation are associated with lower immune activation in SIVmac239-infected northern pig-tailed macaques (*Macaca leonina*)

Ming-Xu Zhang^{1,2,#}, Tian-Zhang Song^{1,2,#}, Hong-Yi Zheng¹, Xue-Hui Wang^{1,3}, Ying Lu^{1,2}, Han-Dan Zhang¹, Ting Li¹, Wei Pang¹, Yong-Tang Zheng^{1,4,*}

¹ Key Laboratory of Animal Models and Human Disease Mechanisms of the Chinese Academy of Sciences/Key Laboratory of Bioactive Peptides of Yunnan Province, National Kunming High Level Biosafety Research Center for Non-human Primates, KIZ-CUHK Joint Laboratory of Bioresources and Molecular Research in Common Diseases, Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming Yunnan 650223, China

² Kunming College of Life Science, University of Chinese Academy of Sciences, Kunming Yunnan 650204, China

³ School of Life Sciences, University of Science and Technology of China, Hefei Anhui 230026, China

⁴ KIZ-SU Joint Laboratory of Animal Models and Drug Development, College of Pharmaceutical Sciences, Soochow University, Suzhou Jiangsu 215123, China

ABSTRACT

Microbial translocation is a cause of systemic immune activation in HIV/SIV infection. In the present study, we found a lower CD8⁺ T cell activation level in *Macaca leonina* (northern pig-tailed macaques, NPMs) than in *Macaca mulatta* (Chinese rhesus macaques, ChRMs) during SIVmac239 infection. Furthermore, the levels of plasma LPS-binding protein and soluble CD14 in NPMs were lower than those in ChRMs. Compared with ChRMs, SIV-infected NPMs had lower Chiu scores, representing relatively normal intestinal mucosa. In addition, no obvious damage to the ileum or colon epithelial barrier was observed in either infected or uninfected NPMs, which differed to that found in ChRMs. Furthermore, no significant

microbial translocation (*Escherichia coli*) was detected in the colon or ileum of infected or uninfected NPMs, which again differed to that observed in ChRMs. In conclusion, NPMs retained superior intestinal integrity and limited microbial translocation during SIV infection, which may contribute to their lower immune activation compared with ChRMs.

Keywords: Immune activation; Intestinal integrity; Microbial translocation; SIVmac239; *Macaca leonina*; Northern pig-tailed macaques; *Macaca mulatta*; Chinese rhesus macaques

Received: 13 December 2018; Accepted: 07 March 2019; Online: 10 April 2019

Foundation items: This work was partly supported by grants from the National Natural Science Foundation of China (U1802284; 81471620, 81671627, 81771770, 81571606), 13th Five-Year Key Scientific and Technological Program of China (2017ZX10304402-002-004, 2017ZX10202102-001-005, 2018ZX10301101-002-003, 2018ZX10301406-003), and Knowledge Innovation Program of the Chinese Academy of Sciences (ZDRW-ZS-2016-4)

#Authors contributed equally to this work

*Corresponding author, E-mail: zhengyt@mail.kiz.ac.cn

DOI: 10.24272/j.issn.2095-8137.2019.047

Open Access

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright ©2019 Editorial Office of Zoological Research, Kunming Institute of Zoology, Chinese Academy of Sciences

INTRODUCTION

Immune activation is a hallmark of HIV/SIV infection and is closely correlated with disease progression (Boasso & Shearer, 2008; Hazenberg et al., 2003). In addition, T cell activation is more predictive of AIDS disease progression than plasma viral load and CD4⁺ T cell count (Giorgi et al., 1999; Liu et al., 1998). Immune activation may accelerate the depletion or impede the reconstitution of CD4⁺ T cells by exacerbating lymphoid tissue fibrosis (Estes et al., 2008; Hunt et al., 2003; Zeng et al., 2011). Furthermore, a number of non-AIDS diseases are also correlated with immune activation, such as cardiovascular disease and cancer (Cadogan & Dalgleish, 2008; Kaplan et al., 2011; Klatt et al., 2013). However, immune activation is difficult to alleviate, even in patients with sustained viral suppression during antiretroviral therapy (French et al., 2009). Therefore, therapeutic strategies to relieve persistent immune activation are needed for patients to improve their quality of life.

Microbial translocation is a significant factor that contributes to persistent immune activation (Brenchley et al., 2006; Sandler & Douek, 2012). HIV/SIV infection can cause severe mucosal CD4⁺ T cell depletion and damage to intestinal integrity, followed by translocation of microbial products into circulation, including 16S rDNA and lipopolysaccharide (LPS) (Brenchley et al., 2004; Dillon et al., 2012; Marchetti et al., 2013; Veazey et al., 1998). As a result, soluble CD14 (sCD14) released by activated monocytes and LPS-binding protein (LBP) levels increase (Brenchley et al., 2006). Plasma sCD14 levels can predict mortality in HIV infection (Sandler et al., 2011). In addition, elevated microbial translocation is associated with increased risk of AIDS-related lymphoma and HIV-associated dementia (Ancuta et al., 2008; Marks et al., 2013). Therefore, therapeutic strategies targeting microbial products and their downstream effects will be of great benefit.

Pig-tailed macaques are divided into three species based on morphological characteristics (Spartaco, 2001). Thereinto, *Macaca nemestrina* (southern pig-tailed macaques, SPMs) are suitable animal models to study the relationship between microbial translocation and disease progression of AIDS due to their compromised intestinal integrity and increased microbial translocation prior to SIV infection (Klatt et al., 2010). Our recent studies indicate, however, that pre-infection peripheral CD4⁺ T cell counts in *Macaca leonina* (northern pig-tailed macaques, NPMs) differ from those in *Macaca nemestrina* (Klatt et al., 2012; Zheng et al., 2014b). In addition, CD4⁺ T cells also exhibit superior homeostasis in SIVmac239-infected NPMs compared with Chinese rhesus macaques (ChRMs), which is not reported in SIV-infected SPMs (Zhang et al., 2017a). This raises the question as to whether compromised intestinal integrity and increased microbial translocation occur in pre- and post-infected NPMs similarly.

In the present study, we analyzed the levels of CD8⁺ T cell activation and plasma biomarkers of microbial translocation in SIVmac239-infected NPMs and ChRMs and normal NPMs. To further analyze the status of microbial translocation, intestinal

integrity and microbial translocation of NPMs were also studied.

MATERIALS AND METHODS

Animals and sample collection

Viral seroprevalence in NPMs and ChRMs enrolled in this study was investigated as described previously (Zhang et al., 2016b). All NPMs and ChRMs were healthy males and were seronegative for SIV, STLV, SRV, and SFV prior to the study. Six NPMs and three ChRMs (6–8 years old) were inoculated with a 3 000 50% tissue culture infectious dose (TCID₅₀) of SIVmac239 intravenously. Blood was collected periodically, and tissues were obtained from the macaques at necropsy. Tissue and blood samples from four uninfected NPMs were also obtained as the control. In addition, blood from five uninfected ChRMs and tissues from one SIVmac239-infected ChRM were also obtained. Blood was partially used for flow cytometry analyses and isolation of plasma and peripheral blood mononuclear cells (PBMCs). All tissues were partially preserved in 4% paraformaldehyde and –80 °C immediately after collection. Detailed information on animals used in histopathological assessment is shown in Table 1.

All animal experiments and procedures were approved by the Ethics Committee of the Kunming Institute of Zoology, Chinese Academy of Sciences (approval No.: SYDW-2015023;

Table 1 Animals enrolled in this study

Animal ID	Species	Status	Age	Survival days post infection
08247	NPM	Infected	7 ^a	491 ^b
08287	NPM	Infected	7 ^a	641 ^c
09203	NPM	Infected	6 ^a	648 ^c
09211	NPM	Infected	6 ^a	476 ^b
10205	NPM	Infected	5 ^a	626 ^c
10225	NPM	Infected	5 ^a	634 ^c
08091	ChRM	Infected	7 ^a	320 ^b
08303	ChRM	Infected	7 ^a	475 ^b
08439	ChRM	Infected	7 ^a	628 ^b
08051	ChRM	Infected	7 ^a	402 ^b
08269	NPM	Uninfected	9 ^{c, d}	–
09223	NPM	Uninfected	8 ^{c, d}	–
09237	NPM	Uninfected	8 ^{c, d}	–
09255	NPM	Uninfected	8 ^{c, d}	–
11049	ChRM	Uninfected	6 ^e	–
105353	ChRM	Uninfected	7 ^e	–
10039	ChRM	Uninfected	7 ^e	–
09003	ChRM	Uninfected	8 ^e	–
090017	ChRM	Uninfected	8 ^e	–

NPM: Northern pig-tailed macaque; ChRM: Chinese rhesus macaque.

^a: Age at infection. ^b: Natural death. ^c: Euthanasia. ^d: Age at necropsy.

^e: Blood collection only. – : Not available.

approval date: 20 June 2015).

Flow cytometry

Multiparameter flow cytometry was performed using peripheral blood, as described previously (Ma et al., 2012; Xia et al., 2009; Zhang et al., 2017b; Zheng et al., 2014a). For the frequency of HLA-DR⁺CD8⁺ T cells, blood was treated with lysing buffer (BD Biosciences, CA, USA) for 10 min and then incubated with a mixture of flow cytometry antibodies at 4 °C for 30 min. For staining with ki67, surface-labeled cells were further treated with fixation and permeabilization solution (BD Biosciences, CA, USA), followed by perm/wash buffer (BD Biosciences, CA, USA). Cross-reactive flow cytometry human antibodies anti-CD3 APC-Cy7 (clone SP34-2), anti-CD8 PE-Cy7 (clone RPA-T8), anti-HLA-DR APC (clone G46-6), and anti-ki67 PE (B56) were purchased from BD Pharmingen (Franklin Lakes, NJ, USA). Anti-CD4 PerCP-Cy5.5 (clone OKT4) was obtained from Biolegend (San Diego, CA, USA). Flow cytometry acquisition was performed on a BD FACSVersTM flow cytometer (BD, Franklin Lakes, NJ, USA) and flow cytometric data analysis was performed using FlowJo vX.0.7 (TreeStar, Ashland, OR, USA).

Enzyme-linked immuno sorbent assay (ELISA)

Plasma used for ELISA was isolated from blood by centrifugation (room temperature, 500 g, 10 min). Plasma was preserved at -80 °C until detection. Plasma sCD14, LBP and I-FABP levels were detected using commercially available ELISA kits obtained from R&D Systems (Minneapolis, MN, USA).

Histopathological assessment

Intestinal tissues fixed in 4% paraformaldehyde were dehydrated in a graded ethanol series and then embedded in paraffin. We collected 4 µm thick paraffin sections, which were stained with hematoxylin and eosin (H&E). Intestinal injury was evaluated using the Chiu score system following the grading principles described by Chiu et al. (1970).

Immunofluorescence

Immunofluorescence was performed as described previously (Zhang et al., 2016a). The 4 µm thick paraffin sections mentioned above were baked at 60 °C for 2 h before dewaxing in xylene and rehydrating through a graded ethanol series. The sections were then washed in flowing water for more than 15 min. Antigen retrieval was performed using the high-pressure method. The sections were then naturally cooled to room temperature and washed in flowing water for 5 min. For immunofluorescence, sections were treated with 1% sodium borohydride solution for 30 min and washed in phosphate buffered saline. After treatment with blocking buffer, sections were incubated with primary and secondary antibodies. The sections were then treated with Sudan Black for 20 min and washed with 50% ethanol. Mounting medium containing DAPI was used to adhere a coverglass to the microscope slide at last. Cross-reactive antibodies, including rabbit anti-claudin-3 antibody (polyclonal), mouse anti-cytokeratin antibody (clone MNF116), donkey anti-rabbit IgG

(Alexa Fluor[®]488, polyclonal), and donkey anti-mouse IgG (Alexa Fluor[®]555, polyclonal), were purchased from Abcam (Cambridgeshire, UK). Rabbit anti-*E. coli* antibody was purchased from Dako (Glostrup, Denmark). Microscopic examinations were performed on a Leica DMI4000B Microsystem (Leica Microsystems, Wetzlar, Germany). All slides were inspected, with photographs taken using the system.

Statistical analysis

Two-way ANOVA was used to compare frequencies of HLA-DR⁺CD8⁺ T cells, ki67⁺CD8⁺ T cells and plasma I-FABP levels between NPMs and ChRMs. The Mann-Whitney nonparametric test was used to compare pre-infection plasma sCD14, LBP and I-FABP levels and Chiu scores. Spearman's rank test was used to determine correlations between CD8⁺ T cell activation and plasma sCD14, LBP and I-FABP levels in NPMs and ChRMs during SIVmac239 infection. All data analyses were performed using GraphPad Prism v6.01 (GraphPad Software, San Diego, CA, USA).

RESULTS

NPMs have lower frequencies of activated and proliferating CD8⁺ T cells than ChRMs during SIVmac239 infection

T cell activation, especially that of CD8⁺ T cells, is more closely correlated with disease progression than plasma viral load or CD4⁺ T cell count during HIV-1 infection (Giorgi et al., 1999; Liu et al., 1998). In our previous study, we also found a close relationship between T cell activation and disease progression in SIVmac239-infected NPMs (Zhang et al., 2017b). Therefore, here we compared the frequencies of activated and proliferating CD8⁺ T cells in NPMs and ChRMs. The infection was divided into an acute and chronic phase at 12 weeks post infection (Figure 1). The frequencies of HLA-DR⁺CD8⁺ T cells and ki67⁺CD8⁺ T cells in NPMs were both lower than those in ChRMs, regardless of acute or chronic phase, which indicated that CD8⁺ T cell activation and proliferation levels of NPMs were lower than those of ChRMs during SIVmac239 infection (Figure 1A, B).

NPMs have lower levels of plasma sCD14 and LBP than ChRMs during SIVmac239 infection

As microbial translocation is an important cause of immune activation, we further analyzed the plasma biomarkers of microbial translocation, including sCD14 and LBP (Figure 2). We found no significant differences in pre-infection plasma sCD14 and LBP levels between NPMs and ChRMs (Figure 2A). These results implied that there was no obvious difference in microbial translocation between NPMs and ChRMs in the absence of SIV infection. Plasma sCD14 levels in NPMs were lower than in those ChRMs during SIVmac239 infection, which indicated that SIVmac239 infection might cause more severe microbial translocation in ChRMs than in NPMs (Figure 2B). However, there was no obvious difference in LBP levels between SIVmac239-infected NPMs and




Figure 1 Comparisons of CD8⁺ T cell activation and proliferation between NPMs and ChRMs

Frequencies of CD8⁺HLA-DR⁺ (A) and CD8⁺ki67⁺ (B) in peripheral blood T cells. Cubic spline curves were created to show the data trends. Infection was divided into acute phase and chronic phase at 12 weeks post infection. Two-way ANOVA was used to compare the frequencies of CD8⁺HLA-DR⁺ and CD8⁺ki67⁺ T cells between NPMs (black, $n=6$) and ChRMs (red, $n=3$). P -values represent differences between the two species. $P < 0.05$ was considered statistically significant.

ChRMs (Figure 2C). Furthermore, plasma sCD14 levels were directly correlated with frequencies of HLA-DR⁺CD8⁺ T cells in both NPMs and ChRMs, which implied that higher sCD14 levels might contribute to higher CD8⁺ T activation (Figure 2D). In contrast, plasma LBP levels were not directly correlated with the frequencies of HLA-DR⁺CD8⁺ T cells in either NPMs or ChRMs (Figure 2E). Thus, not all microbial translocation-related products appear to be closely associated with immune activation.

Intestinal mucosal injuries in ChRMs were more severe than in NPMs post SIVmac239 infection

Microbial translocation mainly occurs in the intestine; thus, we

focused on mucosal injuries in the colon and ileum. The normal NPMs had relatively intact intestinal mucosa, whereas the SIVmac239-infected ChRMs and NPMs presented varying degrees of damage (Figure 3A). We evaluated the severity of mucosal injury using the Chiu score system (Chiu et al., 1970), which indicated that colon mucosal injuries in SIVmac239-infected ChRMs were more severe than those in SIVmac239-infected NPMs, though there was no obvious difference in the Chiu score of the ileum mucosa (Figure 3B, C). In addition, colon and ileum mucosal injuries in SIVmac239-infected NPMs were more severe than those in the normal group. These results indicate that SIVmac239 infection can damage intestinal mucosa and NPMs may be




Figure 2 Plasma microbial translocation biomarker levels and their correlations with frequencies of CD8⁺HLA-DR⁺ T cells

A: Pre-infection plasma sCD14 and LBP levels of NPMs ($n=10$) and ChRMs ($n=8$). Mann-Whitney nonparametric test was used to calculate P -values. $P < 0.05$ was considered statistically significant. Two-way ANOVA was used compared sCD14 (B) and LBP (C) plasma levels during SIVmac239 infection. Plasma sCD14 and LBP levels in NPMs ($n=6$) and ChRMs ($n=3$) from the same post-infection time points were compared. $P < 0.05$ was considered statistically significant. Correlations between frequencies of CD8⁺HLA-DR⁺ T cells and sCD14 (D) and LBP (E) plasma levels in NPMs ($n=6$) and ChRMs ($n=3$) were analyzed. Spearman rank correlation coefficients were calculated to assess correlations. $P < 0.05$ was considered statistically significant.

less affected than ChRMs.

NPMs maintained superior intestinal epithelial integrity post SIVmac239 infection

Tight junctions between the intestinal epithelium play an important role in preventing translocation of gut microbe products from the lumen into the subepithelial region via the paracellular pathway (Milatz et al., 2010). Therefore, we detected the expression of tight junction protein claudin-3, which reflects the integrity of the intestinal epithelium. The intestinal epithelium in SIVmac239-infected ChRMs showed severe discontinuities in both the colon and ileum. In contrast, however, the intestinal epithelium of SIVmac239-infected NPMs showed superior continuities, as did normal NPMs

(Figure 4A). These results indicate that SIVmac239 infection may not severely affect the expression of tight junction protein claudin-3 in NPMs.

Limited microbial translocation levels in SIVmac239-infected NPMs

Although SIVmac239-infected NPMs maintained superior intestinal epithelial integrity, whether the relatively intact intestinal epithelium was concomitant with lower microbial translocation was unclear. Therefore, we selected *Escherichia coli* as a representative microbial product and analyzed its infiltration into the intestinal epithelium (represented by cytokeratin). As expected, translocation of *E. coli* through the intestinal epithelium was obvious in SIVmac239-infected




Figure 3 Intestinal mucosal injuries in the colon and ileum

A: Representative images (100×) of colon and ileum in normal NPMs (NPM Nor, $n=4$) and SIVmac239-infected ChRMs (ChRM SIV, $n=4$) and NPMs (NPM SIV, $n=6$). B: Colon Chiu scores. C: Ileum Chiu scores. Chiu scores range from 0 to 5, with a higher score indicating more severe injury. Mann-Whitney nonparametric test was used to compare Chiu scores between different groups. $P<0.05$ was considered statistically significant.




Figure 4 Tight junctions and translocation of microbial products (*E. coli*) into intestinal lamina propria in the colon and ileum

A: Representative images (100×) of epithelial barriers in the colon and ileum of normal NPMs (NPM Nor, $n=4$) and SIVmac239-infected NPMs (NPM SIV, $n=6$) and ChRMs (ChRM SIV, $n=4$). Tight junction protein claudin-3 (green) was stained to show the integrity of epithelial barriers. Blue is DAPI. White arrows show discontinuities, red arrows show continuities of epithelial barriers. B: Representative images (100×) of microbial translocation in the colon and ileum. Immunofluorescence was performed to stain DAPI (blue), *E. coli* (green), and cytokeratin (red). White arrows show translocation of microbial products (*E. coli*) into lamina propria.

ChRMs, but rare in SIVmac239-infected or normal NPMs (Figure 4B). These results are consistent with the expression of the tight junction protein claudin-3. In summary, NPMs had limited microbial translocation due to superior intestinal epithelial integrity.

DISCUSSION

Microbial translocation is associated with systemic immune activation during HIV/SIV infection, which further accelerates disease progression (Brenchley et al., 2006; Canary et al., 2013; Marchetti et al., 2011).

The pig-tailed macaque is an Old-World monkey known to be susceptible to HIV-1 infection (Agy et al., 1992; Liao et al., 2007). We previously identified a novel TRIM5-cyclophilin fusion gene (TRIMCyp) in the pig-tailed macaque that did not inhibit HIV-1 or SIV replication, thus explaining why the pig-tailed macaque is prone to HIV-1 infection (Kuang et al., 2009; Liao et al., 2007). Recently, three NPM virus models, i.e., HIV-

1_{NL4.3}, HIV-1_{R3A}, and stHIV-1sv, were established in our lab and a long-term viral reservoir was detected in these NPMs, which indicated that NPMs might serve as a potential model for HIV-1 reservoir research (Pang et al., 2017, 2018). In addition, we previously reported on various biological parameters of NPMs (Lian et al., 2016, 2017, 2018; Song et al., 2018; Zhang et al., 2014, 2016b, 2016c; Zheng et al., 2014b; Zhu et al., 2015), thus promoting the application of NPMs in AIDS and other biomedical research. SPMs are appropriate research models due to their compromised intestinal integrity and increased microbial translocation in the absence of SIV infection (Canary et al., 2013; Klatt et al., 2010). In the present study, however, we found lower levels of CD8⁺ T cell activation in SIVmac239-infected NPMs than in ChRMs, which was partly attributed to superior intestinal integrity and limited microbial translocation. These results imply that the intestinal microenvironment of NPMs may differ from that of SPMs. Thus, further study on the differences and potential mechanisms may provide new insight into the therapeutic strategies of microbial translocation during HIV infection.

Several plasma biomarkers are commonly selected to evaluate the level of microbial translocation, including LPS, sCD14, LBP, and EndoCAb (endotoxin-core antibody) (Brenchley et al., 2006; Marchetti et al., 2013). In contrast to uninfected individuals, HIV-infected progressors have significantly higher levels of plasma sCD14 and LBP and significantly lower levels of EndoCAb (Brenchley et al., 2006). In the present study, we also observed direct correlations between plasma sCD14 levels and CD8⁺ T cell activation in SIVmac239-infected NPMs and ChRMs, which implied that microbial translocation was closely associated with immune activation (Figure 2D). However, plasma biomarkers do not consistently reflect the real level of microbial translocation and limitations do exist (Marchetti et al., 2013). Here, we did not observe direct correlations between plasma LBP levels and CD8⁺ T cell activation in either NPMs or ChRMs (Figure 2E). Previous research has demonstrated that gut commensal microbes are not the main contributors to circulating CD4⁺ T cell activation (Zimmermann et al., 2015). Furthermore, CD8⁺ T cells may be activated in an antigen-independent manner that is not associated with the level of microbial translocation-related products (Bastidas et al., 2014). Therefore, the relationship between certain plasma microbial translocation biomarkers and immune activation needs further confirmation.

Histopathological assessment provides the most direct

evidence of intestinal mucosal injuries. In the present study, superior intestinal integrity was only coincident with lower plasma sCD14 levels in NPMs. These results indicate that compromised intestinal integrity may cause severe microbial translocation, followed by systemic immune activation, which accelerates disease progression. Furthermore, microbial translocation resulting from breakdown of intestinal mucosa may result in microbial product translocation into draining or distant lymph nodes, followed by persistent local immune activation (Estes et al., 2010). Therefore, intestinal mucosal damage appears to be the fundamental cause of microbial translocation and subsequent immune activation.

To assess the level of intestinal injury, fatty acid-binding proteins are often used as plasma biomarkers of tissue injury (Pelsers et al., 2003, 2005). In the present study, we selected intestinal-type fatty acid-binding protein (I-FABP) as the plasma biomarker of intestinal injury, as described elsewhere (Jenabian et al., 2015; Pelsers et al., 2003). As the plasma I-FABP levels in most individuals were below the lower limit, we did not find any significant differences in plasma I-FABP levels between NPMs and ChRMs either prior to or post infection (Figure 5A, B). Further study showed that plasma I-FABP levels were directly associated with CD8⁺ T cell activation in NPMs, but not in ChRMs (Figure 5C, D). In summary, although the plasma I-FABP level was correlated with immune




Figure 5 Plasma I-FABP levels and their correlations with CD8⁺ T cell activation

A: Pre-infection plasma I-FABP levels. Mann-Whitney nonparametric test was used to calculate *P*-values. *P*<0.05 was considered statistically significant. B: Plasma I-FABP levels during SIVmac239 infection. Two-way ANOVA compared the plasma I-FABP levels of NPMs and ChRMs. *P*< 0.05 was considered statistically significant. C, D: Correlations between CD8⁺ T cell activation and plasma I-FABP levels in NPMs and ChRMs during SIVmac239 infection. Spearman's rank test was used to determine correlations. *P*<0.05 was considered statistically significant.

activation to some extent, I-FABP may not fully reflect the level of intestinal injury or microbial translocation.

There are some limitations in the present study. First, it is difficult to obtain absolutely accurate results using such a small number of animals. Given the differences between each macaque, more experimental animals would improve accuracy. In addition, infection time varies between individuals, which may also affect pathological consequences.

In the present study, we characterized the relationship between immune activation and microbial translocation in SIVmac239-infected NPMs for the first time. Intriguingly, NPMs maintained superior intestinal integrity and limited microbial translocation after SIV infection, which was different from that observed in ChRMs and SPMs. How NPM intestinal mucosa avoids damage from SIV infection is still an enigma that needs to be resolved. Research on intestinal mucosal immunity in NPMs is likely to provide guidance for therapeutic strategies of microbial translocation in HIV-1-infected patients. In addition, although pig-tailed macaques are susceptible to HIV-1 infection, how HIV-1 affects regional immunity in the intestinal tract of pig-tailed macaques is still unclear (Kuang et al., 2009; Liao et al., 2007; Pang et al., 2017). Studies on this problem may provide deeper insight into the mechanisms of microbial translocation during HIV-1 infection.

COMPETING INTERESTS

The authors declare that they have no competing interests.

AUTHORS CONTRIBUTIONS

Y.T.Z. and M.X.Z. conceived and designed the experiments. M.X.Z., T.Z.S., H.Y.Z., X.H.W., Y.L., H.D.Z., T.L. and W.P. performed the experiments. M.X.Z. analyzed the data. M.X.Z. and Y.T.Z. wrote the paper. All authors read and approved the final version of the manuscript.

ACKNOWLEDGEMENTS

We thank Jin Jiang, Yu Xiao, Xiao-Dong Lian, Ren-Rong Tian, Jia-Hao Song, and Min Chen from the Kunming Institute of Zoology, Chinese Academy of Sciences, for their assistance with the experiments.

REFERENCES

Agy MB, Frumkin LR, Corey L, Coombs RW, Wolinsky SM, Koehler J, Morton WR, Katze MG. 1992. Infection of *Macaca nemestrina* by human immunodeficiency virus type-1. *Science*, **257**(5066): 103–106.

Ancuta P, Kamat A, Kunstman KJ, Kim EY, Autissier P, Wurcel A, Zaman T, Stone D, Mefford M, Morgello S, Singer EJ, Wolinsky SM, Gabuzda D. 2008. Microbial translocation is associated with increased monocyte activation and dementia in AIDS patients. *PLoS One*, **3**(6): e2516.

Bastidas S, Graw F, Smith MZ, Kuster H, Günthard HF, Oxenius A. 2014. CD8⁺ T cells are activated in an antigen-independent manner in HIV-infected individuals. *Journal of Immunology*, **192**(4): 1732–1744.

Boasso A, Shearer GM. 2008. Chronic innate immune activation as a cause of HIV-1 immunopathogenesis. *Clinical Immunology*, **126**(3): 235–242.

Brenchley JM, Price DA, Schacker TW, Asher TE, Silvestri G, Rao S, Kazzaz Z, Bornstein E, Lambotte O, Altmann D, Blazar BR, Rodriguez B, Teixeira-Johnson L, Landay A, Martin JN, Hecht FM, Picker LJ, Lederman MM, Deeks SG, Douek DC. 2006. Microbial translocation is a cause of systemic immune activation in chronic HIV infection. *Nature Medicine*, **12**(12): 1365–1371.

Brenchley JM, Schacker TW, Ruff LE, Price DA, Taylor JH, Beilman GJ, Nguyen PL, Khoruts A, Larson M, Haase AT, Douek DC. 2004. CD4(+) T cell depletion during all stages of HIV disease occurs predominantly in the gastrointestinal tract. *Journal of Experimental Medicine*, **200**(6): 749–759.

Cadogan M, Dalgleish AG. 2008. HIV induced AIDS and related cancers: chronic immune activation and future therapeutic strategies. *Advances in Cancer Research*, **101**: 349–395.

Canary LA, Vinton CL, Morcock DR, Pierce JB, Estes JD, Brenchley JM, Klatt NR. 2013. Rate of AIDS progression is associated with gastrointestinal dysfunction in simian immunodeficiency virus-infected pigtail macaques. *Journal of Immunology*, **190**(6): 2959–2965.

Chiu CJ, Mcardle AH, Brown R, Scott HJ, Gurd FN. 1970. Intestinal mucosal lesion in low-flow states. I. A morphological, hemodynamic, and metabolic reappraisal. *Archives of Surgery*, **101**(4): 478–483.

Dillon SM, Manuzak JA, Leone AK, Lee EJ, Rogers LM, Mccarter MD, Wilson CC. 2012. HIV-1 infection of human intestinal lamina propria CD4⁺ T cells in vitro is enhanced by exposure to commensal *Escherichia coli*. *Journal of Immunology*, **189**(2): 885–896.

Estes JD, Haase AT, Schacker TW. 2008. The role of collagen deposition in depleting CD4⁺ T cells and limiting reconstitution in HIV-1 and SIV infections through damage to the secondary lymphoid organ niche. *Seminars in Immunology*, **20**(3): 181–186.

Estes JD, Harris LD, Klatt NR, Tabb B, Pittaluga S, Paiardini M, Barclay GR, Smedley J, Pung R, Oliveira KM, Hirsch VM, Silvestri G, Douek DC, Miller CJ, Haase AT, Lifson J, Brenchley JM. 2010. Damaged intestinal epithelial integrity linked to microbial translocation in pathogenic simian immunodeficiency virus infections. *PLoS Pathogens*, **6**(8): e1001052.

French MA, King MS, Tschampa JM, Da Silva BA, Landay AL. 2009. Serum immune activation markers are persistently increased in patients with HIV infection after 6 years of antiretroviral therapy despite suppression of viral replication and reconstitution of CD4(+) T cells. *The Journal of Infectious Diseases*, **200**(8): 1212–1215.

Giorgi JV, Hultin LE, Mckeating JA, Johnson TD, Owens B, Jacobson LP, Shih R, Lewis J, Wiley DJ, Phair JP, Wolinsky SM, Detels R. 1999. Shorter survival in advanced human immunodeficiency virus type 1 infection is more closely associated with T lymphocyte activation than with plasma virus burden or virus chemokine coreceptor usage. *The Journal of Infectious Diseases*, **179**(4): 859–870.

Hazenber MD, Otto SA, Van Benthem BHB, Roos MTL, Coutinho RA, Lange JMA, Hamann D, Prins M, Miedema F. 2003. Persistent immune activation in HIV-1 infection is associated with progression to AIDS. *AIDS*, **17**(13): 1881–1888.

Hunt PW, Martin JN, Sinclair E, Bredt B, Hagos E, Lampiris H, Deeks SG. 2003. T cell activation is associated with lower CD4(+) T cell gains in human immunodeficiency virus-infected patients with sustained viral suppression during antiretroviral therapy. *The Journal of Infectious Diseases*, **187**(10): 1534–1543.

Jenabian MA, El-Far M, Vyboh K, Kema I, Costiniuk CT, Thomas R, Baril

- JG, Leblanc R, Kanagaratham C, Radzioch D, Allam O, Ahmad A, Lebouche B, Tremblay C, Ancuta P, Routy JP. 2015. Immunosuppressive tryptophan catabolism and gut mucosal dysfunction following early HIV infection. *The Journal of Infectious Diseases*, **212**(3): 355–366.
- Kaplan RC, Sinclair E, Landay AL, Lurain N, Sharrett AR, Gange SJ, Xue XN, Hunt P, Karim R, Kern DM, Hodis HN, Deeks SG. 2011. T cell activation and senescence predict subclinical carotid artery disease in HIV-infected women. *The Journal of Infectious Diseases*, **203**(4): 452–463.
- Klatt NR, Canary LA, Vanderford TH, Vinton CL, Engram JC, Dunham RM, Cronise HE, Swerczek JM, Lafont BA, Picker LJ, Silvestri G, Brenchley JM. 2012. Dynamics of simian immunodeficiency virus SIVmac239 infection in pigtail macaques. *Journal of Virology*, **86**(2): 1203–1213.
- Klatt NR, Chomont N, Douek DC, Deeks SG. 2013. Immune activation and HIV persistence: implications for curative approaches to HIV infection. *Immunological Reviews*, **254**(1): 326–342.
- Klatt NR, Harris LD, Vinton CL, Sung H, Briant JA, Tabb B, Morcock D, McGinty JW, Lifson JD, Lafont BA, Martin MA, Levine AD, Estes JD, Brenchley JM. 2010. Compromised gastrointestinal integrity in pigtail macaques is associated with increased microbial translocation, immune activation, and IL-17 production in the absence of SIV infection. *Mucosal Immunology*, **3**(4): 387–398.
- Kuang YQ, Tang X, Liu FL, Jiang XL, Zhang YP, Gao G, Zheng YT. 2009. Genotyping of TRIM5 locus in northern pig-tailed macaques (*Macaca leonina*), a primate species susceptible to human immunodeficiency virus type 1 infection. *Retrovirology*, **6**(1): 58.
- Lian XD, Zhang XH, Dai ZX, Zheng YT. 2016. Cloning, sequencing, and polymorphism analysis of novel classical MHC class I alleles in northern pig-tailed macaques (*Macaca leonina*). *Immunogenetics*, **68**(4): 261–274.
- Lian XD, Zhang XH, Dai ZX, Zheng YT. 2017. Characterization of classical major histocompatibility complex (MHC) class II genes in northern pig-tailed macaques (*Macaca leonina*). *Infection, Genetics and Evolution*, **56**: 26–35.
- Lian XD, Zhang XH, Dai ZX, Zheng YT. 2018. Identification of the major histocompatibility complex class-II *DM* and *DO* alleles in a cohort of northern pig-tailed macaques (*Macaca leonina*). *Immunogenetics*, **70**(4): 271–277.
- Liao CH, Kuang YQ, Liu HL, Zheng YT, Su B. 2007. A novel fusion gene, TRIM5-Cyclophilin A in the pig-tailed macaque determines its susceptibility to HIV-1 infection. *AIDS*, **21**: S19–S26.
- Liu Z, Cumberland WG, Hultin LE, Kaplan AH, Detels R, Giorgi JV. 1998. CD8⁺ T-lymphocyte activation in HIV-1 disease reflects an aspect of pathogenesis distinct from viral burden and immunodeficiency. *Journal of Acquired Immune Deficiency Syndromes and Human Retrovirology*, **18**(4): 332–340.
- Ma JP, Xia HJ, Zhang GH, Han JB, Zhang LG, Zheng YT. 2012. Inhibitory effects of chloroquine on the activation of plasmacytoid dendritic cells in SIVmac239-infected Chinese rhesus macaques. *Cellular & Molecular Immunology*, **9**(5): 410–416.
- Marchetti G, Cozzi-Lepri A, Merlini E, Bellistri GM, Castagna A, Galli M, Verucchi G, Antinori A, Costantini A, Giacometti A, Di Caro A, Monforte AD, Grp IFS. 2011. Microbial translocation predicts disease progression of HIV-infected antiretroviral-naïve patients with high CD4(+) cell count. *AIDS*, **25**(11): 1385–1394.
- Marchetti G, Tincati C, Silvestri G. 2013. Microbial translocation in the pathogenesis of HIV infection and AIDS. *Clinical Microbiology Reviews*, **26**(1): 2–18.
- Marks MA, Rabkin CS, Engels EA, Busch E, Kopp W, Rager H, Goedert JJ, Chaturvedi AK. 2013. Markers of microbial translocation and risk of AIDS-related lymphoma. *AIDS*, **27**(3): 469–474.
- Milatz S, Krug SM, Rosenthal R, Gunzel D, Muller D, Schulzke JD, Amasheh S, Fromm M. 2010. Claudin-3 acts as a sealing component of the tight junction for ions of either charge and uncharged solutes. *Biochimica et Biophysica Acta-Biomembranes*, **1798**(11): 2048–2057.
- Pang W, Song JH, Lu Y, Zhang XL, Zheng HY, Jiang J, Zheng YT. 2018. Host restriction factors APOBEC3G/3F and other interferon-related gene expressions affect early HIV-1 infection in northern pig-tailed macaque (*Macaca leonina*). *Frontiers in Immunology*, **9**: 1965.
- Pang W, Zhang GH, Jiang J, Zheng HY, Zhang LT, Zhang XL, Song JH, Zhang MX, Zhu JW, Lei AH, Tian RR, Liu XM, Zhang L, Gao G, Su L, Zheng YT. 2017. HIV-1 can infect northern pig-tailed macaques (*Macaca leonina*) and form viral reservoirs in vivo. *Science Bulletin*, **62**(19): 1315–1324.
- Pelsers MM, Namiot Z, Kisielewski W, Namiot A, Januszkievicz M, Hermens WT, Glatz JFC. 2003. Intestinal-type and liver-type fatty acid-binding protein in the intestine. Tissue distribution and clinical utility. *Clinical Biochemistry*, **36**(7): 529–535.
- Pelsers MMAL, Hermens WT, Glatz JFC. 2005. Fatty acid-binding proteins as plasma markers of tissue injury. *Clinica Chimica Acta*, **352**(1–2): 15–35.
- Sandler NG, Douek DC. 2012. Microbial translocation in HIV infection: causes, consequences and treatment opportunities. *Nature Reviews Microbiology*, **10**(9): 655–666.
- Sandler NG, Wand H, Roque A, Law M, Nason MC, Nixon DE, Pedersen C, Ruxrungtham K, Lewin SR, Emery S, Neaton JD, Brenchley JM, Deeks SG, Sereti I, Douek DC, Grp ISS. 2011. Plasma levels of soluble CD14 independently predict mortality in HIV infection. *The Journal of Infectious Diseases*, **203**(6): 780–790.
- Song TZ, Zhang MX, Xia YJ, Xiao Y, Pang W, Zheng YT. 2018. Parasites may exit immunocompromised northern pig-tailed macaques (*Macaca leonina*) infected with SIVmac239. *Zoological Research*, **39**(1): 42–51.
- Spartaco G. 2001. Notes on the taxonomy of *Macaca nemestrina leonina* Blyth, 1863 (Primates: Cercopithecidae). *Hystrix the Italian Journal of Mammalogy*, **12**(1): 51–54.
- Veazey RS, Demaria M, Chalifoux LV, Shvetz DE, Pauley DR, Knight HL, Rosenzweig M, Johnson RP, Desrosiers RC, Lackner AA. 1998. Gastrointestinal tract as a major site of CD4⁺ T cell depletion and viral replication in SIV infection. *Science*, **280**(5362): 427–431.
- Xia HJ, Zhang GH, Wang RR, Zheng YT. 2009. The influence of age and sex on the cell counts of peripheral blood leukocyte subpopulations in Chinese rhesus macaques. *Cellular & Molecular Immunology*, **6**(6): 433–440.
- Zeng M, Smith AJ, Wietgreffe SW, Southern PJ, Schacker TW, Reilly CS, Estes JD, Burton GF, Silvestri G, Lifson JD, Carlis JV, Haase AT. 2011. Cumulative mechanisms of lymphoid tissue fibrosis and T cell depletion in HIV-1 and SIV infections. *Journal of Clinical Investigation*, **121**(3): 998–1008.
- Zhang LT, Tian RR, Zheng HY, Pan GQ, Tuo XY, Xia HJ, Xia XS, Pang W, Zheng YT. 2016a. Translocation of microbes and changes of immunocytes in the gut of rapid- and slow-progressor Chinese rhesus macaques infected

- with SIVmac239. *Immunology*, **147**(4): 443–452.
- Zhang MX, Zheng HY, Jiang J, Pang W, Zhang GH, Zheng YT. 2016b. Viral seroprevalence in northern pig-tailed macaques (*Macaca leonina*) derived from Ho Chi Minh City, Vietnam. *Primates*, **57**(3): 413–419.
- Zhang MX, Zheng HY, Jiang J, Song JH, Chen M, Xiao Y, Lian XD, Song TZ, Tian RR, Pang W, Zheng YT. 2017a. Northern pig-tailed macaques (*Macaca leonina*) maintain superior CD4⁺ T-cell homeostasis during SIVmac239 infection. *European Journal of Immunology*, **48**(2): 384–385.
- Zhang MX, Zheng HY, Jiang J, Song JH, Chen M, Xiao Y, Lian XD, Song TZ, Tian RR, Pang W, Zheng YT. 2017b. Predict disease progression from T-cell phenotypes in northern pig-tailed macaques (*Macaca leonina*) during SIVmac239 infection. *Immunology*, **152**(4): 574–579.
- Zhang XL, Pang W, Deng DY, Lv LB, Feng Y, Zheng YT. 2014. Analysis of immunoglobulin, complements and CRP levels in serum of captive northern pig-tailed macaques (*Macaca leonina*). *Zoological Research*, **35**(3): 196–203.
- Zhang XL, Song JH, Pang W, Zheng YT. 2016c. Molecular cloning and anti-HIV-1 activities of APOBEC3s from northern pig-tailed macaques (*Macaca leonina*). *Zoological Research*, **37**(4): 246–51.
- Zheng HY, Zhang MX, Pang W, Zheng YT. 2014a. Aged Chinese rhesus macaques suffer severe phenotypic T- and B-cell aging accompanied with sex differences. *Experimental Gerontology*, **55**: 113–119.
- Zheng HY, Zhang MX, Zhang LT, Zhang XL, Pang W, Lyu LB, Zheng YT. 2014b. Flow cytometric characterizations of leukocyte subpopulations in the peripheral blood of northern pig-tailed macaques (*Macaca leonina*). *Zoological Research*, **35**(6): 465–473.
- Zhu L, Lei AH, Zheng HY, Lyv LB, Zhang ZG, Zheng YT. 2015. Longitudinal analysis reveals characteristically high proportions of bacterial vaginosis-associated bacteria and temporal variability of vaginal microbiota in northern pig-tailed macaques (*Macaca leonina*). *Zoological Research*, **36**(5): 285–298