

Evaluation of the Effectiveness of the Use of Papaya Fruit Latex in Making Herbal Medicated Multinutrition Block as a Local Goat Feed Supplement

Retno Iswarin Pujaningsih, Baginda Iskandar Moeda Tampoebolon, Widiyanto and Dian Wahyu Harjanti

Department of Animal Husbandry, Faculty of Animal and Agriculture Science, University of Diponegoro,
Jl. Prof. Soedarto, Tembalang, Semarang, 50275 Central Java, Indonesia
*Corresponding author e-mail: retno.marwoto@gmail.com

Abstract. The use of papaya latex formulated with nitrogen supplementation and minerals in the form of multinutrition blocks that can pass through the rumen is expected to improve nutritional quality and feed digestibility. This study aims to evaluate the effectiveness of the use of fermented rice straw, papaya fruit latex and blood clamshell flour as a main raw material in the manufacture of multinutrition blocks plus herbal anthelmintic medicine (papaya fruit latex) on physical quality (texture, color, aroma, hardness), chemical quality (proximate composition and mineral content of Ca, P, Mg, Cu, Zn). Microbiological tests were carried out on total bacteria, Salmonella and Escherichia coli. The research design used was a completely randomized design with 4 treatment levels of papaya fruit administration and 4 replications. Data is processed statistically using ANOVA. The results of organoleptic physical quality analysis showed significant differences in the texture of the product given the highest papaya latex (0.007ml / kg). Chemical analysis of the product showed a quantitative increase in protein, although statistically the treatment did not show any significant differences. The results of the analysis of the microbiological quality of the product quantitatively showed that the resulting multinutrition block herbs were not contaminated with negative bacteria. It was concluded that multinutrition block herbs can be given to livestock as supplementary feed.

Key words: papaya latex, multinutrition blocks, anthelmintic herbal, local goat

Abstrak. Pemberian getah pepaya yang diformulasikan dengan pemberian suplementasi nitrogen dan mineral dalam bentuk multinutrisi blok yang dapat melewati rumen diharapkan mampu meningkatkan kualitas nutrisi dan pencernaan pakan. Penelitian ini bertujuan untuk mengevaluasi efektifitas penggunaan jerami padi yang difermentasi, getah buah pepaya serta tepung cangkang kerang darah sebagai bahan baku andalan pada pembuatan multinutrisi blok plus anthelmintic medicine herbal (getah buah pepaya) terhadap kualitas fisik (tekstur, warna, aroma, kekerasan), kualitas kimia (komposisi proksimat dan kandungan mineral Ca, P, Mg, Cu, Zn). Uji mikrobiologis dilakukan terhadap total bakteri, Salmonella dan Escherichia coli. Desain penelitian yang digunakan adalah rancangan acak lengkap dengan 4 perlakuan level pemberian getah buah pepaya dan 4 ulangan. Data diolah secara statistik menggunakan anova. Hasil analisis kualitas fisik secara organoleptik menunjukkan perbedaan yang nyata pada tekstur produk yang diberi getah pepaya tertinggi (0,007ml/kg). Analisis produk secara kimiawi menunjukkan adanya peningkatan protein secara kuantitatif meskipun secara statistik perlakuan tidak menunjukkan perbedaan yang nyata. Hasil analisis kualitas mikrobiologi produk secara kuantitatif menunjukkan bahwa herbal multinutrisi blok yang dihasilkan tidak tercemar bakteri negatif. Disimpulkan bahwa herbal multinutrisi blok dapat diberikan kepada ternak sebagai pakan pelengkap.

Kata kunci: getah pepaya, multinutrisi blok, anthelmintic herbal, kambing lokal

Introduction

The unbalanced effects of nutrient supply on ruminants include low productivity, susceptibility to disease infestations that cause high mortality rates. Nematode worm infestation in many cases reported by

Suhardono et al., (2012) occurs in young goats during growth. According to Malesi (2010) worms that commonly interfere with the health of goats and sheep are from the Nematode class, and the most dangerous type is *Haemonchus contortus* (tapeworm) which lives in the

abomasum and lives as a blood sucking with infestation rates according to Lastuti et al., (2006) can reach 80%. This worm will lead to death in livestock, the growth of livestock will be disrupted to cause reproductive problems (Malesi, 2010)

Goats that show symptoms of intestinal worms can be treated with traditional medicine or with manufacturing drugs. Traditional medicines such as papaya latex or papaya leaf juice are cheaper, easier to obtain, can be done by farmers, and have no harmful side effects compared to manufacturer's drugs. According to Permana (2013) papaya (*Carica papaya* L.) is a plant that almost all parts of the plant can be used for various purposes. Almost all parts of this plant from the roots, leaves, sap, to the seeds, have been empirically used as anthelmintics. Anthelmintic is an ingredient used to destroy worms in the digestive tract of goats.

Increasing the productivity of ruminants through the provision of practical nutrition must be developed so that it is easily applied by smallholder farmers. This includes feed supplementation using block multinutritions containing minerals according to the standard requirements of goats and microbial growth factors to increase feed digestibility. In this study, multinutrition blocks were given additional herbal anthelmintic to overcome nematode worm infestations which can interfere with the growth of livestock.

Materials and Methods

Providing multinutrition block herbs was done by using raw materials in the form of rice straw fermented with EM4, blood clam shells, young papaya sap, molasses, urea, bentonite, salt and lime. The main equipment used is mold blocks, grinders and pans to heat molasses.

Providing herbal medicated multinutrition block

Before being mixed with other raw materials, rice straw is fermented for 14 days with EM4 and

rice bran, then dried and ground. Clamshell also needs to be grinded first before mixed together in the formulation. Multinutrition block herbal formulations showed on Table 1. Molasses were heated for 10 minutes at 40°C then mixed with other raw materials. Papaya fruit latex is added according to research treatment. After homogeneous the mixture is weighed 600 grams each and formed using a 10 cm diameter pipe. Then after being solid and hardened, divided into 4 with the same weight. Each mixture is made 4 times for each treatment.

Table 1. Formulation of Herbal Medicated Multinutrition Block

Raw material	%
Molasses	50
Fermented rice straw	30
Salt	3
Clamshell	6
Urea	4
Bentonite	7

Tapping papaya fruit latex

To get young papaya latex, young papaya fruit is sliced as deep as a maximum of 2 mm. Each fruit has a maximum of 4 slices with a distance of 2 cm. Young papaya sap is collected for 2 hours every 3 days on the same fruit.

Treatment

The level of papaya fruit latex that ws given is T0: 0 ml / kg, T1: 0.003 ml / kg, T2: 0.005 ml / kg and T3: 0.007 ml / kg for the treatment. Variables were measured organoleptically (texture, color, hardness and aroma), chemistry (proximate composition and minerals Ca, P, Mg, Cu, Zn) and microbiology (total bacteria, *Escherichia coli*, *Salmonella sp.*). The research design used was a completely randomized design with 4 treatments and 4 replications. Data analysis using ANOVA.

Results and Discussion

Physical quality test results

The organoleptic block multinutrition quality test was carried out by 10 trained semi-panelists. Organoleptic test results are shown in Table 2.

Table 2. Physical quality test results of herbal medicated multinutrition block

Treatment	Colour	Texture	Aroma	Hardness
T0	4	3.64 ^a	3.88	3.28 ^a
T1	4	3.64 ^a	3.88	3.3 ^a
T2	4	3.78 ^a	3.78	3.35 ^a
T3	4	3.98 ^b	3.82	3.65 ^b

Different superscripts in the same column show different treatments ($P < 0.05$)

Notes: T0: without papaya latex, T1 with 0,003 ml/kg of papaya latex, T2: with 0,005 ml/kg of papaya latex, T3: with 0,007 ml/kg of papaya latex

Based on Table 2 there are significant differences in texture and hardness of herbal medicated multinutrition block (T3) treatment compared to other treatments (T0, T1, T2). The addition of 0.007 ml / kg of latex papaya is thought to cause a tighter texture and denser hardness than other treatments. Papaya latex is a white liquid that is sticky and released from plants that have laticifer or sap producers (Kusumadjaja and Dewi, 2005). Refer to Yusha'u et al., (2009) the milky juice of papaya (latex papaya) is used as chewing gum. The ability of papaya latex to bind the ingredients of the herbal medicated multinutrition block affects the texture and hardnesses to become more tight and dense.

The sensor of color and aroma has no significant difference since molasses becomes the dominant amount of herbal medicated multinutrition block. Molasses is responsible for the dark color of the product. Its dark color caused by the second boiling cycle of sugarcane (Kaushik et al., 2018)

Chemical quality test results

The proximate composition of multinutrition block herbs is shown in Table 3. From Table 3 it

can be seen that the nutritional content of multinutrition herbs in the treatment blocks of T1 and T2 is quantitatively different from the percentage of protein content. While the other nutritional content is not significantly different. The fermentation process of rice straw using EM4 and fermented for 14 days can increase the crude protein content. During the fermentation process there is a change in structured carbohydrates and non-structural carbohydrates proven by the decrease in the content of crude fiber in fermented rice straw by 30.88% of crude fiber content of unfermented rice straw by 32.14% (Table 4). The addition of papaya fruit latex although not statistically significant, quantitatively tends to increase the protein content of the herbal multinutrition block. According to Rodrigues et al., (2009) papaya fruit latex contains about 30% of protease enzymes and 10% of other proteins.

Although the mineral content of rice straw is quite complete, in general it is relatively lower compared to the mineral content of grasses and legumes. Low quality feed which has a high content of fiber mostly contains very low minerals so that it becomes a limiting factor for microbial growth in the rumen such as P and S content. Therefore, it is necessary to add minerals to the livestock ration if rice straw is used as the main feed.

Mineral content in rice straw is used by microorganisms found in EM4 to help metabolize activity in digesting crude fiber. Stimulatory factors that significantly increase the value of substrate digestibility by stem bacteria ($P < 0.05$) when compared to controls are Fe; Zn; Se; Cu; Co, Mo, and Mn. There are similarities and differences in effects between substances used in the activity of bacterial cocci and stems. Zn; Co; Cu and Mo increase the activity of bacterial cocci and stem (Thalib et al., 2000). In general, the mineral content of multinutrition herbal blocks in each treatment was not significantly different.

Table 3. Proximate composition of herbal medicated multinutrition block

Treatment	Dry		Crude	Crude	Protein
	Matter	Ash	fiber	Fat	
T0	75.54	15.66	9.34	1.77	4.69
T1	75.60	15.68	9.31	1.80	4.72
T2	75.75	15.72	9.40	1.82	4.76
T3	75.80	15.74	9.35	1.83	4.81

Table 4. Nutrient content of rice straw

Sample	Dry		Crude	Crude	Protein
	matter	Ash	fiber	fat	
RS	84.22	22.25	32.14	3.32	5.31
FRS	79.11	21.34	30.88	2.41	7.72

Note: RS: Rice straw, FRS: Fermented Rice Straw

Table 5. Herbal medicated multinutrition block mineral content

Treatment	Ca	P	Mg	Cu	Zn
T0	147.53	0.16	8.01	1.92	0.16
T1	147.80	0.16	8.15	1.95	0.16
T2	148.27	0.16	8.19	1.98	0.16
T3	148.11	0.17	8.21	2.00	0.17

Microbiological quality test results

The addition of EM4 to the rice straw fermentation process affected the total number of multinutrition block bacteria (Table 6). In general no *Escherichia coli* and *Salmonella sp* bacteria were identified. on block multinutrition. The presence of flavonoids in papaya fruit latex which functions as antimicrobials plays a role in preventing the development of pathogenic bacteria (Kaushik et al., 2018)

Table 6. Total Bacteria, Gram Bacteria, Identification of *Escherichia coli* and *Salmonella sp*.

Parameter	T0	T1	T2	T3
Total Bakteri (cfu/g)	1,26 × 10 ⁴	1,26 × 10 ⁴	1,29 × 10 ⁴	1,28 × 10 ⁴

Bakteri Gram *	3,4	3,7	3,7	3,6
<i>E. coli</i> **	Negatif	Negatif	Negatif	Negatif
<i>Salmonella sp.</i> **	Negatif	Negatif	Negatif	Negatif

Notes : * : score 1 – 4

** : qualitatively

Conclusions

From the results of this work, it can be concluded that herbal medicated multinutrition block can be given to livestock as supplementary feed.

Acknowledgement

I would like to thank the Research and Community Service Institutions of UNDIP for the chance to have funding and support through my research. I would like to thank my teams for accepting nothing less than excellence from me. Last but not the least, I would like to thank my students and Mr. Julianto, the owner of the goat farm, for supporting me finishing this work.

References

- Amin, M, S.D Hasan, O. Yanuarianto, M. Iqbal dan I.W Karda. 2016. Peningkatan Kualitas Jerami Padi Menggunakan Teknologi Amoniasi Fermentasi. Jurnal Ilmu dan Teknologi Peternakan Indonesia, Volume 2 (1): 96–103; Juni 2016 ISSN: 2460-6669
- Babale D. M., Kibon A and Yahaya M. S. 2015. Performance and linear body measurements of Red Sokoto male goats on replacement levels of corn cobs for maize bran with cowpea husk basal diet. Net Journal of Agricultural Science. Vol. 3(2), pp.35-40, April 2015. ISSN: 2315-9766
- Bakshi, M.P.S and Wadhwa, M. 2011. Nutritional status of dairy animals in different regions of Punjab State in India. Indian Journal of Animal Science 81: 52–58.
- Bheekhee, H., B. Bulman, RK. Boodoo, R Rammuth, L. Heung, R Yeun, R Fakim and B. Dobe, 1999. Development and field evaluation of animal feed supplementation packages for improving meat and milk production in ruminant livestock using locally available feed resources. Livestock Research Department, Agricultural Research and

- Extension Unit, Food and Agricultural Research Council, Quatre Bomes, Mauritius, pp: 1-9.
- Dinas Peternakan dan Kesehatan Hewan Provinsi Jawa Tengah. 2016. Sensus Hasil Pertanian Provinsi Jawa Tengah 2013. http://www.pertanian.go.id/dinakeswan_jateng
- Fayomi, A., Ahmed, A., Musa, U., Salami-Shinaba, J.O., Ogedegbe, S.A and Akanni, K. 2014. Moringa Multi-Nutrient Blocks: Formulation, Production and Feeding Trial Under a Tropical Environment. International Journal of Science, Environment and Technology, Vol. 3, No 1, 2014, 67 – 84 ISSN 2278-3687
- Garcia, L.O and J.I.R. Restrepo, 1995. Multinutrient Block Handbook (FAO Better Farming Series no. 45). Food and Agriculture Organization of United Nation, Rome.
- Kaushik, A., S.Basu, V.S.Batra and M.Balakrishnan. 2018. Fractionation of sugarcane molasses distillery wastewater and evaluation of antioxidant and antimicrobial characteristics. Industrial Crops and Products. Volume 118: 73-80. <https://doi.org/10.1016/j.indcrop.2018.03.040>
- Kusumadaja, PA and R.P. Dewi. 2005. Determination of Optimum Condition of Papain Enzyme from Papaya Var Java. Indo. J. Chem., 2005, 5 (2), 147 – 151. <https://doi.org/10.22146/ijc.21822>
- Lukiwati, D.R dan R.I. Pujaningsih. 2014. Efek Sisa Pupuk Kandang Diperkaya Fosfat Alam terhadap Produksi Jagung Manis dan Jerami di Lahan Kering. Jurnal Lahan Suboptimal. 3(2): 152-160.
- Lukiwati, D.R. dan R.I. Pujaningsih. 2015. Efek sisa pupuk kandang diperkaya fosfat alam dalam bentuk granular dan di inokulasi biodekomposer terhadap nutrisi jerami jagung manis di lahan kering. Jurnal Pastura,(terindeks DOAJ dan Google Scholar). 4(2): 78-82.
- McAllan, A.B. 1991. Optimizing the Use of Poor Quality Forage Feed Resources for Ruminant Production: Supplementation with Bypass Nutrients. In Proceeding of a Symposium : Isotope and Related Techniques in Animal Production and Health. Vienna, 15-19 April, 1991. Jointly organized by IAEA and FAO. pp.25-41
- Mohammed, I.D, M. Baulube and A. Adeyinka. 2007. Multi-nutrient Blocks I: Formulation and Production under a Semi-arid Enviromuent of North East Nigeria. Journal of Biological Sciences 7 (2): 389-392. ISSN 1727-3048
- Mubi, AA., Kibon, A and Mohammed, I.D. 2011. Effects of multinutrient blocks supplementation on the performance of grazing Yankaza sheep in the wet season of guinea savanna region of Nigeria. International Journal of Sustainable Agriculture 3: 103-06
- Pujaningsih, R. I, Mukodiningsih, Sri, Pakpahan, Irjon. 2017. Cassava waste processing technology to support the provision of alternative feed on zero waste management system of livestock. Advanced Science Letters, Volume 23, Number 3, March 2017, pp. 2595-2597(3). American Scientific Publishers. DOI: <https://doi.org/10.1166/asl.2017.8726>
- Rodrigues S.P., da Cunha M., Ventura J.A. and Fernandes P.M.B. 2009. Effects of the *Papaya meleira virus* on papaya latex structure and composition. Plant Cell Rep.28:861–871. doi: 10.1007/s00299-009-0673-7.
- Simbolon, Nico, R.I. Pujaningsih dan S. Mukodiningsih. 2016. Pengaruh berbagai pengolahan kulit singkong terhadap pencernaan bahan kering dan bahan organik secara *in vitro*, protein kasar dan asam sianida. Jurnal Ilmu-Ilmu Peternakan 26 (1): 58 - 65
- Sulistiyanto, B, S.Sumarsih, C.S Utama, R.I.Pujaningsih, C.I.Sutrisno. 2012. Effect of drying methods to the lactic acid bacteria of enriched cassava meals. Bul. Sintesis. Vol 16 (2). Hal : 23-26. Desember 2012
- Suttle, N.F. 1991. Mineral supplementation of low quality roughage. In Proceeding of a Symposium : Isotope and Related Techniques in Animal Production and Health. Vienna, 15-19 April, 1991. Jointly organized by IAEA and FAO. pp.113-126.
- Syahrani, Syahrir, M. Zain Mide, Rohmiyatul Islamiyati dan Ani Asrianie. 2015. Efektifitas Fermentasi Rumen thd Pakan Campuran Jerami Padi dan Biomassa Murbei dengan Penambahan Urea Mineral Molases Liquid. JITP Vol. 4 No. 1, Januari 2015
- Tampoebolon, BIM, Z. Bachruddien, L.M. Yusiati dan S. Margino. 2013. Pengaruh Aplikasi Bakteri Lignoselulolitik Usus Rayap Terseleksi Pada Fermentasi Jerami Padi Terhadap Komposisi Serat. Jurnal Sintesis (Ilmu-ilmu Pertanian) Yayasan Dharma Agrika Semarang Volume 17, no 1. Juni.

- Van Soest, P.J. 1994. Nutritional Ecology of the Ruminant. 2nd Ed. Cornell Univ. Press. Ithaca and London.
- Wadhwa, M and M.P.S. Bakshi. 2014. Nutritional evaluation of urea molasses multi-nutrient blocks containing agro-industrial wastes in buffaloes. Indian Journal of Animal Sciences 84 (5): 544–548.
- Widiyanto, E. Kusumanti, M. Mulyono and S. Surahmanto. 2012. The Influence of Season and Topography on Manganese (Mn) Status of Grazing Java Thin-Tailed Sheep in the Agriculture Area in Mijen of Semarang-Central Java. Journal Of Indonesian Tropical Animal Agriculture. Vol 38, No 2 (2013). DOI: 10.14710/jitaa.38.2.131-138
- Widiyanto, E. Pangestu, Surahmanto, V.D. Yuniyanto, B.I.M. Tampobolon and B.W.H.E. Prasetyono. 2015a. Effect of Mineral Supplementation and Introduction of *Setaria sphacelata* Grass and *Gliricidia sepium* Legume on Productivity of Kacang Goat at Serang River Basin Upland Area, Central Java, Indonesia. Pakistan Journal of Nutrition. Volume: 14(8) pp. 440-446. DOI: 10.3923/pjn.2015.440.446
- Widiyanto, Sumarsono, Sudjatmogo, B.W.H.E Prasetyono, A.Setiadi and Surahmanto. 2015b. Iodine Mineral Status of Etawah Crossbred Goat at Different Physiological Stages Fed Elephant Grass and Tofu Byproduct. Journal of Animal Production. Vol 17, No 1 (2015)
- Yusha’u, M., Onuorah, F. C. and Murtala, Y. 2009. N-Vitro Sensitivity Pattern of Some Urinary Tract Isolates to Carica Papaya Extracts. Bayero Journal of Pure and Applied Sciences, 2(2): 75 – 78.