Dental Chronicles

Case 1: Class II Div 2 Case

Dr. Devinder Preet Singh

DOI: https://doi.org/10.5281/zenodo.10369366

fifteen year old male patient reported to the dental clinic with a chief complaint of irregular upper front teeth. Intraorally the patient had Class II molar and canine relationship, division 2 incisors with deep overbite. The patient was skeletal Class II with deficient mandible. The patient was treated with comprehensive fixed mechanotherapy with self ligating metal brackets having 0.022" slot. After leveling and alignment of the teeth which took about 9 months, the patient was given Forsus Fixed Functional Appliance, to stimulate the growth of the mandible, for 6 months to correct the skeletal Class II. The total treatment took about I.5 years with the achievement of Class I molar and skeletal relationship and ideal overjet and overbite. Permanent lower bonded retention from canine to canine was given with upper removable anterior inclined plane to retain the Class II correction.

Extra-oral Photographs (Pre-Treatment)



Frontal View







Profile View



Right Lateral

Intra-oral Photographs (Pre-Treatment)



Left Lateral



Frontal View





Maxillary Occlusal



Mandibular Occlusal



OPG (Orthopantomogram)



Lateral Cephalogram

Intra-Oral Photographs (Forsus Appliance)



Frontal View



Left Lateral



Right Lateral



Extra-oral Photographs (Post-Treatment)



Frontal View





Smiling View

Profile View

Intra-oral Photographs (Post-Treatment)



Right Lateral



Left Lateral



Frontal View



Maxillary Occlusal



Mandibular Occlusal







OPG (Orthopantomogram)

Lateral Cephalogram

Case 2 Anterior Crossbite

A twenty two year old female patient reported to the dental clinic with a chief complaint of irregular upper front teeth. Intraorally the patient had Class I molar and canine relationship with anterior crossbite irt. 11-41, 21-31 and rotated 12,22, 32,42. The patient was skeletal Class I with horizontal growth pattern. The patient was treated with comprehensive fixed mechanotherapy with conventional metal brackets having 0.022" slot. The leveling and alignment of the teeth took about 9 months with the correction of anterior cross bite.. The total treatment took about I.2 years with the achievement of Class I incisor relationship and ideal overjet and overbite. Permanent lower bonded lingual retention from canine to canine was given with upper Hawley's retainer.

Extra-oral Photographs (Pre-Treatment)



Frontal View





Smiling View

Profile View



Intra-oral Photographs (Pre-Treatment)







Frontal View



Left Lateral



Maxillary Occlusal



Mandibular Occlusal



OPG (Orthopantomogram)

26



Lateral Cephalogram

Extra-oral Photographs (Post-Treatment)







Frontal View

Smiling View

Profile View

Intra-oral Photographs (Post-Treatment)



Right Lateral







Frontal View



Maxillary Occlusal



Mandibular Occlusal





OPG (Orthopantomogram)



Lateral Cephalogram

