

## A note on uniquely (nil) clean ring

Shervin Sahebi<sup>a</sup>, and Mina Jahandar <sup>a,\*</sup>

<sup>a</sup>Department of Mathematics, Islamic Azad University, Central Tehran Branch, PO.  
Code 14168-94351, Iran

---

**Abstract.** A ring  $R$  is uniquely (nil) clean in case for any  $a \in R$  there exists a uniquely idempotent  $e \in R$  such that  $a - e$  is invertible (nilpotent). Let  $C = \begin{pmatrix} A & V \\ W & B \end{pmatrix}$  be the Morita Context ring. We determine conditions under which the rings  $A, B$  are uniquely (nil) clean. Moreover we show that the center of a uniquely (nil) clean ring is uniquely (nil) clean.

---

**Keywords:** Full element, uniquely clean ring, nil clean ring

---

### 1. Introduction

We say that an element  $a \in R$  is uniquely (nil) clean provided that there exists a unique idempotent  $e \in R$  such that  $a - e \in R$  is invertible (nilpotent). A ring  $R$  is uniquely (nil) clean in case every element in  $R$  is uniquely (nil) clean. As is well known, every uniquely nil clean ring is uniquely clean. Many authors have studied such rings, see [? ? ? ? ?].

A Morita Context  $(A, B, W, V, \psi, \varphi)$  consists two rings  $A, B$ , two bimodules  ${}_A V_B$ ,  ${}_B W_A$  and a pair of bimodule homomorphisms  $\psi : V \otimes_B W \rightarrow A$ ,  $\phi : W \otimes_A V \rightarrow B$ , such that  $\psi(v \otimes w)v' = v\phi(w \otimes v')$ ,  $\phi(w \otimes v)w' = w\psi(v \otimes w')$ . We can form

$$C = \left\{ \begin{pmatrix} a & v \\ w & b \end{pmatrix} \mid a \in A, b \in B, v \in V, w \in W \right\}$$

and define a multiplication on  $C$  as follows:

$$\begin{pmatrix} a & v \\ w & b \end{pmatrix} \begin{pmatrix} a' & v' \\ w' & b' \end{pmatrix} = \begin{pmatrix} aa' + \psi(v \otimes w') & av' + vb' \\ wa' + bw' & \phi(w \otimes v') + bb' \end{pmatrix}$$

A routine check shows that, with this multiplication (and entry-wise addition),  $C$  becomes an associative ring. We call  $C$  a Morita Context ring [?]. Obviously, the class of the rings of Morita Contexts includes all  $2 \times 2$  matrix rings and all formal triangular matrix rings. In recent years, many authors studied Morita Contexts from different points of view [? ?].

---

\*Corresponding author. Email: m66.jahandar@gmail.com

In this paper in the first section we obtain the relationship of uniquely (nil) cleanness between Morita Context ring  $C$  and  $A, B$ . At last in the second section, we investigate if the center of a uniquely (nil) clean rings are uniquely (nil) clean? Throughout, all rings are associative rings with identity.  $Z(R)$  will denote, the center of  $R$ .

### 1.1 Morita Context ring

The following results are useful tools needed in the proof of main results.

**THEOREM 1.1** (see [?, Theorem 2.2] and [?, Corollary 3.3.7]) *Every factor ring of uniquely (nil) clean ring is again uniquely (nil) clean.*

**LEMMA 1.2** *Every idempotent in a uniquely clean ring is central.*

*Proof* Let  $e^2 = e \in R$ . If  $r \in R$ , then  $e + (er - ere)$  is an idempotent. Hence  $1 + (er - ere)$  is a unit, so the fact that  $[e + (er - ere)] + 1 = e + [1 + (er - ere)]$  implies that  $e + (er - ere) = e$  because  $R$  is uniquely clean. It follows that  $er = ere$ , and similarly  $re = ere$ . ■

**LEMMA 1.3** *Every idempotent in uniquely nil clean ring is central.*

*Proof* Let  $e \in R$  be an idempotent and let  $r$  be any element of  $R$ . Notice that the element  $e + er(1 - e)$  can be written as  $e + (er(1 - e))$  or as  $(e + er(1 - e)) + 0$  as the sum of an idempotent and a nilpotent. Since  $R$  is uniquely nil clean, this shows that  $e = e + er(1 - e)$ , implying that  $er(1 - e) = 0$ . It can likewise be shown that  $(1 - e)re = 0$ , so  $e$  is central. ■

**THEOREM 1.4** *Let  $C = \begin{pmatrix} A & V \\ W & B \end{pmatrix}$  be the Morita Context with  $\varphi, \psi = 0$ . If  $C$  is a uniquely (nil) clean ring then  $A, B$  are uniquely (nil) clean rings.*

*Proof* Let  $I = \begin{pmatrix} 0 & V \\ W & B \end{pmatrix}$ ,  $J = \begin{pmatrix} A & V \\ W & 0 \end{pmatrix}$ . One can check that  $I, J$  are ideals of  $C$  and  $C/I \simeq A$ ,  $C/J \simeq B$ . The uniquely (nil) cleanness of  $A, B$  follows from Theorem 2.1. ■

The following example shows that the converse of Theorem 2.4 is not true.

**Example 1.5** Let  $C = \begin{pmatrix} \mathbb{Z}_2 & \mathbb{Z}_2 \\ 0 & \mathbb{Z}_2 \end{pmatrix}$ . One can check that  $\mathbb{Z}_2$  is uniquely (nil) clean. Since  $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$  is a noncentral idempotent in  $C$ , then  $C$  is not uniquely (nil) clean, by Lemma 2.2 and Lemma 2.3.

**COROLLARY 1.6** *Let  $R, S$  be two rings, and  $M$  be an  $(R, S)$ -bimodule. Let  $E = \begin{pmatrix} R & M \\ 0 & S \end{pmatrix}$  be the formal triangular matrix ring. If  $E$  is a uniquely (nil) clean ring then  $R$  and  $S$  are uniquely (nil) clean rings.*

*Proof* Formal triangular matrix rings are special cases of the Morita Context rings with zero morphisms, therefore the result follows by Theorem 2.1. ■

## 2. The center of uniquely (nil) clean rings

It is interesting to know if the center of a ring shares the same property with the ring. We don't know if the center of a clean ring is necessarily clean? But we have:

**THEOREM 2.1** *The center of a uniquely (nil) clean ring is uniquely (nil) clean.*

*Proof* Let  $R$  be a uniquely (nil) clean ring and  $x \in Z(R)$ . Then there exists a unique idempotent  $e \in R$  such that  $x - e \in R$  is invertible (nilpotent). Since  $e \in Z(R)$  by Lemma 2.2 and Lemma 2.3, then  $x - e \in Z(R)$ . Thus  $x$  is uniquely (nil) clean in  $Z(R)$ . ■

## References

- M. Y. Ahn, (2003). Weakly clean rings and almost clean rings. Ph.D. Thesis, University of Iowa.
- D. D. Anderson, V. P. Camillo, Commutative rings whose elements are a sum of unit and idempotent. *Comm. Algebra* 30 (2002), pp. 3327–3336.
- B. Li, L. Feng, F-clean rings and rings having many full elements. *J. Korean Math. Soc.* 2 (2010), pp. 247–261.
- J. Che, W. K. Nicholson, Y. Zhou, Group rings in which every element is uniquely the sum of a unit and idempotent. *J. Algebra.* 306 (2006), pp. 453–460.
- H. Chen, Morita contexts with many units. *Comm. Algebra.* 30 (3) (2002), pp. 1499–1512.
- A. J. Diesl, Classes of strongly clean rings. Ph.D. Thesis, University of California, Berkeley, (2006).
- A. Haghany, Hopficity and co-hopficity for Morita Contexts. *Comm. Algebra.* 27(1)(1999), pp. 477–492.
- W. K. Nicholson, Y. Zhou, Rings in which elements are uniquely the some of an idempotent and unit. *Clasy. Math. J.* 46(2004), pp. 227–236.