

History of study, updated checklist, distribution and key of scorpions (Arachnida: Scorpiones) from China

Zhi-Yong DI¹, Zi-Zhong YANG², Shi-Jin YIN³, Zhi-Jian CAO¹, Wen-Xin LI^{1,*}

1. College of Life Sciences, Wuhan University, Wuhan Hubei 430072, China

2. Yunnan Provincial Key Laboratory of Entomological Biopharmaceutical Dali University, Dali Yunnan 671003, China

3. College of pharmacy, South-Central University for Nationalities, Wuhan Hubei 430074, China

Abstract: This review describes the history of taxonomic research on scorpions and provides an updated checklist and key of the scorpions currently known in China. This checklist is based on a thorough review of the extant literatures on scorpion species whose presence has been confirmed in China through field expeditions and examination of scorpion collections, excepting a few members that have no clear distribution or are currently in doubt. Totally, the scorpion fauna of China consists of 53 species and subspecies belonging to 12 genera crossing five families, with 33 species (62.3%) and one genus being recorded as endemic. Additionally, identification key and the distribution of scorpions from China are provided.

Keywords: Scorpion; Taxonomy; Checklist; Key; Distribution; China

China is comprised of a vast territory crossing frigid, temperate, and tropical zones, as well as complex topography (80% of which are mountains) including rich rivers, lakes, and diverse climate. These features contribute to China's possession of high species and habitat diversity, making it to be one of 12 megadiversity countries. For example, there are at the least 1.5×10^4 species estimated conservatively, which is about 1/10 of the sum of the entire world (88 328, 929 050 species recorded in China and the world respectively, Yu, 2004).

Of these species, Scorpions, though as a small arachnid group, are quite interesting. To date, there are 15 families, 197 genera and 2 069 species recorded in the world (6/20/2013, <http://www.ntnu.no/ub/scorpion-files/index.php>). Aristotle (384–322 BC) was the first scorpion researcher who studied zoological information about biogeography of scorpions in the western world (Fet et al, 2009), a search enhanced by the systematics research began by Carl von Linne (1758), who recorded five scorpion species in the tenth edition of *Systema Naturae*, and classified them as the members of the genus *Scorpio* in Insect Aptera (Fet et al, 2002).

Chinese reported scorpions from 2 000 years ago, as

the simple information in *Er ya* (published in 221 BC–9 AD) (Wang & Chen, 2007). Scorpions are found in some multifarious ancient books and local chronicles, especially the traditional Chinese medicine books, such as, *Shu ben cao*, *Ben cao tu jing*, *Ben cao jing shu*, and *Ben cao bei yao* (Zhang et al, 2009). However, all of these historical data contained no value on classification. The beginning of Chinese scorpion taxonomy research was started by foreign scientists. Prior to 2003, there were 19 species and subspecies reported in China, as noted by Zhu et al (2004). *Buthus confucius* Karsch (1879) (= *Buthus martensii* = *Mesobuthus martensii*, belonging to the family Buthidae) was the first species described from China (Zhu et al, 2004). Simon (1880) described a new species, *Buthus confucius* (Buthidae), on the basis of specimens collected in Beijing (cited in Qi et

Received: 8 November 2013; Accepted: 13 December 2013

Foundation items: This work was supported by grants from the National Natural Sciences Foundation of China (31071942), the Basic Project of Ministry of Science and Technology of China (2007FY210800), and the 973 program (2010CB529800).

* Corresponding author, E-mail: liwxlab@whu.edu.cn

al, 2004; Shi & Zhang, 2005). Kraepelin (1899) reviewed the order Scorpiones, including 13 species and subspecies from China. Karsch (1881) also suggested that *Buthus confucius* might be a synonym of *Buthus martensii* (Qi et al, 2004; Shi & Zhang, 2005). Birula (1897, 1911) named three subspecies from China: *Buthus caucasicus intermedius* (Birula, 1897) (= *Mesobuthus caucasicus intermedius*); *Buthus caucasicus przewalskii* (Birula, 1897) (= *Meso-buthus caucasicus przewalskii*); and *Buthus eupeus mongolcus* (Birula, 1911) (= *Mesobuthus eupeus mongolcus*). Birula's descriptions were backward and without normal figures. Birula (1904) described a new subspecies, *Buthus confucius hainanensis* (= *Mesobuthus martensii hainanensis*), based on a single specimen from Hainan Island. Wu (1936) identified two families, four genera and four species of scorpions from China. Kishida (1939) recorded Chinese scorpions and re-described *Buthus martensii*, however, his figures were very inaccurate, and he provided no characteristics of trichobothria. Takashima (1942, 1948, 1951) listed scorpions from Hainan and Shanxi, while Stahnke (1967) re-described *Mesobuthus eupeus mongolcus*, but did not compare this species with its close relatives. Song et al (1982) finished his research on external morphology, reproductive system, life habits and development of *Buthus martensii*. Kovařík (1994) named a new subspecies of *Scorpiops hardwickii* (Gervais, 1843): *Scorpiops hardwickii jendeki* Kovařík, 1994, and later elevated this subspecies to species in his revision of the family Scorpiopidae ((transferred to Euscorpiidae by Sologlad & Sissom (2001)).

Though an impressive compilation of earlier scholarship, there were many limitations to these earlier studies. Thankfully, the number of research papers on Chinese scorpions increased gradually since the early 21st Century. Zhu et al (2004) published a list of Chinese scorpions based on Fet et al (2000) and Kishida (1939) mainly, listing five families, nine genera, 19 species and subspecies. Qi et al (2004) re-described *Mesobuthus martensii martensii* (Karsch, 1879) and provided the detailed study history of this species. Qi et al (2005) published the first comprehensive report on scorpions from Tibet, the authors discovered eight new species belonging to the family Chaerilidae (with one monotypic genus *Chaerilus*) and the family Euscorpiidae (the genera *Euscorpiops* and *Scorpiops*): *Chaerilus tessellatus* Qi, Zhu & Lourenço, 2005; *Euscorpiops karschi* Qi, Zhu & Lourenço, 2005; *Euscorpiops shidian*

Qi, Zhu & Lourenço, 2005; *Euscorpiops vachoni* Qi, Zhu & Lourenço, 2005; *Scorpiops atomatus* Qi, Zhu & Lourenço, 2005; *Scorpiops langxian* Qi, Zhu & Lourenço, 2005; *Scorpiops luridus* Qi, Zhu & Lourenço, 2005; and *Scorpiops pococki* Qi, Zhu & Lourenço, 2005. *Scorpiops petersii* Pocock, 1893 also appeared in this paper which followed on the report of Kishida (1939). Lourenço et al (2005) identified two new species: *Mesobuthus songi* Lourenço, Qi & Zhu, 2005 (= *Hottentotta songi* (Lourenço et al, 2005)), and *Heterometrus tibetanus* Lourenço, Qi, & Zhu, 2005 (belonging to the family Scorpionidae). Shi & Zhang (2005) summarized the research history of taxonomy of the family Buthidae and listed eight species and subspecies, four genera in China: *Mesobuthus martensii* (Karsch, 1879); *Mesobuthus eupeus* (Koch, 1839) (with two subspecies: *Mesobuthus eupeus mongolicus* (Birula, 1911); *Mesobuthus eupeus thersites* (C. L. Koch, 1839)); *Mesobuthus caucasicus* (Nordmann, 1840) (with two subspecies: *Mesobuthus caucasicus intermedius* (Birula, 1897); and *Mesobuthus caucasicus przewalskii* (Birula, 1897)); *Lychas mucronatus* (Fabricius, 1798); *Isometrus maculatus* (DeGeer, 1778); and *Orthochirus scrobiculosus* (Grube, 1873). Lourenço & Qi (2006) described one new genus and new species based on specimens from Tibet: *Tibetiomachus* Lourenço & Qi, 2006 (belonging to the family Hemiscorpiidae), *Tibetiomachus himalayensis* Lourenço & Qi, 2006. Bastawade (2006) reported two new species and four new records based on specimens from Zangnan (the South Tibet, China): *Chaerilus dibangvalleycus* Bastawade, 2006; *Chaerilus pictus* (Pocock, 1890); *Chaerilus tricostatus* Pocock, 1899; *Euscorpiops asthenurus* (Pocock, 1900); *Euscorpiops kamengensis* Bastawade, 2006; and *Scorpiops leptochirus* Pocock, 1893. Zhu et al (2007) described a new species of *Euscorpiops* from Yunnan: *Euscorpiops yangi* Zhu, Zhang & Lourenço, 2007. Shi et al (2007) reported the geographical distribution of two species of *Mesobuthus*: *Mesobuthus eupeus* and *Mesobuthus martensii*. Zhu et al (2008) summarized the chaerilid scorpions of China, and provided the re-descriptions for *Chaerilus tessellatus* Qi, Zhu & Lourenço, 2005 and *Chaerilus triznai* Kovařík, 2000. They also pointed out that *Chaerilus pictus* (Pocock, 1890), which was described by Qi, Zhu & Lourenço (2005) was misidentified and described it as one new species: *Chaerilus conchiformis* Zhu, Han & Lourenço, 2008. Lourenço & Zhu (2008) discovered a new species

belonging to *Isometrus* (belonging to the family Buthidae): *Isometrus (Reddyanus) tibetanus* Zhu & Lourenço, 2008. Di & Zhu (2009a, b) described two new species: *Scorpiops lhasa* Di & Zhu, 2009; and *Chaerilus mainlingensis* Di & Zhu, 2009. Zhang & Zhu (2009) analysed the morphological variation of *Mesobuthus martensii* (Karsch, 1879) from Northern China and found that although this species is widespread in northern China, its morphology does not vary significantly and the variation both in males and females is below species level. Di & Zhu (2009c) described the male of *Euscorpiops karschi* for the first time. Di et al (2009) analysed the genus *Chaerilus* Simon, 1877 of China, with a description of the female *Chaerilus tricostatus* Pocock, 1899 for the first time (*Chaerilus assamensis* Kraepelin, 1913 was an erroneous record in this paper). Di & Zhu (2010) provided a redescription of *Scorpiops margerisonae* Kovařík, 2000 and described the female for the first time. Teruel & Rein (2010) transferred *Mesobuthus songi* Zhu, Qi & Lourenço, 2005 to the genus *Hottentotta* (belonging to the family Buthidae): *Hottentotta songi* (Zhu, Qi & Lourenço, 2005). Sun et al (2010) reported a new species of the genus *Mesobuthus* from Xinjiang, China: *Mesobuthus bolensis* Sun, Zhu & Lourenço, 2010, redescribed the characters of *Mesobuthus songi* Lourenço, Qi & Zhu, 2005, and transferred it to genus *Hottentotta* too. Di et al (2010a, b) discovered two new species in Yunnan: *Euscorpiops validus* Di, Cao, Wu & Li, 2010; *Euscorpiops puerensis* Di, Wu, Cao, Xiao & Li, 2010. Sun & Zhu (2010a, b) described new species of the

genera *Euscorpiops* and *Mesobuthus* from Yunnan and Xinjiang, China: *Euscorpiops xui* Sun & Zhu, 2010, and *Mesobuthus longichelus* Sun & Zhu, 2010. Lourenço et al (2010) published a new record genus and a new species in China: *Razianus xinjianganus* Lourenço, Sun & Zhu, 2010. Di et al (2011a) reviewed the scorpions from Yunnan, and described a new record species to China: *Euscorpiops kubani* Kovařík, 2004. Di et al (2011b) recorded the *Scorpiops* species distributing in Central China, and provided a key for the Chinese species of the genus. Sun & Sun (2011) reviewed the genus *Mesobuthus* (Scorpiones: Buthidae) in China, and described a new species: *Mesobuthus karshius* Sun & Sun, 2011. Recently, Kovařík (2012a) reported five new species of genus *Chaerilus*, including one new species from China: *Chaerilus wrzecionkoi* Kovařík, 2012. Di et al (2013a) reviewed the scorpions from Hainan Island and listed five species belonging to four genera of two families. Di & Fet (2012, personal communication) questioned the validity of *Mesobuthus martensii hainanensis* in Hainan. Di et al (2013b) analysed the scorpions from Tibet: 26 species of seven genera of five families were recorded in Tibet, all with distribution in south and the north shore of Yarlung Zangbo Jiang.

Given both the earlier literature and the most recent and detailed study, the total count for scorpions in China includes 53 species of 12 genera of five families. Particularly worth noting at four scorpion genera with higher diversity in China than elsewhere: *Chaerilus* (8), *Euscorpiops* (11), *Mesobuthus* (9) and *Scorpiops* (11) with more species.

Taxonomy

Phylum: Arthropoda

Subphylum: Chelicerata

Class: Arachnida

Order: Scorpionida

Family Buthidae C. L. Koch, 1837

Buthidae: Fet & Lowe, 2000: 54–57; Söleglad & Fet, 2003: 89–91.

Chinese members: six genera, 18 species and subspecies.

Distribution in China (provinces): most areas of China, except Guangdong, Guizhou, Heilongjiang, Hunan, Jiangxi, Jilin, Sichuan, Zhejiang.

Genus *Hottentotta* Birula, 1908

Hottentotta: Fet & Lowe, 2000: 134–135; Kovařík, 2007: 2–3, 8–10; Sun et al, 2010: 40.

Chinese members: two species.

Distribution in China: Tibet.

1. *Hottentotta alticola* (Pocock, 1895)

Hottentotta (Hottentotta) alticola minusalta: Fet & Lowe, 2000: 135.

Hottentotta (Hottentotta) alticola nigrifrons: Fet & Lowe, 2000: 136.

Hottentotta alticola: Kovařík, 2007: 10, figs. 21–22; Zhu et al, 2004: 111.

Distribution in China: Unknown.

2. *Hottentotta songi* (Lourenço, Qi & Zhu, 2005)

Mesobuthus songi Lourenço, Qi & Zhu, 2005: 3–8, figs. 1–17, tab. 1.

Hottentotta songi: Teruel & Rein, 2010: 7; Sun et al, 2010: 40–42, figs. 25–29.

Distribution: Tibet (endemic (unique to China)).

Genus *Isometrus* Ehrenberg, 1828

Isometrus: Thorell, 1876: 8; Sissom, 1990: 101; Fet & Lowe, 2000: 146; Kovařík, 2003: 1–2.

Chinese members: two subgenera, three species.

Distribution in China: Hainan, Taiwan, Tibet.

Subgenus *Isometrus* Ehrenberg, 1829

Isometrus (Isometrus): Fet & Lowe, 2000: 146; Kovařík, 2003: 2.

Chinese members: one species.

Distribution in China: Hainan, Taiwan, Tibet.

3. *Isometrus (Isometrus) maculatus* (DeGeer, 1778)

Isometrus (Isometrus) maculatus: Fet & Lowe, 2000: 147; Kovařík, 2003: 2–4; Zhu et al, 2004: 112.

Di et al, 2013a: 4, 7, figs. 1–29, tabs. 1–2.

Isometrus (Raddyanus) europaeus: Tikader & Bastawade, 1983: 286–292, figs. 824–840.

Distribution: Hainan, Taiwan.

Subgenus *Reddyanus* Vachon, 1972

Isometrus (Reddyanus): Fet & Lowe, 2000: 151; Kovařík, 2003: 5.

Chinese members: two species.

Distribution in China: Hainan, Tibet.

4. *Isometrus (Reddyanus) hainanensis* Lourenço, Qi & Zhu, 2005

Isometrus (Reddyanus) hainanensis Lourenço, Qi & Zhu, 2005: 58, 60, figs. 1–15, 17–19, tab. 1.

Distribution in China: Hainan (endemic).

5. *Isometrus (Reddyanus) tibetanus* Zhu & Lourenço, 2008

Isometrus (Reddyanus) tibetanus Zhu & Lourenço, 2008: 268–270, figs. 14–26, 32, tab. 1.

Distribution in China: Tibet (endemic).

Genus *Lychas* C. L. Koch, 1845

Lychas: Tikader & Bastawade, 1983: 40; Kovařík, 1997: 312–314.

Chinese members: two species.

Distribution in China: Guangxi, Hainan, Shanghai (?), Yunnan.

6. *Lychas mucronatus* (Fabricius, 1798)

Lychas mucronatus: Kovařík, 1997: 341–344, figs. 10, 12, 29, 31, 80–82, 93, 98; Fet & Lowe, 2000: 164–165; Zhu et al, 2004: 112; Di et al, 2013a: 7, 12, 15, figs. 48–69, tabs. 1–2.

Distribution in China: Guangxi, Hainan, Yunnan.

7. *Lychas scutilus* C. L. Koch, 1845

Lychas scutilus: Kovařík, 1997: 351–352, figs. 41, 47–76, tabs. 1–3; Fet & Lowe, 2000: 166; Zhu et al, 2004: 112.

Distribution in China: Shanghai?

Comments: Fet et al (2000) recorded this species' distribution in China, but they questioned its authenticity. Kovařík & Whitman (2004) reported *Lychas scutilus* C.L. Koch, 1845 in China, and provided the information: “1 ♀ (591), Cina: Shanghai, 1878, d. [G.] Branchi. Note: È la prima segnalazione per la Cina”.

Genus *Mesobuthus* Vachon, 1950

Mesobuthus Vachon 1950: Sun, Zhu & Lourenço, 2010: 35.

Chinese members: nine species and subspecies.

Distribution in China: the north of the Changjiang River (except Heilongjiang and Jilin), Hainan (?), Shanghai (?).

8. *Mesobuthus bolensis* Sun, Zhu & Lourenço, 2010

Mesobuthus bolensis Sun, Zhu & Lourenço, 2010: 36, 40, figs. 2–3, 5–11, 14–18, 21–22, tab. 1; Sun & Sun, 2011: 59–60.

Distribution in China: Xinjiang (endemic).

9. *Mesobuthus caucasicus intermedius* (Birula, 1897)

Mesobuthus caucasicus intermedius: Shi & Zhang, 2005: 475; Sun & Zhu, 2010b: 3–4, 7–8, figs. 2, 11–13; Sun & Sun, 2011: 61–63, figs. 3–4, tab.1.

Olivierus caucasicus intermedius: Fet & Lowe, 2000:191; Zhu et al, 2004:113.

Distribution in China: Xinjiang.

10. *Mesobuthus caucasicus przewalskii* (Birula, 1897)

Mesobuthus caucasicus przewalskii: Shi & Zhang, 2005: 475; Sun & Zhu, 2010b: 4–5, 7–8, figs. 3, 14–16; Sun & Sun, 2011: 60–61, figs. 1–2, tab. 1.

Olivierus caucasicus przewalskii (Birula): Fet & Lowe, 2000: 192; Zhu et al, 2004: 113.

Distribution in China: Xinjiang.

11. *Mesobuthus eupeus mongolicus* (Birula, 1911)

Mesobuthus eupeus mongolicus: Sun & Sun, 2011: 67–70, figs. 7–8, tab. 1; Zhu et al, 2004: 112.

Distribution in China: Gansu, Inner Mongolia (Neimenggu), Ningxia.

12. *Mesobuthus eupeus thersites* (C. L. Koch, 1839)

Mesobuthus eupeus thersites: Sun & Sun, 2011: 70–72, tab. 1; Zhu et al, 2004: 113.

Distribution in China: Xinjiang.

13. *Mesobuthus karshius* Sun & Sun, 2011

Mesobuthus karshius: Sun & Sun, 2011: 63–67, figs. 5–6, tab.1.

Distribution in China: Xinjiang (endemic).

14. *Mesobuthus longichelus* Sun & Zhu, 2010

Mesobuthus longichelus Sun & Zhu, 2010b: 5–10, figs. 1, 4–10, 17–21.

Distribution in China: Xinjiang (endemic).

15. *Mesobuthus martensii martensii* (Karsch, 1879)

Buthus martensi Karsch: Kraepelin, 1899: 25–26; Wu, 1936: 115–117, fig. 1; Song et al, 1982: 22–25, figs. 1–7; Song, 1998: 508, fig. 30: 1.

Mesobuthus martensii (Karsch): Kovařík, 1998: 115; Shi & Zhang, 2005: 474; Shi et al, 2007: 216–223, figs. 1–3, tab. 1; Zhang & Zhu, 2009: 1–17, tabs. 1–8; Sun & Zhu, 2010b: 10.

Mesobuthus martensii martensii (Karsch): Fet & Lowe, 2000: 178; Qi et al, 2004: 137–143, figs. 1–19, tab. 1; Zhu et al, 2004: 113; Sun & Sun, 2011, 72, fig. 9, tab. 1.

Distribution in China: Shi et al (2007) summarized the distribution range of *M. martensii* (Karsch, 1879) in China as the south side of N43° and the north side of the Yangtze River, bordered by the Helan Mountains and the Tengger and Mo Us sand desert in the west and limited by the sea in the east. Kovařík & Whitman (2004) reported *Mesobuthus martensii* (Karsch, 1879) distributed in Shanghai, and the simple information as follows: “2♀♀ (592), Cina: Shanghai, 1878, d. [G.] Branchi”.

16. *Mesobuthus martensii hainanensis* (Birula, 1904)

Buthus confucius hainanensis Birula, 1904: 27; Sun & Sun, 2011: 72.

Mesobuthus martensii hainanensis: Fet et al, 2000: 178; Zhu et al, 2004: 113.

Distribution in China: Hainan (endemic ?).

Comments: Di & Fet questioned the facticity of its distribution in Hainan (2012, personal communication). Birula (1904) described this subspecies, but did not provide a detailed description or illustrations; he did not report the gender or discuss relationship between this species and nominotypic *M. martensii*. We have not found any *Mesobuthus* species in Hainan, and we question the authenticity of this record. Birula (1904) studied specimens labeled “Hainan” and collected by Alfred Otto Herz (St. Petersburg, Russia) who conducted entomological expeditions in the 1890s to China, Korea, Japan, and Siam. Herz did indeed visit Hainan; his collections of reptiles and insects from Hainan were among the first ever made by Europeans. Therefore, *M. martensii hainanensis* could be a case of a mistaken label. Shi et al (2007) confirmed the distribution range of *M. martensii* in China by extensive field surveys and predictive models, and considered this species restricted to latitudes south of N43° and the north side of the Yangtze River, bordered by the Helan Mountains and the Tengger and Mo Us sand desert in the west and limited by the sea in the east.

Genus *Orthochirus* Karsch, 1891

Orthochirus: Kovařík, 2004c; Lourenço & Leguin, 2011a & b.

Chinese members: one species.

Distribution in China: general northwest.

17. *Orthochirus scrobiculosus* (Grube, 1873)

Orthochirus scrobiculosus: Shi & Zhang, 2005: 475.

Distribution in China: general northwest.

Genus *Razianus* Farzanpay, 1987

Razianus: Lourenço et al, 2010: 307–308.

Chinese members: one species.

Distribution in China: Xinjiang.

18. *Razianus xinjianganus* Lourenço, Sun & Zhu, 2010

Razianus xinjianganus Lourenço, Sun & Zhu, 2010: 308–309, 311–312, figs. 1–2, tab. 1.

Distribution in China: Xinjiang (endemic).

Family Chaerilidae Pocock, 1893

Chaerilidae: Fet, 2000a: 323. Kovařík, 2000a: 40–41; Soleglad & Fet, 2003: 92.

Chinese members: one genus, eight species.

Distribution in China: Tibet.

Genus *Chaerilus* Simon, 1877

Chaerilus: Fet, 2000: 323; Kovařík, 2000a: 38; Kovařík, 2005: 1; Qi et al, 2005: 29; Lourenço & Zhu, 2008: 462.

Chinese members: eight species.

Distribution in China: Tibet.

19. *Chaerilus conchiformis* Zhu, Han & Lourenço, 2008

Chaerilus pictus: Qi et al, 2005: 34, 38, figs.126–144.

Chaerilus conchiformis Zhu, Han & Lourenço, 2008: 38–42, figs.1–29, tab.1.

Distribution in China: Tibet (endemic).

20. *Chaerilus dibangvalleyicus* Bastawade, 2006

Chaerilus dibangvalleyicus Bastawade, 2006: figs. 1–16.

Distribution in China: Tibet (endemic).

21. *Chaerilus mainlingensis* Di & Zhu, 2009

Chaerilus mainglingensis Di & Zhu, 2009b: 97–98, 101, figs. 1–16.

Distribution in China: Tibet (endemic).

22. *Chaerilus pictus* (Pocock, 1890)

Chaerilus pictus: Fet, 2000: 327; Kovařík, 2000a: 53–54; figs. 21–22, 39, 42–43, tabs. 1–2; Zhu et al, 2004: 113–114.

Distribution in China: Tibet.

23. *Chaerilus tessellatus* Qi, Zhu & Lourenço, 2005

Chaerilus tessellatus Qi, Zhu & Lourenço, 2005: 30, 34, figs. 109–125; Zhu et al, 2008: 44, 47, figs. 30–44, tab. 1.

Distribution in China: Tibet (endemic).

24. *Chaerilus tricostatus* Pocock, 1899

Chaerilus tricostatus: Fet, 2000a: 327; Kovařík, 2000a: 61–62, figs. 27–28, tabs. 1–2; Di et al, 2009: 133, 136–137, figs. 1–18, tab. 1.

Distribution in China: Tibet.

25. *Chaerilus tryznai* Kovařík, 2000

Chaerilus tryznai Kovařík, 2000a: 65–66, figs. 32–33, tabs. 1–2.

Chaerilus tryznai: Zhu et al, 2008: 47–48, 50–51, figs. 45–59, tab. 1.

Distribution in China: Tibet (endemic).

26. *Chaerilus wrzecionkoi* Kovařík, 2012

Chaerilus wrzecionkoi Kovařík, 2012b: 11, 13, figs. 62–77.

Distribution: Tibet (endemic).

Family Euscorpiidae Laurie, 1896

Euscorpiidae: Fet & Sissom, 2000: 355; Soleglad & Fet, 2003: 105.

Scorpionidae: Fet, 2000c: 487; Kovařík, 2000b: 154.

Chinese members: two genera, 22 species.

Distribution in China: Yunnan, Tibet.

Genus *Euscorpiops* Vachon, 1980

Euscorpiops: Fet & Sissom, 2000: 488; Kovařík, 2000b: 154; Kovařík, 2005: 1, 4; Kovařík, 2012a: 1, 3.

Chinese members: 11 species.

Distribution in China: Yunnan, Tibet.

27. *Euscorpiops asthenurus* (Pocock, 1900)

Euscorpiops asthenurus: Fet & Sissom, 2000: 488.

Scorpiops asthenurus: Kovařík, 2000b: 167, figs. 15, 28, 31, tabs. 1–3.

Distribution in China: Tibet.

28. *Euscorpiops kamengensis* Bastawade, 2006

Euscorpiops kamengensis Bastawade, 2006: 454, 456–457, figs. 17–26.

Distribution in China: Tibet (endemic).

29. *Euscorpiops karschi* Qi, Zhu & Lourenço, 2005

Euscorpiops karschi Qi, Zhu & Lourenço, 2005: 25, figs. 94–108; Di & Zhu, 2009c: 11, 14–15, figs. 1–27, tab. 1.

Distribution in China: Tibet (endemic).

30. *Euscorpiops kubani* Kovařík, 2004

Euscorpiops kubani Kovařík, 2004a: 14–16, figs. 1–6, tab. 1.

Euscorpiops kubani: Di et al, 2011a: 5–9, figs. 10–28, tabs. 1–2.

Distribution in China: Yunnan.

31. *Euscorpiops novaki* Kovařík, 2005

Euscorpiops novaki Kovařík, 2005: 4, 6, figs. 8, 11, 15–16, tab. 1.

Distribution in China: Tibet (endemic).

32. *Euscorpiops puerensis* Di, Wu, Cao, Xiao & Li, 2010

Euscorpiops puerensis Di et al, 2010b: 49–52, 54, 56, 58–59, figs. 1–34, tabs. 1–2.

Euscorpiops puerensis: Di et al, 2011a: 9, 12–15, figs. 29–49.

Distribution in China: Yunnan (endemic).

33. *Euscorpiops shidian* Qi, Zhu & Lourenço, 2005

Euscorpiops shidian Qi, Zhu & Lourenço, 2005: 18, 22, 25, figs. 78–93.

Euscorpiops shidian: Di et al, 2011a: 9, 15–19, figs. 50–68, tabs. 1–2.

Distribution in China: Yunnan (endemic).

34. *Euscorpiops vachoni* Qi, Zhu & Lourenço, 2005

Euscorpiops vachoni Qi, Zhu & Lourenço, 2005: 18, figs. 62–77.

Euscorpiops vachoni: Di et al, 2011a: 19–21, figs. 69–72.

Distribution in China: Yunnan (endemic).

35. *Euscorpiops validus* Di, Cao, Wu & Li, 2010

Euscorpiops validus Di et al, 2010: 14–17, 19, 21, figs. 1–32, tabs. 1–2.

Euscorpiops validus Di et al, 2011: 21, figs. 73–91.

Distribution in China: Yunnan (endemic).

36. *Euscorpiops xui* Sun & Zhu, 2010

Euscorpiops xui Sun & Zhu, 2010: 62, 67, figs. 1–14, tab. 1.

Euscorpiops xui: Di et al, 2011a: 21–25, figs. 92–110, tabs. 1–2.

Distribution in China: Yunnan (endemic).

37. *Euscorpiops yangi* Zhu, Zhang & Lourenço, 2007

Euscorpiops yangi Zhu et al, 2007: 20–22, 25, figs. 1–22, tab. 1.

Euscorpiops yangi: Di et al, 2011a: 26–28, figs. 111–117.

Distribution in China: Yunnan (endemic).

Genus *Scorpiops* Peters, 1861

Scorpiops: Fet & Sissom, 2000: 491; Kovařík, 2000b: 162, 164, 166; Qi et al, 2005: 2; Di & Zhu, 2009a: 40; Di et al, 2011b, 1–2. Kovařík, 2009: 1.

Chinese members: 11 species.

Distribution in China: Hubei, Yunnan, Tibet.

38. *Scorpiops atomatus* Qi, Zhu & Lourenço, 2005

Scorpiops atomatus Qi, Zhu & Lourenço, 2005: 6, 10, figs. 16–31.

Distribution in China: Tibet (endemic).

39. *Scorpiops hardwickii* (Gervais, 1843)

Scorpiops hardwickii: Kovařík, 2000b: 175–179, figs. 14, 46, 56–57.

Scorpiops hardwickii hardwickii: Fet & Sissom, 2000: 492.

Distribution in China: Tibet.

40. *Scorpiops jendeki* Kovařík, 1994

Scorpiops hardwickii jendeki Kovařík, 1994: 62, figs. 7–13, tab. 1; Fet, 2000: 492.

Scorpiops jendeki: Kovařík, 2000b: 180, 182, figs. 59–60, tabs. 1–3; Di et al, 2013b: 90, 93–94, figs. 119–135, tab. 3.

Distribution in China: Yunnan (endemic).

41. *Scorpiops langxian* Qi, Zhu & Lourenço, 2005

Scorpiops langxian Qi, Zhu & Lourenço, 2005: 10, 14, figs. 32–46.

Distribution: Tibet (endemic).

42. *Scorpiops leptochirus* Pocock, 1893

Scorpiops leptochirus Pocock, 1893: Fet & Sissom, 2000: 493.

Distribution: Tibet.

43. *Scorpiops lhasa* Di & Zhu, 2009

Scorpiops lhasa Di & Zhu, 2009a: 40–41, 45, 47, figs. 1–33, tab. 1.

Distribution: Tibet (endemic).

44. *Scorpiops luridus* Qi, Zhu & Lourenço, 2005

Scorpiops luridus Qi, Zhu & Lourenço, 2005: 2, 6, figs. 1–15.

Distribution: Tibet (endemic).

45. *Scorpiops margerisonae* Kovařík, 2000

Scorpiops margerisonae Kovařík, 2000b: 189, figs. 66, 70, tabs. 1–3; Di & Zhu, 2010: 1–8, figs. 1–23, tabs. 1–2.

Distribution: Tibet (endemic).

46. *Scorpiops petersii* Pocock, 1893

Scorpiops petersii: Kovařík, 2000b: 192–194, figs. 35, 42, tabs. 1–3; Fet & Sissom, 2000: 494.

Distribution: Tibet.

47. *Scorpiops pococki* Qi, Zhu & Lourenço, 2005

Scorpiops pococki Qi, Zhu & Lourenço, 2005: 14, figs. 47–61.

Distribution: Tibet (endemic).

48. *Scorpiops tibetanus* Hirst, 1911

Scorpiops tibetanus Hirst, 1911: 472–473; Kovařík, 2000b: 197, figs. 47, 68–69, tab.1–3; Fet & Sissom, 2000: 495; Di et al, 2013b: 75, 77, 80–81, 83, 85, figs. 102–118, tab. 2.

Distribution: Tibet (endemic).

Family Hemiscorpiidae Pocock, 1893

Chinese members: two genera, two species.

Distribution in China: Hainan, Tibet.

Genus *Liocheles* Sundevall, 1833

Liocheles: Monod & Volschenk, 2004: 677.

Chinese members: one species.

Distribution in China: Hainan.

49. *Liocheles australasiae* (Fabricius, 1775)

Hormurusaustra lasiae: Wu, 1936: 121–123, fig. 4; Tikader & Bastawade, 1983: 501–505, figs. 1362–1375.

Liocheles australasiae: Monod & Volschenk, 2004: 677; Di et al, 2013a: 15–16, 19–21, figs. 70–88, tabs. 1–2.

Liocheles australasiae australasiae: Fet, 2000b: 397.

Distribution: Hainan.

Genus *Tibetiomachus* Lourenço & Qi, 2006

Tibetiomachus Lourenço & Qi, 2006: 291.

Chinese members: one species.

Distribution in China: Tibet.

50. *Tibetiomachus himalayensis* Lourenço & Qi, 2006

Tibetiomachus himalayensis Lourenço & Qi, 2006: 291, 293–294, figs. 1: 5–14, 2: 5–26.

Distribution: Tibet (endemic).

Family Scorpionidae Latreille, 1802

Scorpionidae: Fet, 2000c: 427–428; Soleglad & Fet, 2003: 113–114.

Chinese members: one genus, three species.

Distribution in China: Tibet.

Genus *Heterometrus* Ehrenberg, 1828

Heterometrus: Fet, 2000d: 431; Kovařík, 2004b: 2, 4; Lourenço et al, 2005: 9.

Chinese members: three species.

Distribution in China: Tibet (?)

51. *Heterometrus longimanus* (Herbst, 1800)

Heterometrus petersii (Thorell, 1876): Zhu et al, 2004: 114.

Distribution in China: some indeterminate parts.

52. *Heterometrus tibetanus* Lourenço, Qi & Zhu, 2005

Heterometrus tibetanus Lourenço, Qi & Zhu, 2005: 10–14, figs. 18–34, tab. 1.

Distribution: Tibet (endemic).

53. *Heterometrus petersii* (Thorell, 1876)

Heterometrus petersii (Thorell, 1876): Zhu et al, 2004: 114.

Distribution in China: some indeterminate parts.

Key to scorpions from China (Di & Zhu, 2010; Di et al, 2009, 2010, 2011a, 2011b, 2013a, 2013b; Monod & Volschenk, 2004; Kovařík, 1997, 2000a, 2000b, 2003, 2004a, 2004b, 2005, 2007, 2009, 2012a, 2012b; Prendini, 2000; Soleglad & Fet, 2003; Sun et al, 2010; Sun & Sun, 2011; Zhu & Lourenço, 2005):

1. Orthobothriotaxic pattern type A; ventral aspect of leg tarsus with multiple irregular rows of setae, no trace of spinules (*configuration* 2); dorsal edge of cheliceral movable finger with two basal denticles; hemispermatophore is *flagelliform* (**Buthidae**) 2
1. Orthobothriotaxic pattern type B or C; ventral aspect of leg tarsus with or without irregular setal rows, spinules present medially; dorsal edge of cheliceral movable finger with a *single* basal denticle; hemi-spermatophore is either *fusiform* or *lamelliform* 19
2. Tibial spurs absent on all legs (**Isometrus**) 3
2. Tibial spurs present on legs III & IV 5
3. Trichobothrium *db* of the fixed finger in a distal position in relation to the trichobothria *et* and *est*; the distance between external trichobothria of the femur, e_1 and e_2 being at least two to five times the distance between trichobothria e_1 and d_3 of the femur (**subgenus Isometrus**); telson with 2 granules on the ventral surface *Isometrus maculatus*
3. Trichobothrium *db* in a basal position to *et*, situated between *et* and *est*; the distance between external trichobothria of the femur, e_1 and e_2 always less than two times the distance between e_1 and d_3 (**subgenus Reddyanus**) 4
4. Telson with 5 granules on the ventral surface *Isometrus hainanensis*
4. Telson with 2 granules on the ventral surface *Isometrus tibetanus*
5. Telson with subaculear tooth (**Lychas**) 6
5. Telson without subaculear tooth 7
6. In adults, total length of males longer than females, metasomes of males notable elongated *Lychas scutillus*
6. In adults, with similar body and metasome length in both sexes (or males small) *Lychas mucronatus*
7. Metasome with punctuate (**Orthochirus**) *Orthochirus scrobiculosus*
7. Metasome with granule or smooth 8
8. In adults, with smaller body (usually shorter than 20mm), carinae of prosome and mesosome weak (**Razianus**)
..... *Razianus xinjianganus*
8. Medium to large body length (over 30 mm), carinae of prosome and mesosome strong 9
9. Ventrolateral carinae of fifth metasomal segment with all granules more or less equal in size and never lobate (**Hottentotta**) 10
9. Ventrolateral carinae of Metasoma segment V formed of disjunct and unequal granules, often enlarged posteriorly (**Mesobuthus**) 11
10. Movable finger of pedipalp-chela with distinct granules divided into 14–16 rows, pedipalps without intense setation *Hottentotta alticola*
10. Movable finger of pedipalp-chela with distinct granules divided into 13 rows, body and pedipalps covered with very intense setation *Hottentotta songi*

11. Ventrolateral carinae of segment V on metasoma with several markedly large and extroversive lobed granules 12
11. Ventrolateral carinae of segment V on metasoma without markedly large and extroversive lobed granules 14
12. Ventral carinae of segment II and III of metasoma gradually stronger posteriorly 13
12. Ventral carinae of segment II and III of metasoma not stronger posteriorly *Mesobuthus longichelus*
13. Anterior margin of carapace with a very weak median concavity, chelae more robust
..... *Mesobuthus eupeus thersites*
13. Anterior margin of carapace with a very weak median projection or approximately straight, chelae relatively less robust *Mesobuthus eupeus mongolicus*
14. Ventral surface of segment V on metasoma without brown pigment 15
14. Ventral surface of segment V on metasoma with markedly brown pigment 16
15. Surfaces of carapace with relatively dense small granules, tarsus of legs with two long longitudinal rows of setae positioned ventrally *Mesobuthus bolensis*
15. Surfaces of carapace between median carinae almost smooth, but the external surfaces with comparatively dense small granules, tarsus of legs with two short longitudinal rows of setae positioned ventrally *Mesobuthus karshius*
16. Dorsal surfaces of metasomal segments I–IV and each surface of segment V with irregular net-like dark pigmentation 17
16. Only surfaces of segment V on metasoma with irregular net-like dark pigmentation, dorsal surfaces of segments I–IV without net-like pigmentation (*Mesobuthus martensii* (Karsch, 1879)) 18
17. Pectinal teeth number 20–25 in females and 26–30 in males; dentate margins of movable and fixed fingers with 12 and 11 oblique rows of granules respectively *Mesobuthus caasicus intermedius*
17. Pectinal teeth numbers 15–19 in females and 19–23 in males; dentate margins of movable and fixed fingers with 11 and 10 oblique rows of granules respectively *Mesobuthus caasicus przewalskii*
18. Distributed on the north side of the Yangtze River *Mesobuthus martensii martensii*
18. Distributed on Hainan Island *Mesobuthus martensii hainanensis*
19. Orthobothriotaxic pattern type B; sternum is type 1; hemispermatophore is fusiform (**Chaerilidae: Chaerilus**) 20
19. Orthobothriotaxic pattern type C; sternum is type 2; hemispermatophore is lamelliform 27
20. Movable finger of pedipalp with 7–8 rows of granules 21
20. Movable finger of pedipalp with 10–14 rows of granules 25
21. Chela length to width ratio in adults 1.6–1.8 *Chaerilus conchiformus*
21. Chela length to width ratio in adults higher than 2.0 22
22. Ventral side of seventh mesosomal segment with 2 pair of granular carina, anterior margin straight with a median notch 23
22. Ventral side of seventh mesosomal segment with many granules but without carina, anterior margin straight without median notch 24
23. Pedipalp femur short than carapace; 8–9 minute teeth on inner ventral margins of movable and immovable fingers respectively *Chaerilus dibangvalleycus*
23. Pedipalp femur longer than carapace, 7–8 minute teeth on inner ventral margins of movable and immovable fingers respectively *Chaerilus mainlingensis*
24. Manus of pedipalp in male narrow and long. Chela length/width ratio in male higher than 3
..... *Chaerilus tryznai*
24. Manus of pedipalp in male robust (Fig. 68). Chela length/width ratio in adults lower than 2.6
..... *Chaerilus wrzecionkoi*
25. Movable finger of pedipalp with 13–14 rows of granules; telson of male rather long and about 4.7 times greater length than width, with obvious sexual dimorphism *Chaerilus pictus*
25. Movable finger of pedipalp with 11–12 rows of granules, telson of male and female without sexual dimorphism, manus lacks 1 dorsal carina 26
26. Carapace, tergites nearly smooth in adults, chelicerae dorsal aspect without granules *Chaerilus tessellatus*
26. Carapace, tergites with many big granules in adults, chelicerae dorsal aspect with granules *Chaerilus tricostatus*

27. Legs with two pedal spurs (though one or more pedal spurs are lost in many troglobitic species); ventral aspect of leg tarsus equipped with moderately developed setal pairs and/or median row of spinules (configuration 5); paraxial organ without reflection of internobasal sperm duct (**Chactoidea**, see Soleglad & Fet, 2003, p. 92–93: Key to the superfamilies of parvorder **Iurida**); chelal fingers equipped with inner accessory denticles (IAD), outer denticles (OD) situated outside of median denticle (MD) row; major variable neobothriotaxy present, types Eu1 and Eu2; chelal palm is flat in appearance, carinae D3 and V2 essentially obsolete, angle formed by carinae D3:D4:D5 greater than 90° (**Euscorpidae**, see Soleglad & Fet, 2003, p. 94: Key to the families of superfamily **Chactoidea**) 28
27. Legs with one pedal spur (retrolateral spur absent, though this character is reversed in some bothriurid genera); ventral aspect of leg tarsus equipped with pairs of large limbated socketed setae, median spinule row optional (configuration 4); paraxial organ with reflection of internobasal sperm duct (**Scorpionoidea**, see Soleglad & Fet, 2003, p. 92–93: Key to the superfamilies of parvorder **Iurida**) 47
28. Tricho-bothrium *Eb*₃ on external surface of chela is located between trichobothria Dt and Est. Telson vesicle/aculeus juncture with annular ring (**Euscorpions**) 29
28. Trichobothrium *Eb*₃ on the external aspect of pedipalp chela located basally from trichobothrium Dt. Annular ring at vesicle/aculeus juncture absent (**Scorpiops**) 39
29. Number of trichobothria on external surface of pedipalp patella: 19 (5 *eb*, 2 *esb*, 2 *em*, 5 *est*, 5 *et*) 30
29. Number of trichobothria on external surface of pedipalp patella: 17–19 (5–6 *eb*, 1–2 *esb*, 2 *em*, 4 *est*, 5 *et*) 32
30. Pattern of trichobothria on external surface of pedipalp patella: 19 (6 *eb*, 2 *esb*, 2 *em*, 4 *est*, 5 *et*) *Euscorpions xui*
30. Pattern of trichobothria on external surface of pedipalp patella: 19 (5 *eb*, 2 *esb*, 2 *em*, 5 *est*, 5 *et*) 31
31. Number of trichobothria on ventral surface of patella: 7; number of pectinal teeth: 4–5; movable finger longer than carapace and as long as pedipalp femur *Euscorpions kamengensis*
31. Number of trichobothria on ventral surface of patella: 9; pectinal teeth number 8; movable finger as long as carapace and shorter than pedipalp femur *Euscorpions novaki*
32. Male pedipalp chela finers strongly scalloped: with a pronounced lobe on the movable finger and a corresponding notch on fixed finger 33
32. Male pedipalp chela fingers slightly scalloped or straight: lobe and corresponding notch reduced to absent 37
33. Chela length to width ratio higher than 2.9 34
33. Chela length to width ratio lower than 2.9 36
34. Female pedipalp fingers nearly straight *Euscorpions asthenurus*
34. Female pedipalp fingers obviously scalloped 35
35. Chela length to width ratio 2.9–3.2, carapace with dense, minute granules, total length over than 50 mm, coloration basically dark brown *Euscorpions validus*
35. Chela length to width ratio 3.4–3.5, carapace with sparse, nearly equal granules, total length less than 50 mm (small species), coloration basically dark red-brown *Euscorpions karschi*
36. Chela manus short, stout and rounded *Euscorpions vachoni*
36. Chela manus flattened dorsoventrally *Euscorpions puerensis*
37. Chela length to width ratio higher than 3.2 38
37. Chela length to width ratio lower than 3.2 *Euscorpions kubani*
38. Number of trichobothria on ventral surface of patella: 11, chela length to carapace length ratio less than 1.4 *Euscorpions shidian*
38. Number of trichobothria on ventral surface of patella: 10, chela length to carapace length ratio higher than 1.9 *Euscorpions yangi*
39. Fingers of pedipalps are straight or only slightly flexed in both sexes 40
39. Fingers of pedipalps are flexed (or curved) in both sexes 41
40. Ventral trichobothria on patella number 6 (7 rarely), total length 30–42.1 mm, pectinal teeth number 4–5, chela length to width ratio about 2.2 *Scorpiops jendeki*
40. Ventral trichobothria on patella number 7, total length 40–58 mm, pectinal teeth number 7–9, chela length to width ratio about 3.3–3.5 *Scorpiops leptochirus*

41. Male chela length to width ratio about 1.8–2.2; the manus with same or very similar length and width, fingers of pedipalps are very strongly flexed in the male. Ventral trichobothria on patella number 6–8 *Scorpiops hardwickii* “complex” group (including *S. hardwickii*, *S. langxian*, and *S. pococki*)
41. Manus length to width ratio visibly higher than 1.0 42
42. Total length more than 65 mm 43
42. Total length less than 65 mm 44
43. Mostly yellowish to yellow in adults, ventral patella of pedipalps with 9 trichobothria *Scorpiops luridus*
43. Mostly red brown in adults, ventral patella of pedipalps with 7 (rarely 6 or 8) trichobothria *Scorpiops petersii*
44. Dorsally flat manus of pedipalps and chela of both sexes with length/width ratio: 2.1–2.2 (mean about 2.1 in males and 2.2 in females), total length 40.0–50.0 mm in adults *Scorpiops margerisonae*
44. Dorsally round manus of pedipalps or at least the chela of one sex with length to width ratio higher than 2.2 or total length higher than 50 mm 45
45. Total length less than 40 mm 46
45. Total length more than 50 mm, chela strong, with length/width ratio: 2.0 in male and 2.5 in female *Scorpiops tibetanus*
46. Chela of pedipalp length to width ratio about 2.6–3.0, dorsal surface of chela of pedipalp coarse *Scorpiops lhasa*
46. Chela of pedipalp length to width ratio lower than 2.5, dorsal surface of chela of pedipalp smooth with luster *Scorpiops atomatus*
47. Median ocular tubercle of carapace shallow, not raised above carapace surface; 2 pairs of lateral eyes; telotarsus with lateral lobes truncated; *Est* located in middle of hand (**Hemiscorpiidae**, see Stockmann & Ythier, 2010, p. 201) 48
47. Median ocular tubercle raised up; 3 pairs of lateral eyes; telotarsus with lateral margins ending in rounded lobes; *Est* located in distal of hand (**Scorpionidae**, see Stockmann & Ythier, 2010, p. 201); pedipalp femur with three trichobothria; patella of pedipalp with 19 trichobothria, three on ventral and 13 on external surface; chela of pedipalp with 26 trichobothria; retrolateral pedal spurs absent; lateroapical margins of tarsi produced into rounded lobes; metasomal segments I to IV with paired ventral submedian carinae; stridulatory organ located on opposing surfaces of pedipalp coxa and first leg; total length 60 to 180 mm (**Heterometrus**, see Kovařík, 2004, p. 2) 49
48. Chela trichobothrium *dt* present (**Liocheles**) *Liocheles australasiae*
48. Chela trichobothrium *dt* absent (**Tibetiomachus**) *Tibetiomachus himalayensis*
49. Chela round and strong *Heterometrus petersii*
49. Chela narrow and long 50
50. Chela manus with weak granules on the dorso-internal aspect *Heterometrus longimanus*
50. Chela manus with strong spinoid granules on the dorso-internal aspect *Heterometrus tibetanus*

DISCUSSION

The distribution of scorpions is quite well demarcated by very distinct features arising from the geography and topography of China: there are markedly different constituents between east and west, north and south. Only four recorded species can be found living in both East China and Chinese islands (based on recently finished investigations): *Isometrus maculatus* (DeGeer, 1778), *Lychas mucronatus* (Fabricius, 1798), *Liocheles australasiae* (Fabricius, 1775), and *Mesobuthus martensii martensii* (Karsch, 1879); Three species maybe extinct or invalid: *Isometrus hainanensis* Lourenço, Qi & Zhu,

2005, *Lychas scutillus* C. L. Koch, 1845, and *Mesobuthus martensii hainanensis* (Birula, 1904). In Central China, only one species is reported: *Mesobuthus martensii martensii* (Karsch, 1879).

All of the species of genera *Chaerilus*, *Euscorpiops*, *Scorpiops*, and most of *Mesobuthus* live in the west of China, which is unsurprising as Tibet, Yunnan, and Xinjiang are the richest areas of scorpion biodiversity in all of China. Meanwhile, there are only nine species of two genera reported in northern China (including Xinjiang): *Mesobuthus* (eight species) and *Razianus* (one species).

Zhu et al (2004) published a list of Chinese scorpions based on the information of Fet et al (2000) and Kishida (1939), which included 5 families, 9 genera, 19 species and subspecies. Qi et al (2005) discovered eight new species of Chaerilidae (*Chaerilus*) and Euscorpiidae (*Euscorpiops* and *Scorpiops*). Di et al (2011a) reviewed the scorpions of Yunnan, recording eight previously known species and described one new species to China: *Euscorpiops kubani* Kovařík, 2004. Sun & Sun (2011) reviewed the genus *Mesobuthus* (Scorpiones: Buthidae) in China, recorded Nine species and subspecies including a new species: *Mesobuthus karshius* Sun & Sun, 2011. Di et al (2013a, b) reviewed the scorpions from Hainan Island and Tibet: 5 species and 26 species were recorded in Hainan and Tibet respectively.

While not completely exhaustive, this study nonetheless gives an excellent overview on the

general situation of scorpion biodiversity in China. At present, 53 scorpion species of 12 genera of 5 families are recorded in China, which has greatly helped clarify the characters of biodiversity and distribution of scorpions. This overview—including a history of taxonomic research, an updated checklist and key of the scorpions of China—accordingly is aimed at producing a catalogue of scorpions in China. While the survey on scorpion species diversity in China is not completed yet, we hope that this overview can provide researchers in both mainland China and abroad with enough basic data for taxonomy, fauna, and resources to help further their efforts.

Acknowledgements: We are grateful to Prof. Victor Fet, Prof. Wilson R. Lourenço, Mr. Kovařík, and Mr. Jan Ove Rein for providing references.

References

- Bastawade DB. 2006. Arachnida: Scorpionida, Uropygi, Schizomida and Onco podid Opiliones (Chelicerata). Zool. Surv. India. Fauna of Arunachal Pradesh, *State Fauna Series*, **13**(2): 449-465.
- Birula AA. 1897. Miscellanea scorpologica. II Zur Synonymie der russischen Scorpione. *Annuaire du Muse'e Zoologique de l'Acade'mie Imp'e'riale des Sciences de St.-Pe'tersbourg*, **2**: 377-391.
- Birula AA. 1904. Miscellanea scorpologica VI. Ueber einige Buthus-Arten Centralasiens nebst ihrer geographischen Verbreitung. *Annuaire du Muse'e Zoologique de l'Acade'mie Imp'e'riale des Sciences de St.-Pe'tersbourg*, **9**: 20-27.
- Birula AA. 1911. Arachnologische Beiträge I Zur Scorpionen- und Solifugen-Fauna des Chinesischen Reiches. *Revue Russe d'Entomologie*, **11**(2): 195-201.
- Cloudsley-Thompson JL. 1990. Scorpions in mythology, folklore and history. In: Polis GA. *The Biology of Scorpions*. Stanford, CA: Stanford University Press, 462-485.
- Di ZY, Zhu MS. 2009a. One new species of the Genus *Scorpiops* Peters, 1861 (Scorpiones: Euscorpiidae, Scorpiopinae) from Tibet, China. *Zootaxa*, **2030**: 39-48.
- Di ZY, Zhu MS. 2009b. A new species of *Chaerilus* Simon, 1877 (Scorpiones, Chaerilidae) from China. *Acta Arachnologica*, **58**(2): 97-102.
- Di ZY, Zhu MS. 2009c. The male of *Euscorpiops karschi* (Scorpiones: Euscorpiidae, Scorpiopinae) from China (Tibet). *Arthropoda Selecta*, **18**(1/2): 9-16.
- Di ZY, Zhu MS. 2010. Redescription of *Scorpiops margerisonae* Kovařík, 2000, with the first record of its female, from China (Scorpiones: Euscorpiidae: Scorpiopinae). *Euscorpius*, **104**: 1-9.
- Di ZY, Wu YL, Cao ZJ, Fan LQ, Li WX. 2009. The genus *Chaerilus* Simon, 1877 (Scorpiones: Chaerilidae) in China, with a description of the female *C. tricostatus* Pocock, 1899. *Arthropoda Selecta*, **18**(3/4): 131-138.
- Di ZY, Cao ZJ, Wu YL, Li WX. 2010a. A new species of the Genus *Euscorpiops* Vachon, 1980 (Scorpiones: Euscorpiidae, Scorpiopinae) from Yun'nan, China. *Zootaxa*, **2361**: 13-22.
- Di ZY, Wu YL, Cao ZJ, Xiao H, Li WX. 2010b. A catalogue of the genus *Euscorpiops* Vachon, 1980 (Scorpiones: Euscorpiidae, Scorpiopinae) from China, with description of a new species. *Zootaxa*, **2477**: 49-61.
- Di ZY, He YW, Wu YL, Cao ZJ, Liu H, Jiang DH, Li WX. 2011a. The scorpions of Yunnan (China): updated identification key, new record and redescriptions of *Euscorpiops kubani* and *E. shidian* (Arachnida, Scorpiones). *ZooKeys*, **82**: 1-33.
- Di ZY, He YW, Cao ZJ, Wu YL, Li WX. 2011b. The first record of the family Euscorpiidae (Arachnida: Scorpiones) from Central China, with a key of Chinese species of the genus *Scorpiops*. *Euscorpius*, **116**: 1-11.
- Di ZY, Cao ZJ, Wu YL, Zhu L, Liu H, Li WX. 2013a. The scorpions of Hainan Island, China (Arachnida: Scorpiones). *Euscorpius*, **153**: 1-22.
- Di ZY, Xu XB, Cao ZJ, Wu YL, Li WX. 2013b. Notes on the scorpions (Arachnida, Scorpiones) from Tibet with the redescription of *Scorpiops jendeki* Kovařík, 2000 (Scorpiones, Euscorpiidae) from Yunnan (China). *ZooKeys*, **301**: 51-99.
- Fet V. 2000a. Family Chaerilidae Pocock, 1893. In: *Catalog of the Scorpions of the World (1758–1998)*. New York: The New York Entomological Society, 323-328.
- Fet V. 2000b. Family Ischnuridae Simon, 1879. In: *Catalog of the Scorpions of the world (1758–1998)*. New York: The New York Entomological Society, 383-408.

- Fet V. 2000c. Family Scorpionidae Latreille, 1802. In: Catalog of the Scorpions of the world (1758–1998). New York: The New York Entomological Society, 427-486.
- Fet V. 2000d. Family Scorpionidae Kraepelin, 1905. In: Catalog of the Scorpions of the World (1758–1998). New York: The New York Entomological Society, 487-495.
- Fet V, Braunwalder ME, Cameron HD. 2002. Scorpions (Arachnida, Scorpiones) described by Linnaeus. *Bulletin of the British Arachnological Society*, **12**(4): 176-182.
- Fet V, Lowe G. 2000. Family Buthidae C. L. Koch, 1837. In: Catalog of the Scorpions of the world (1758–1998). New York: The New York Entomological Society, 54-286.
- Fet V, Sissom WD. 2000. Family Euscorpionidae Laurie, 1896. In: Sissom FWD, Lowe G, Braunwalder M. Catalog of the Scorpions of the world (1758–1998). New York: The New York Entomological Society, 355-380.
- Fet V, El-Hennawy H, Braunwalder ME, Cloudsley-Thompson JL. 2009. The first observation on scorpion biogeography by Aristotle. *Boletín Sociedad Entomológica Aragonesa*, **44**: 147-150.
- Fet V, Sissom WD, Lowe G, Braunwalder ME. 2000. Catalog of the Scorpions of the World (1758–1998). New York: The New York Entomological Society.
- Hirst S. 1911. Descriptions of new scorpions. *Annals and Magazine of Natural History*, **8**(46): 462-473.
- Kishida K. 1939. Arachnida of Jehol, Scorpiones. In: Report of the first scientific expedition to Mandchoukuo under the leadership of Shigeyasu Tokunaga, 1933, 5, 1, **4**(10): 1-66.
- Kovařík F. 1994. *Scorpiops irenae* sp. n. from Nepal and *Scorpiops hardwickei jendeki* subsp. n. from Yunnan, China (Arachnida: Scorpionida. Vaejovidae). *Acta Societatis Zoologicae Bohemicae*, **58**: 61-66.
- Kovařík F. 1997. Revision of the genera *Lychas* and *Hemilychas*, with descriptions of six new species (Scorpiones, Buthidae)". *Acta Societatis Zoologicae Bohemicae*, **61**: 311-371.
- Kovařík F. 2000a. Revision of family Chaerilidae (Scorpiones), with descriptions of three new species. *Serket*, **7**(2): 38-77.
- Kovařík F. 2000b. Revision of family Scorpionidae (Scorpiones), with descriptions of six new species. *Acta Societatis Zoologicae Bohemicae*, **64**: 153-201.
- Kovařík F. 2003. A review of the genus *Isometrus* Ehrenberg, 1828 (Scorpiones: Buthidae) with descriptions of four new species from Asia and Australia. *Euscorpius*, **10**: 1-19.
- Kovařík F. 2004a. *Euscorpionops kubani* sp. nov. from Laos (Scorpiones, Euscorpionidae, Scorpioninae). *Acta Musei Moraviae, Scientiae biologicae (Brno)*, **89**: 13-18.
- Kovařík F. 2004b. A review of the genus *Heterometrus* Ehrenberg, 1828, with descriptions of seven new species (Scorpiones, Scorpionidae). *Euscorpius*, **15**: 1-60.
- Kovařík F. 2004c. Revision and taxonomic position of genera *Afghanorthochirus* Lourenço & Vachon, *Baloothochirus* Kovařík, *Butheolus* Simon, *Nanobuthus* Pocock, *Orthochiroides* Kovařík, *Pakistanorthochirus* Lourenço, and Asian *Orthochirus* Karsch, with descriptions of twelve new species (Scorpiones, Buthidae). *Euscorpius*, **16**: 1-33.
- Kovařík F. 2005. Three new species of the genera *Euscorpionops* Vachon, 1980 and *Scorpiops* Peters, 1861 from Asia (Scorpiones: Euscorpionidae, Scorpioninae). *Euscorpius*, **27**: 1-10.
- Kovařík F. 2007. A revision of the genus *Hottentotta* Birula, 1908, with descriptions of four new species (Scorpiones: Buthidae). *Euscorpius*, **58**: 1-107.
- Kovařík F. 2012a. *Euscorpionops thaomischi* sp. n. from Vietnam and a key to species of the genus (Scorpiones: Euscorpionidae: Scorpioninae). *Euscorpius*, **142**: 1-8.
- Kovařík F. 2012b. Five new species of *Chaerilus* Simon, 1877 from China, Indonesia, Malaysia, Philippines, Thailand, and Vietnam (Scorpiones: Chaerilidae). *Euscorpius*, **149**: 1-14.
- Kovařík F, Whitman S. 2004. Cataloghi del Museo di Storia Naturale dell'Università di Firenze - Sezione di Zoologia «La Specola». XXII. Arachnida scorpiones. Tipi. Addenda (1988–2004) e checklist della collezione (Euscorpioninae esclusi). *Atti Soc. To sc. Sci. nat., Mem., Serie B*, **111**: 103-119.
- Kraepelin K. 1899. Scorpiones und Pedipalpi. In: *Herausgegeben von der Deutschen zoologischen Gesellschaft*. Berlin: R. Friedlander und Sohn Verlag, 265.
- Lourenço WR, Leguin EA. 2011a. Further Considerations on the species of the genus *Orthochirus* Karsch, 1891 from Africa, with description of three new species (Scorpiones: Buthidae). *Euscorpius*, **123**: 1-19.
- Lourenço WR, Leguin EA. 2011b. One more new species of the genus *Orthochirus* Karsch, 1891 from Africa (Scorpiones: Buthidae). *Euscorpius*, **135**: 1-6.
- Lourenço WR, Qi JX. 2006. Mountain scorpions: a new genus and species from Tibet (China). *Comptes Rendus Biologies*, **329**(4): 289-295.
- Lourenço WR, Qi JX, Zhu MS. 2005. Description of two new species of scorpions from China (Tibet) belonging to the genera *Mesobuthus* Vachon (Buthidae) and *Heterometrus* Ehrenberg (Scorpionidae). *Zootaxa*, **985**: 1-16.
- Lourenço WR, Qi JX, Zhu MS. 2005. Description of a new species of *Isometrus* Ehrenberg 1828 (Scorpiones, Buthidae) from the Island of Hainan, China. *Boletín de la Sociedad Entomológica Aragonesa*, **36**: 57-63.
- Lourenço WR, Sun D, Zhu MS. 2010. *Razianus xinjianganus* sp. nov.: A new record Genus and new species of (Scorpiones, Buthidae) from China. *Journal of Hebei University: Natural Science Edition*, **30**(3): 307-312.
- Lourenço WR, Zhu MS. 2008. A new species of the genus *Isometrus* Ehrenberg 1828 from China (Scorpiones, Buthidae). *Acta Zootaxonomica Sinica*, **33**(2): 264-271.
- Monod L, Volschenk ES. 2004. *Liocheles litodactylus* (Scorpiones: Liochelidae): An unusual new Liocheles species from the Australian wet tropics (Queensland). *Memoirs of the Queensland Museum*, **49**: 675-687.

- Prendini L. 2000. Phylogeny and classification of the superfamily Scorpionoidea Latreille, 1802 (Chelicerata, Scorpiones): An exemplar approach. *Cladistics*, **16**(1): 1-78.
- Qi JX, Zhu MS, Lourenço WR. 2004. Redescription of *Mesobuthus martensii martensii* (Karsch, 1879) (Scorpiones: Buthidae) of China. *Revista Ibérica de Aracnología*, **10**: 137-144.
- Qi JX, Zhu MS, Lourenço WR. 2005. Eight new species of the genera *Scorpiops* Peters, *Euscorpiops* Vachon, and *Chaerilus* Simon (Scorpiones: Euscorpiidae, Chaerilidae) from Tibet and Yunnan, China. *Euscorpius*, **32**: 1-40.
- Shi CM, Zhang DX. 2005. A review of the systematic research on buthid scorpions (Scorpiones: Buthidae). *Acta Zootaxonomica Sinica*, **30**: 470-477.
- Shi CM, Huang ZS, Wang L, He LJ, Hua YP, Leng L, Zhang DX. 2007. Geographical distribution of two species of *Mesobuthus* (Scorpiones, Buthidae) in China: insights from systematic field surveys and predictive models. *The Journal of Arachnology*, **35**(2): 215-226.
- Sissom WD. 1990. Systematics, biogeography and paleontology. In: The Biology of Scorpions. California: Stanford University Press, 64-160.
- Soleglad ME, Fet V. 2003. High-level systematics and phylogeny of the extant scorpions (Scorpiones: Orthosterni). *Euscorpius*, **11**: 1-175.
- Soleglad ME, Sissom WD. 2001. Phylogeny of the family Euscorpiidae Laurie, 1896: a major revision. In: Scorpions 2001. In Memorium Gary A. Polis. Bucks: British Arachnological Society, 25-111.
- Song DX. 1998. Arachnida Scorpiones. In: Pictorial Keys to Soil Animals of China. Beijing: Science Press, 131-132, 507-508.
- Song DX, Lv XY, Shang JW. 1982. Morphology and habits of *Buthus martensii*. *Bulletin of Biology*, **1**: 22-25.
- Stahnke HL. 1967. Scorpiones. Ergebnisse der zoologischen Forschungen von Dr. Z. Kaszab in der Mongolei. 93. Skorpiones. *Reichenbachia*, **9**(6): 59-68.
- Sun D, Sun ZN. 2011. Notes on the genus *Mesobuthus* (Scorpiones: Buthidae) in China, with description of a new species. *The Journal of Arachnology*, **39**(1): 59-75.
- Sun D, Zhu MS. 2010a. One new species of scorpion belonging to the genus *Euscorpiops* Vachon, 1980 from Yunan, China (Scorpiones: Euscorpiidae, Scorpiopinae). *Zootaxa*, **2399**: 61-68.
- Sun D, Zhu MS. 2010b. A new species of the genus *Mesobuthus* Vachon, 1950 (Scorpiones: Buthidae) from Xinjiang, China. *ZooKeys*, **37**: 1-12.
- Sun D, Zhu MS, Lourenço WR. 2010. A new species of *Mesobuthus* (Scorpiones: Buthidae) from Xinjiang, China, with notes on *Mesobuthus songi*. *The Journal of Arachnology*, **38**(1): 35-43.
- Takashima H. 1942. The survey on scorpions and Whip amphipods in East Asia (V). *Acta Arachnologica*, **7**(1): 24-30.
- Takashima H. 1948. Scorpions of Shansi, North China. *Acta Arachnologica*, **10**(3-4): 112-116.
- Takashima H. 1951. Notes on the scorpion of Hainan Island. *Acta Arachnologica*, **12**(3-4): 83-87.
- Teruel R, Rein JO. 2010. A new species of *Hottentotta* Birula, 1908 from Afghanistan, with a note on the generic position of *Mesobuthus songi* Lourenço, Qi & Zhu, 2005 (Scorpiones: Buthidae). *Euscorpius*, **94**: 1-8.
- Tikader BK, Bastawade DB. 1983. The Fauna of India. Vol. 3. Scorpions (Scorpionida: Arachnida). Calcutta: Zoo logical Survey of India.
- Thorell T. 1876. On the classification of Scorpions. *Annals and Magazine of Natural History*, **4** (17): 1-15.
- Wang XL, Chen SM. 2007. The Explaining about How the Words of "Er Ya · Shi Chong" about Name and Thing Were Named. *Journal of Xuzhou Normal University (Philosophy and Social Sciences Edition)*, **133**(3): 65-69.
- Wu HW. 1936. A review of the scorpions and whip-scorpions of China. *Sinensia*, **7**: 113-127.
- Yu GY. 2004. The diversity of Chinese insect species. *Contemporary entomological research* (The Entomological Society of China to establish 60 anniversary conference and Academic Symposium), 177-179.
- Zhang CB, Fang CW, Liu SJ, Ji KM. 2009. The pharmacognosy and pharmacology research of East Asia scorpion *Mesobuthus martensii*. *Journal of Anhui TCM College*, **28**(2): 57-59.
- Zhang L, Zhu MS. 2009. Morphological variation of *Mesobuthus martensii* (Karsch, 1879) (Scorpiones: Buthidae) in Northern China. *Euscorpius*, **81**: 1-17.
- Zhu MS, Han GX, Lourenço WR. 2008. The chaerilid scorpions of China (Scorpiones: Chaerilidae). *Zootaxa*, **1943**: 37-52.
- Zhu MS, Qi JX, Song DX. 2004. A Checklist of Scorpions from China (Arachnida: Scorpiones). *Acta Arachnologica Sinica*, **13**(2): 111-118.
- Zhu MS, Zhang L, Lourenço WR. 2007. One new species of scorpion belonging to the genus *Euscorpiops* Vachon, 1980 from South China (Scorpiones: Euscorpiidae, Scorpiopinae). *Zootaxa*, **1582**: 19-25.