

## Effects of temperature acclimation on body mass and energy budget in the Chinese bulbul *Pycnonotus sinensis*

Yu-Nan WU<sup>1</sup>, Lin LIN<sup>1</sup>, Yu-Chao XIAO<sup>1</sup>, Li-Meng ZHOU<sup>1</sup>, Meng-Si WU<sup>1</sup>, Hui-Ying ZHANG<sup>1</sup>, Jin-Song LIU<sup>1,2,3,\*</sup>

1. School of Life and Environmental Sciences, Wenzhou University, Wenzhou 325035, China

2. Zhejiang Provincial Key Lab for Subtropical Freshwater Environment and Marine Biological Resources Protection, Wenzhou 325035, China

3. Institute of Applied Ecology, Wenzhou University, Wenzhou 325035, China

**Abstract:** Chinese bulbuls (*Pycnonotus sinensis*) are small passerine birds that inhabit areas of central, southern and eastern China. Previous observations suggest that free-living individuals of this species may change their food intake in response to seasonal changes in ambient temperature. In the present study, we randomly assigned Chinese bulbuls to either a 30 °C or 10 °C group, and measured their body mass (BM), body temperature, gross energy intake (GEI), digestible energy intake (DEI), and the length and mass of their digestive tracts over 28 days of acclimation at these temperatures. As predicted, birds in the 30 °C group had lower body mass, GEI and DEI relative to those in the 10 °C group. The length and mass of the digestive tract was also lower in the 30 °C group and trends in these parameters were positively correlated with BM, GEI and DEI. These results suggest that Chinese bulbuls reduced their absolute energy demands at relatively high temperatures by decreasing their body mass, GEI and DEI, and digestive tract size.

**Keywords:** Body mass; Energy budget; *Pycnonotus sinensis*; Temperature acclimation

Thermogenesis and adjustment of energy intake are important for the survival of winter—particularly active birds in their natural environment (O'Connor, 1995, Yuni & Rose, 2005). The daily energy expenditure of many birds is usually higher in winter (Bryant et al, 1985; O'Connor, 1995; Stokkan et al, 1986) and this change may be triggered by environmental factors such as photoperiod, ambient temperature, and diet quality and/or quantity (Starck, 1996; 1999; Boon et al, 2000; Klaassen et al, 2004). Temperature is an important environmental *zeitgeber* for seasonal acclimatization in birds (McKechnie et al, 2007; Swanson, 2001; Tieleman et al, 2003; Vézina et al, 2006). Low-temperature stress during winter in temperate zones may induce an increase in energy expenditure during a season of generally decreased food availability (Krams et al, 2010). Many birds have a variety of strategies to cope with low temperature stress, such as increasing body mass, food intake and thermogenesis (Bednekoff et al, 1994; Goymann et al, 2006; Krams et al, 2010; Webster & Weathers, 2000; Williams & Tieleman, 2000; Swanson, 2010). The body mass of a bird is the sum of its energy

intake and energy expenditure. In a bird maintaining constant body mass, time-averaged energy intake equals time-averaged energy use (Hammond & Diamond, 1997). This balance depends on the interplay between the intake and digestive processing of matter and energy, and their allocation among diverse functions, including thermoregulation, growth and reproduction (Caviedes-Vidal et al, 2007). Accordingly, morphological changes in the digestive tract provide a useful indication of energy expenditure (Williams & Tieleman, 2000; Karasov et al, 2011).

Phenotypic flexibility is a response to environmental conditions varying predictably, or of more stochastic fluctuations in the environment (Piersma & Drent, 2003; Starck & Rahmaan, 2003). Such responses to variable conditions ought to be reversible and repeatable

Received: 28 March 2013; Accepted: 26 May 2013

Foundation items: This study was supported by the National Natural Science Foundation of China (31070366) and the 'Xinmiao' Project in Zhejiang Province.

\*Corresponding author, E-mail: ljs@wzu.edu.cn

(McWilliams & Karasov, 2001; Starck & Rahmaan, 2003). Several authors suggest that the avian digestive tract is a suitable model for the study of phenotypic plasticity because its morphology is positively correlated with food composition (frugivorous and nectar-feeding birds have smaller and shorter digestive tracts than granivorous and insectivorous species), and depends not only on resorption and assimilation processes, but also on fine adjustment of its morphology, such as epithelial resorptive surface dimensions, volume and transport efficiency (DeGoulier *et al.*, 1999; Goymann *et al.*, 2006; Karasov *et al.*, 2004; Lavin *et al.*, 2008; Starck & Rahmaan, 2003; Webster & Weathers, 2000).

Chinese bulbuls (*Pycnonotus sinensis*) are small passerines that are distributed over vast areas of central, southern and eastern China (MacKinnon & Phillipps, 2000). They have been reported to have high body temperature ( $T_b$ ), high upper critical temperature ( $T_{uc}$ ), low BMR, a relatively wide thermal neutral zone (TNZ) (Zhang *et al.*, 2006), and they increase their body mass and BMR in response to colder temperatures (Zheng *et al.*, 2008b). There is evidence that this species enhances its thermogenic capacity by increasing organ mass and respiratory enzyme activity (Zhang *et al.*, 2008; Zheng *et al.*, 2010). For this study, we selected Chinese bulbuls as a study species because they are resident in Zhejiang Province where we are based, and global warming appears to have allowed this species to colonize northeastern and northwestern China (Li *et al.*, 2006; Song, 2006), and previous studies (Ni *et al.*, 2010; Peng *et al.*, 2010; Zhang *et al.*, 2006, 2008; Zheng *et al.*, 2008a, 2010; Zhou *et al.*, 2010) provide critical background information required for our research. In this report, we investigated changes in body mass, energy budget and digestive tract morphology under two different metabolic loads, or energy demands, elicited by ambient temperature. We hypothesized that phenotypic variation in body mass, energy budget and digestive organs may play important roles in the adaptation of Chinese bulbuls to changing environmental conditions. We predicted that Chinese bulbuls will decrease their body mass and energy intake in response to warm ambient temperatures similar to those typically experienced in summer. Furthermore, when individuals have low energy requirements we predict that the mass and length of their digestive tract will decrease accordingly.

## MATERIALS AND METHODS

In Wenzhou, the climate is warm-temperate with an average annual rainfall of 1 700 mm spread across all months, with slightly more precipitation during winter and spring. The mean annual temperature is 18 °C. Mean maximum daily temperature ranges from 39 °C in July to

8 °C in January, and mean minimum daily temperature from 28 °C in July to 3 °C in the same month. Mean winter and summer temperatures are 6 °C and 31 °C, respectively (Zheng *et al.*, 2008b).

### Animals

Chinese bulbuls were captured in mist nets in Wenzhou city during March 2011. Body mass to the nearest 0.1 g was determined immediately upon capture with a Sartorius balance (model BT25S). Birds were transported to the laboratory on the day of capture and caged outdoors for 1 or 2 d in 50×30×20 cm enclosures under natural photoperiods and temperatures of 13±1 °C before temperature acclimatization and measurements commenced. Food and water were supplied *ad libitum*. Birds were moved into individual cages for at least two weeks, after which 16 birds were randomly assigned to either the 30 °C or 10 °C group (eight birds in each group). Both groups were acclimated to these temperatures for 28 days. Body temperature was measured at 21:00 and 23:00 using a digital thermometer (Beijing Normal University Instruments Co.). The probe of the thermometer was inserted 3 cm into each bird's cloaca and a reading taken within 30 s.

### Energy budget

We regarded digestible energy intake as an index of total daily energy expenditure. During the 28 day experimental period, food was provided quantitatively and water was provided *ad libitum*. Food residues and feces were collected during the two days before temperature acclimation began (week 0) and once a week (every seventh day) thereafter throughout the four week experimental period. These residues were separated manually, then oven-dried at 60 °C until a constant mass was obtained. The caloric content of the residual food and feces were determined using a C200 oxygen bomb calorimeter (IKA Instrument, Germany). Gross energy intake (GEI), feces energy (FE), digestible energy intake (DEI), and digestibility of energy were calculated according to Grodzinski & Wunder (1975) and Ni *et al.* (2010):

$$\text{GEI (kJ/day)} = \text{dry food intake (g/day)} \times \text{caloric value of dry food (kJ/g)}$$

$$\text{FE (kJ/day)} = \text{dry mass of feces (g/day)} \times \text{caloric value of dry feces (kJ/g)}$$

$$\text{DEI (kJ/day)} = \text{GEI (kJ/day)} - \text{FE (kJ/day)}$$

$$\text{Digestibility (\%)} = \text{DEI (kJ/day)} / \text{GEI (kJ/day)} \times 100\%$$

### Measurements of organ masses

Following the observation period, all birds were killed humanely and their digestive tracts (gizzard, small intestine and rectum) were removed, measured (±1 mm) and weighed (±0.1 mg) at the end of the 28 day acclimation period. The gizzard, small intestine and

rectum were then rinsed with saline to remove all gut contents before being dried and reweighed. These organs were then dried to a constant mass over 2 d at 75 °C and reweighed to the nearest 0.1 mg (Liu & Li, 2006; Williams & Tieleman, 2000).

### Statistics

Data were analyzed using SPSS 12.0. All variables were tested for normality using the Kolmogorov-Smirnov test. Non-normally distributed data were transformed to natural logarithms. Repeated-measures analysis of variance (RM-ANOVA) was used to determine the significance of changes in body mass, body temperature, GEI, FE, DEI and digestibility over time. Least significant difference (LSD) *post hoc* tests were used to detect significant differences in the above parameters within the same group of birds between different days of temperature acclimation. The independent sample *t*-test was used to assess the significance of any differences in the body mass between 10 °C and 30 °C birds measured on the same day. Differences in the above variables between 10 °C and 30 °C birds, except for those in body mass, were evaluated using ANOVA or ANCOVA with body mass as a covariate where appropriate. In the case of digestibility, which was a percentage value, the arcsine-square-root transformation was performed prior to analysis to normalize the data. All results are expressed as mean±SEM, with  $P < 0.05$  being considered statistically significant.

## RESULTS

There were no significant differences in body mass, body temperature, GEI, FE, DEI or digestibility between the 10 °C and 30 °C birds prior to the experiment.

### Body mass and body temperature

The body mass of the 30 °C birds decreased significantly throughout the course of the experiment (RM-ANOVA,  $F_{4,35}=2.718$ ,  $P < 0.05$ , Figure 1A). Although this decrease in body mass was not significant after day 7 or day 14 (*post hoc*,  $P > 0.05$ ), it was after day 21 (*post hoc*,  $P < 0.05$ ) and by day 28 the body mass of the 30 °C group had decreased by 11.2% relative to that measured on day 0 (*post hoc*,  $P < 0.05$ ). In contrast, the 10 °C birds maintained relatively constant body masses over the duration of the experiment (RM-ANOVA,  $F_{4,35}=0.420$ ,  $P > 0.05$ ).

There was no significant difference in body mass between the 30 °C and 10 °C birds after day 7 ( $t_{14}=0.550$ ,  $P > 0.05$ ), day 14 ( $t_{14}=0.196$ ,  $P > 0.05$ ) and day 21 ( $t_{14}=0.2126$ ,  $P > 0.05$ ), but there was after day 28 ( $t_{14}=2.371$ ,  $P < 0.05$ ). At the end of experiment, the body mass of the 30 °C group was 7.5% lower than that of the 10 °C group (Figure 1A).




Figure 1 Trend in body mass and body temperature in Chinese bulbuls acclimated to either 30 °C or 10 °C

Data are presented as mean±SE; Different letters (a or b) indicate significant differences ( $P < 0.05$ ) between measurements made on birds acclimated to 30 °C on different days; \*,  $P < 0.05$ .

Neither the 10 °C nor the 30 °C group underwent significant changes in body temperature during the course of the experiment (RM-ANOVA, 10 °C,  $F_{4,35}=1.861$ ,  $P > 0.05$ ; 30 °C,  $F_{4,35}=1.349$ ,  $P > 0.05$ , Figure 1B), nor were any significant between-group differences in body temperature detected (day 7,  $F_{1,14}=2.679$ ,  $P > 0.05$ ; day 14,  $F_{1,14}=0.009$ ,  $P > 0.05$ ; day 21,  $F_{1,14}=0.260$ ,  $P > 0.05$ ; day 28,  $F_{1,14}=0.531$ ,  $P > 0.05$ ).

### Energy intake and digestibility

Birds in the 10 °C group did not undergo significant changes in FE (RM-ANOVA,  $F_{4,34}=2.239$ ,  $P > 0.05$ , Figure 2B) or digestibility (RM-ANOVA,  $F_{4,34}=1.283$ ,  $P > 0.05$ , Fig. 2D), but did undergo significant changes in GEI (RM-ANOVA,  $F_{4,34}=6.494$ ,  $P < 0.01$ , Figure 2A) and DEI (RM-ANOVA,  $F_{4,34}=6.494$ ,  $P < 0.001$ , Figure 2C). Although no significant change in these two parameters occurred during the first three weeks of the experiment, both had decreased significantly by day 28 (*post hoc*,  $P < 0.05$ ).

Birds in the 30 °C group underwent significant changes in GEI, FE, DEI and digestibility (RM-ANOVA, GEI,  $F_{4,34}=24.508$ ,  $P < 0.001$ , Fig. 2A; FE,  $F_{4,34}=15.741$ ,  $P < 0.001$  Figure 2B; DEI  $F_{4,34}=22.236$ ,  $P < 0.001$  Figure 2C; digestibility,  $F_{4,34}=3.438$ ,  $P < 0.05$ ; Figure 2D). Significant changes in GEI, FE and DEI were evident by




Figure 2 Trends in GEI, FE, DEI and digestibility in Chinese bulbuls acclimated to either 30 °C or 10 °C

Data are presented as mean±SEM; Different letters (a, b, c, or d for the 30 °C group; and A, B or C for the 10 °C group) indicate significant differences ( $P<0.05$ ) in measurements made on birds in either the 30 °C or 10 °C group on different days; \*\*:  $P<0.01$ ; \*\*\*:  $P<0.001$  indicate significant differences between the group of birds acclimated to 30°C and those acclimated to 10 °C.

day 7 (*post hoc*, GEI,  $P<0.05$ ; FE,  $P<0.05$ ; DEI,  $P<0.05$ ) and continued over the subsequent three weeks. The GEI, FE and DEI of the 30 °C group were significantly lower than those of the 10 °C group on days 7, 14, 21 and 28 (ANCOVA, day 7, GEI,  $F_{1,13}=24.271$ ,  $P<0.001$ ; FE,  $F_{1,13}=14.812$ ,  $P<0.01$ ; DEI  $F_{1,13}=14.395$ ,  $P<0.01$ ; day 28, GEI,  $F_{1,13}=46.130$ ,  $P<0.001$ ; FE,  $F_{1,13}=18.252$ ,  $P<0.01$ ; DEI  $F_{1,13}=96.832$ ,  $P<0.001$ ). No significant difference in digestibility between the 10 °C and 30 °C groups was observed (ANCOVA, day 28,  $F_{1,13}=0.305$ ,  $P>0.05$ ). Log GEI and DEI were positively correlated with log body mass at day 28 (GEI:  $r^2=0.372$ ,  $P<0.01$ ; DEI:  $r^2=0.458$ ,  $P<0.01$ , Figure 3).

### Digestive tract morphology

Although no significant changes in gizzard and rectum length (ANCOVA, gizzard,  $F_{1,13}=0.027$ ,  $P>0.05$ ; rectum,  $F_{1,13}=1.108$ ,  $P>0.05$ , Figure 4) were detected, the length of the total digestive tract and small intestine varied significantly over the course of the experiment (ANCOVA, total digestive tract,  $F_{1,13}=5.325$ ,  $P<0.05$ ;




Figure 3 Least squares regression of GEI and DEI as dependent variable of body mass in Chinese bulbuls acclimated to 30 °C or 10 °C

small intestine,  $F_{1,13}=5.626$ ,  $P<0.05$ , Figure 4), becoming shorter in 30 °C birds than in 10 °C birds. The log of total digestive tract length was positively correlated with log body mass, GEI and DEI at day 28 (body mass:  $r^2=0.267$ ,  $P<0.05$ , Figure 6; GEI:  $r^2=0.237$ ,  $P<0.05$ , Figure 7; DEI:  $r^2=0.399$ ,  $P<0.01$ , Figure 8).

There was no significant difference in gizzard mass between the 10 °C and 30 °C birds at the end of the experiment (ANCOVA, wet mass with content,  $F_{1,13}=0.706$ ,  $P>0.05$ ; wet mass without contents,  $F_{1,3}=0.396$ ,  $P>0.05$ , dry mass,  $F_{1,13}=0.434$ ,  $P>0.05$ , Figure 5).

However, the mass of the small intestine changed significantly (ANCOVA, wet mass with contents,  $F_{1,13}=7.158$ ,  $P<0.05$ ; wet mass without contents,  $F_{1,13}=13.491$ ,  $P<0.01$ ; dry mass,  $F_{1,13}=15.569$ ,  $P<0.01$ , Figure 5). Small intestine wet mass with contents, wet mass without contents and dry mass were 30.1%, 35.0% and 36.3%

lower, respectively, in the 30 °C group than in their 10 °C counterparts.




Figure 4 Effect of temperature acclimation on digestive tract length in the Chinese bulbul. Data are presented as mean $\pm$ SEM; \*:  $P<0.05$




Figure 5 Effect of temperature acclimation on digestive tract mass in the Chinese bulbul

Data are presented as mean $\pm$ SEM. \*:  $P<0.05$ ; \*\*:  $P<0.01$ .

No significant difference in rectal mass between 10 °C and 30 °C birds was evident at day 28 (ANCOVA, wet mass with contents,  $F_{1,13}=1.832$ ,  $P>0.05$ ; wet mass without contents,  $F_{1,3}=1.311$ ,  $P>0.05$ , dry mass,  $F_{1,13}=0.484$ ,  $P>0.05$ , Figure 5).

However, bulbuls in the 30 °C group had a lighter digestive tract overall than 10 °C birds (ANCOVA, wet mass with contents,  $F_{1,13}=7.580$ ,  $P<0.05$ ; wet mass without contents,  $F_{1,13}=15.858$ ,  $P<0.01$ ; dry mass,  $F_{1,13}=16.275$ ,  $P<0.01$ , Figure 5). The log of wet and dry mass of the complete digestive tract was positively

correlated with log body mass, GEI and DEI at day 28 (body mass, wet mass:  $r^2=0.462$ ,  $P<0.01$ ; dry mass:  $r^2=0.626$ ,  $P<0.001$ , Figure 6; GEI, wet mass:  $r^2=0.635$ ,  $P<0.001$ ; dry mass:  $r^2=0.726$ ,  $P<0.001$ , Figure 7; DEI, wet mass:  $r^2=0.615$ ,  $P<0.001$ ; dry mass:  $r^2=0.712$ ,  $P<0.001$ , Figure 8).

## DISCUSSION

Our results suggest that ambient temperature had significant effects on the body mass, GEI, FE, and DEI of

Chinese bulbuls, all of which decreased significantly in birds acclimatized to 30 °C. These birds also underwent a significant decrease in the length and mass of the digestive tract.

Ambient temperature is known to have significant effects on many morphological physiological and biochemical parameters of birds, including body mass (Tieleman et al, 2003; Cooper, 2007; McKechnie et al, 2007; Zheng et al, 2008a). Our body mass results were consistent with those of previous reports on the response of Chinese bulbuls to seasonal changes (Zhang et al, 2008; Zheng et al, 2008b, 2010). We found that bulbuls acclimatized to 10 °C maintained a relatively constant body mass but that those acclimatized to 30 °C decreased in body mass over the four week experimental period, and by the end of the experiment the body mass of these birds was 7.5% lower than that of those kept at 10 °C (Figure 1A). The decrease in the body mass of bulbuls acclimatized to 30 °C reflected an imbalance between energy intake and expenditure, a hypothesis supported by changes in other parameters, such as GEI, DEI, and the length and mass of the digestive tract.

Environmental temperature can alter the energy intake of birds. Cold winter temperatures will increase energy consumption because of the increased need for energy to maintain body temperature. Warmer temperatures on the other hand decrease energy requirements because less heat needs to be produced to maintain body temperature (Cain, 1973; Syafwan et al,

2012). There is marked variation across avian species in the reported energy budgets associated with acclimatization or acclimation to changing ambient temperatures (Salvante et al, 2010; Syafwan et al, 2012). For example, the GEI of warm (38 °C) acclimated Muscovy ducks *Cairina moschata* decreased by 49% relative to that of control (20 °C) birds, and that of cold (-10 °C) acclimated ducks increased by 62% relative to that of control birds in an 11 hour photoperiod (Cain, 1973). Similarly, the GEI of summer acclimatized Elliot's pheasants *Syrnaticus ellioti* was 29% and 34% lower than that of autumn and winter acclimatized birds, respectively (Lou et al, 2003). This suggests that changes in GEI and DEI are a common response to variation in ambient temperature for small birds and mammals in both warm and temperate habitats (Stokkan et al, 1986). We found that Chinese bulbuls acclimatized to 30 °C underwent a 47% decrease in GEI, and a 45% decrease in DEI, relative to those kept at 10 °C. These changes in GEI and DEI were positively correlated with body mass (Figure 3). These results suggest that bulbuls acclimatized to 30 °C experienced an obvious decrease in energy requirements. One could predict that, if different physiological systems have to compete for energy, the combination of these energetically challenging processes would affect heat production, as we found. Chinese bulbuls acclimatized to 30 °C for four weeks decreased their basal metabolic rate (BMR), liver, kidney and small intestine mass, and mitochondrial state-4




Figure 6 Least square regression of length and wet and dry mass of the digestive tract as dependent variables of body mass in Chinese bulbuls acclimatized to 30 °C or 10 °C




Figure 7 Least square regression of length and wet and dry mass of the digestive tract as dependent variables of GEI in Chinese bulbuls acclimatized to 30 °C or 10 °C

respiration and cytochrome c oxidase (COX) activity in liver and muscle, compared to those kept at 10 °C (Zheng et al., 2013). In view of the mass-specific energy metabolism of these organs and/or tissues, the observed decreases in body mass, GEI and DEI were not surprising.

Environmental temperature also influences morphological and physiological functions of the digestive tract in birds (Williams & Tieleman, 2000; Starck & Rahmaan, 2003; Karasov, 2011; Karasov et al., 2011). During cold conditions, when energy demands increase, small birds may increase their energy intake, which simultaneously compromises their digestive efficiency unless they also undergo associated changes in gut size, enzyme activity, nutrient uptake and/or food transit time (McKinney & McWilliams, 2005; Karasov et al., 2011). For example, warm (35 °C) acclimated woodlarks, skylarks, spike-heeled larks *Chersomanes albobasciata* and Dunn's lark underwent a respective 21%, 22%, 9%, and 24% decrease in stomach mass, and a 21%, 23%, 23%, and 21% decrease in intestinal mass, relative to that of cold (15 °C) acclimated conspecifics (Tieleman et al., 2003). Similarly, summer acclimatized spruce grouse *Canachites canadensis* had a 44% lower ventricular mass and a 21% shorter small intestine than winter acclimatized birds (Pendergast & Boag, 1973). According to Sibly's (1981) model of optimal digestion, one of the advantages of increasing digestive tract size is that it allows an increase in the mean retention time of digesta, thereby increasing digestibility if the ingestion rate is constant. Alternately, it allows a constant mean retention time of digesta, which maintains digestive efficiency if the ingestion rate increases. Our results showed that bulbuls acclimatized to 30 °C decreased the size of their gastrointestinal tracts relative to those bulbuls kept at 10 °C. The significant, positive relationship between log body mass, GEI and DEI (Figure 6, Figure 7 and Figure 8), suggests that this was an adaptive response to decreased body mass and energy intake.

In conclusion, Chinese bulbuls in the 30 °C group had lower body mass, GEI and DEI relative to those in




Figure 8 Least square regression of length wet and dry mass of the digestive tract as dependent variables of DEI in Chinese bulbuls acclimated to 30 °C or 10 °C

the 10 °C group. The length and wet and dry mass of the digestive tract of bulbuls were also lower in the 30 °C group and had significant positive relationships with body masses, GEI and DEI. Chinese bulbuls reduced their energy demands at high temperatures by decreasing their GEI and DEI, and digestive tract size.

**Acknowledgements:** We are grateful to Dr David L. Swanson for providing several references. We thank Dr Ron Moorhouse for revising the English and giving some suggestions.

## References

- Bednekoff PA, Biebach H, Krebs J. 1994. Great tit fat reserves under unpredictable temperatures. *Journal of Avian Biology*, **25**(2): 156-160.
- Boon P, Visser GH, Daan S. 2000. Effect of photoperiod on body weight gain, and daily energy intake and energy expenditure in Japanese quail (*Coturnix c. japonica*). *Physiology and Behavior*, **70**(3-4): 249-260.
- Bryant DM, Hails CJ, Prys-Jones R. 1985. Energy expenditure by free-living dippers (*Cinclus cinclus*) in winter. *The Condor*, **87**(2): 177-186.
- Cain BW. 1973. Effect of temperature on energy requirements and northward distribution of the black-bellied tree duck. *The Wilson Bulletin*, **85**(3): 308-317.

- Caviedes-Vidal E, McWhorter TJ, Lavin SR, Chediack JG, Tracy CR, Karasov WH. 2007. The digestive adaptation of flying vertebrates: High intestinal paracellular absorption compensates for smaller guts. *Proceedings of the National Academy of Sciences of the United States of America*, **104**(48): 19132-19137.

- Cooper SJ. 2007. Daily and seasonal variation in body mass and visible fat in mountain chickadees and juniper titmice. *The Wilson Journal of Ornithology*, **119**(4): 720-724.

- DeGolier TF, Mahoney SA, Duke GE. 1999. Relationship of avian cecal lengths to food habits, taxonomic position, and intestinal lengths. *The*

- Condor*, **101**(3): 622-634.
- Goymann W, Trappschuh M, Jensen W, Schwabl I. 2006. Low ambient temperature increases food intake and dropping production, leading to incorrect estimates of hormone metabolite concentrations in European stonechats. *Hormones and Behavior*, **49**(5): 644-653.
- Grodzinski W, Wunder BA. 1975. Ecological energetics of small mammals. In: Golley FB, Petruszewicz K, Ryszkowski L, eds. *Small Mammals: Their Productivity and Population Dynamics*. Cambridge: Cambridge University Press, 173-204.
- Hammond KA, Diamond J. 1997. Maximal sustained energy budgets in humans and animals. *Nature*, **386**(6624): 457-462.
- Karasov WH. 2011. Digestive physiology: a view from molecules to ecosystem. *American Journal of Physiology*, **301**(2): R276-R284.
- Karasov WH, Pinshow B, Starck JM, Afik D. 2004. Anatomical and histological changes in the alimentary tract of migrating Blackcaps (*Sylvia atricapilla*): A comparison among fed, fasted, food-restricted, and refed birds. *Physiological and Biochemical Zoology*, **77**(1): 149-160.
- Karasov WH, Martínez del Río C, Caviedes-Vidal E. 2011. Ecological physiology of diet and digestive systems. *Annual Review of Physiology*, **73**(1): 69-93.
- Klaassen M, Oltrogge M, Trost L. 2004. Basal metabolic rate, food intake, and body mass in cold-and warm-acclimated Garden Warblers. *Comparative Biochemistry and Physiology Part A*[0], **137**(4): 639-647.
- Krams I, Cirule D, Suraka V, Krama T, Rantala MJ, Ramey G. 2010. Fattening strategies of wintering great tits support the optimal body mass hypothesis under conditions of extremely low ambient temperature. *Functional Ecology*, **24**(1): 172-177.
- Lavin SR, Karasov WH, Ives AR, Middleton KM, Garland TJr. 2008. Morphometrics of the avian small intestine compared with that of nonflying mammals: a phylogenetic approach. *Physiological and Biochemical Zoology*, **81**(5): 526-550.
- Li DM, Wang AZ, Lei FM. 2006. A new bird record in Qinghai Province-*Pycnonotus sinensis*. *Chinese Journal of Zoology*, **41**(4): 70. (in Chinese)
- Liu JS, Li M. 2006. Phenotypic flexibility of metabolic rate and organ masses among tree sparrows *Passer montanus* in seasonal acclimatization. *Acta Zoologica Sinica*, **52**(3): 469-477.
- Lou Y, Yu TL, Huang CM, Zhao T, Li HH, Li CJ. 2013. Seasonal variations in the energy budget of Elliot's pheasant (*Syrnaticus ellioti*) in cage. *Zoological Research*, **34**(E1): E19-E25.
- MacKinnon J, Phillipps K. 2000. *A Field Guide to the Birds of China*. London: Oxford University Press, 491-493.
- McWilliams SR, Karasov WH. 2001. Phenotypic flexibility in digestive system structure and function in migratory birds and its ecological significance. *Comparative Biochemistry and Physiology Part A*[0], **128**(3): 577-591.
- McKinney RA, McWilliams SR. 2005. A new model to estimate daily energy expenditure for wintering waterfowl. *The Wilson Bulletin*, **117**(1): 44-55.
- McKechnie AE, Chetty K, Lovegrove BG. 2007. Phenotypic flexibility in basal metabolic rate in laughing doves: responses to short-term thermal acclimation. *Journal of Experimental Biology*, **210**(1): 97-106.
- Ni XY, Lin L, Zhou FF, Wang XH, Liu JS. 2010. Effects of photoperiod on body mass, organ masses and energy metabolism in Chinese bulbul (*Pycnonotus sinensis*). *Acta Ecologica Sinica*, **31**(6): 1703-1713. (in Chinese)
- O'Connor TP. 1995. Metabolic characteristics and body composition in House Finches: effects of seasonal acclimatization. *Journal of Comparative Physiology B*, **165**(4): 298-305.
- Pendegast BA, Boag DA. 1973. Seasonal changes in the internal anatomy of spruce grouse in Alberta. *The Auk*, **90**(2): 307-317.
- Peng LJ, Tang XL, Liu JS, Meng HT. 2010. The effects of thyroid hormone on basal thermogenesis (*Pycnonotus sinensis*). *Acta Ecologica Sinica*, **30**(6): 1500-1507. (in Chinese)
- Piersma T, Drent J. 2003. Phenotypic flexibility and the evolution of organismal design. *Trends Ecology and Evolution*, **18**(5): 228-233.
- Salvante KG, Vézina F, Williams TD. 2010. Evidence for within-individual energy reallocation in cold-challenged, egg-producing birds. *Journal of Experimental Biology*, **213**(12): 1991-2000.
- Sibly RM. 1981. Strategies in digestion and defecation. In: Townsend CR, Calow P. *Physiological Ecology: An Evolutionary Approach to Resource Use*. Oxford: Blackwell, 109-139.
- Song ZM. 2006. A new bird record in Northeast: *Pycnonotus sinensis*. *Journal of Mudanjiang Normal University: Natural Science*, **54**: 1-2. (in Chinese)
- Starck JM. 1996. Phenotypic plasticity, cellular dynamics, and epithelial turnover of the intestine of Japanese quail (*Coturnix coturnix japonica*). *Journal of Zoology*, **238**(1): 53-79.
- Starck JM. 1999. Phenotypic flexibility of the avian gizzard: rapid, reversible and repeated changes of organ size in response to changes in dietary fibre content. *Journal of Experimental Biology*, **202**(22): 3171-3179.
- Starck JM, Rahmaan GHA. 2003. Phenotypic flexibility of structure and function of the digestive system of Japanese quail. *Journal of Experimental Biology*, **206**(11): 1887-1897.
- Stokkan KA, Mortensen A, Blix AS. 1986. Food intake, feeding rhythm, and body mass regulation in Svalbard rock ptarmigan. *American Journal of Physiology*, **251**(2): R264-R267.
- Swanson DL. 2001. Are summit metabolism and thermogenic endurance correlated in winter-acclimated passerine birds? *Journal of Comparative Physiology B*, **171**(6): 475-481.
- Swanson DL. 2010. Seasonal metabolic variation in birds: functional and mechanistic correlates. In: Thompson CF. *Current Ornithology*. Berlin: Springer, 17: 75-129.
- Syafwan S, Wermink GJD, Kwakkel RP, Verstegen MWA. 2012. Dietary self-selection by broilers at normal and high temperature changes feed intake behavior, nutrient intake, and performance. *Poultry Science*, **91**(3): 537-549.
- Tieleman BI, Williams JB, Buschur ME, Brown CR. 2003. Phenotypic variation of larks along an aridity gradient: are desert birds more flexible? *Ecology*, **84**(7): 1800-1815.
- Vézina F, Jalvingh K, Dekinga A, Piersma T. 2006. Acclimation to different thermal conditions in a northerly wintering shorebird is driven by body mass-related changes in organ size. *Journal of Experimental Biology*, **209**(16): 3141-3154.



- Webster MD, Weathers WW. 2000. Seasonal changes in energy and water use by verdins, *Auriparus flaviceps*. *Journal of Experimental Biology*, **203**(21): 3333-3344.
- Williams J, Tieleman BI. 2000. Flexibility in basal metabolic rate and evaporative water loss among hoopoe larks exposed to different environmental temperatures. *Journal of Experimental Biology*, **203**(20): 3153-3159.
- Yuni LPEK, Rose RW. 2005. Metabolism of winter-acclimatized New Holland honeyeaters *Phylidonyris novaehollandiae* from Hobart, Tasmania. *Acta Zoologica Sinica*, **51**(2): 338-343.
- Zhang GK, Fang YY, Jiang XH, Liu JS, Zhang YP. 2008. Adaptive plasticity in metabolic rate and organ masses among *Pycnonotus sinensis*, in seasonal acclimatization. *Chinese Journal of Zoology*, **43**(4): 13-19. (in Chinese)
- Zhang YP, Liu JS, Hu XJ, Yang Y, Chen LD. 2006. Metabolism and thermoregulation in two species of passerines from south-eastern China in summer. *Acta Zoologica Sinica*, **52**(4): 641-647.
- Zheng WH, Li M, Liu JS, Shao SL. 2008a. Seasonal acclimatization of metabolism in Eurasian tree sparrows (*Passer montanus*). *Comparative Biochemistry and Physiology Part A*[0], **151**(4): 519-525.
- Zheng WH, Liu JS, Jang XH, Fang YY, Zhang GK. 2008b. Seasonal variation on metabolism and thermoregulation in Chinese bulbul. *Journal of Thermal Biology*, **33**(6): 315-319.
- Zheng WH, Fang YY, Jang XH, Zhang GK, Liu JS. 2010. Comparison of thermogenic character of liver and muscle in Chinese bulbul *Pycnonotus sinensis* between summer and winter. *Zoological Research*, **31**(3): 319-327. (in Chinese)
- Zheng WH, Lin L, Liu JS, Pan H, Cao MT, Hu YL. 2013. Physiological and biochemical thermoregulatory responses of Chinese bulbuls *Pycnonotus sinensis* to warm temperature: Phenotypic flexibility in a small passerine. *Journal of Thermal Biology*, **38**(5): 483-490.
- Zhou W, Wang YP, Chen DH, Liu JS. 2010. Diurnal rhythms of Chinese bulbul (*Pycnonotus sinensis*) body temperature, body mass, and energy metabolism. *Chinese Journal of Ecology*, **29**(12): 2395-2400. (in Chinese)