

The Effects of Progesterone on Oocyte Maturation and Embryo Development

Mojdeh Salehnia, Ph.D.^{1*}, Saeed Zavareh, Ph.D.²

1. Department of Anatomy, Tarbiat Modares University, Tehran, Iran

2. Department of Cellular and Molecular Biology, School of Biology, Damghan University, Damghan, Iran

Abstract

Oocyte maturation and embryo development are controlled by intra-ovarian factors such as steroid hormones. Progesterone (P4) exists in the follicular fluid that contributes to normal mammalian ovarian function and has several critical functions during embryo development and implantation, including endometrial receptivity, embryonic survival during gestation and transformation of the endometrial stromal cells to decidual cells.

It is well known that the physiological effects of P4 during the pre-implantation stages of some mammal's embryos are mediated by P4 receptors and their gene expression is determined. The effects of P4 on oocytes and embryo development have been assessed by some investigations, with contradictory results. P4, a dominant steroid in follicular fluid at approximately 18 hours after the luteinizing hormone (LH) surge may have a critical role in maturation of oocytes at the germinal stage. However, it has been shown that different concentrations of P4 could not improve *in vitro* maturation rates of germinal vesicles (GV) in cumulus oocyte complexes (COCs) and cumulus denuded oocytes (CDOs). Culture media supplemented with P4 significantly improved mouse embryo development. In addition, an *in vivo* experimental design has shown high blastocyst survival and implantation rates in P4-treated mice.

In this review we explain some of the findings that pertain to the effects of P4 on oocyte maturation and embryo development both *in vitro* and *in vivo*.

Keywords: Progesterone, Oocyte, Embryo, *In Vitro* Maturation

Citation: Salehnia M, Zavareh S. The effects of progesterone on oocyte maturation and embryo development. *Int J Fertil Steril.* 2013; 7(2): 74-81.

Introduction

Oocyte maturation and embryo development are controlled by steroid hormones as well as intra-ovarian factors such as cytokines and growth factors (1-4). *In vivo*, oocyte maturation takes place in the presence of follicular fluid which is composed of plasma exudates and secretions of follicular cells. With each follicular developmental stage, the steroid contents of follicular fluids change and the ratio of progesterone (P4) to estradiol (E2) is related to the maturation stage of the oocytes (5-7).

During folliculogenesis the oocyte gains its developmental competence in a gradual and se-

quential manner, after which it becomes a fully mature oocyte with the capability to become fertilized and develop into a high quality embryo (8).

The process of mammalian oocyte meiosis takes place in several steps. Initiation of the first meiotic division leads to primary oocytes that occur in the fetal development period or around the time of birth. Oocytes progress through zygotene, pachytene and early diplotene stages but arrest at the dictyate stage of prophase I. At puberty the first meiotic division is completed by a surge of luteinizing hormone during the menstrual cycle; the second meiotic arrest of the oocytes occurs at ovulation. Re-



sumption of the second meiotic division occurs after penetration of the sperm (9).

P4 structure and production

P4 is a cholesterol-derived, phylogenetically old steroid hormone (10). It is synthesized during the steroid hormone metabolizing pathways from androgens, estrogens, and glucocorticoids within several cell types such as the corpus luteum, placenta and adrenal gland. In addition it is produced from a plant steroid precursor, diosgenin (11, 12).

Within the ovary, cholesterol is converted by an enzyme to pregnenolone, another precursor steroid, after which it can follow one of two pathways ($\Delta 4$ or $\Delta 5$). In the $\Delta 4$ path way pregnenolone is converted to P4. P4 not only serves as a precursor for other steroids, but enters the female's blood and acts as a hormone on target tissue (13).

The level of plasma P4 varies with sex and reproductive age. P4 is mainly bound to albumin, however it has an affinity to bind to corticosteroid-binding globulin. In the normal menstrual cycle, its levels rise during the follicular phase and reach a maximum level after ovulation. Its half-life in serum is about 5 minutes (10-12, 14).

P4 function

P4 is an intra follicular steroid that plays critical roles in ovulation, implantation and maintenance of pregnancy (15, 16). P4 is the dominant content of follicular fluid steroids in mammalian preovulatory follicles, which are temporary and elevated at 18 hours after the luteinizing hormone (LH) surge (17).

P4 was initially studied as a contraceptive agent by inhibition of the luteinizing hormone surge and ovulation (18). However, it has a critical function in pregnancy maintenance and in the regulation of different biological functions in the ovarian tissue and feto-maternal unit such as resumption of meiosis, fertilization, embryonic development and implantation (19-21).

Clinically, it can be used in the female reproductive system as luteal support during *in vitro* fertilization (IVF) (22), hormone replacement therapy for older women (23), and as treatment for endometriosis and polycystic ovarian syndrome in younger women (24). In addition, P4 has immu-

nological functions for the maintenance of a feto-maternal allograft (19).

Some investigations showed that P4 administration for luteal support improved uterine receptivity at the ultra structure levels (25-27) and enhanced the implantation rate in mice (28).

In our experiments we used ovariectomized animal models and injected exogenous hormones to evaluate the effects of P4 on endometrial morphology and gene expression. Our observations showed that exogenous P4 administration affected expression of endometrial integrin molecules (29, 30).

P4 receptors in oocytes and embryos

The biological actions of P4 are mediated by three genomic isoforms of P4 receptors (PR), PR-A, PR-B and PR-C, in addition to three non-genomic isoforms, alpha, beta, and gamma (31, 32). Although PR-A and PR-B arise from a single gene, PR-A is a more important repressor than PR-B (33). The PR-C isoform is the shortest isoform. PR-C does not have transcriptional activity, however it has a role in decidual cells during late pregnancy (31, 34).

Both PR-A and PR-B are expressed in preovulatory folliclegranulosa cells (35-37). The membrane PR or non-genomic PGR are particularly notable as promoters of oocyte meiosis. They are expressed in neural, kidney, and intestinal tissues in addition to the reproductive tract (32).

Mice that lack PR-A and PR-B isoforms are infertile as a result of ovulation failure (38). It has been suggested that induction of PR isoforms in cumulus cells and their binding to P4 appear to affect follicular growth, oocyte maturation, and embryo development (39, 40).

Although PR was identified in granulosa cells, there was no evidence of this receptor on the oocyte (41, 42) with the exception of one report which observed the PR receptor in *Xenopus laevis* oocytes (43). Canine oocytes have been shown to express estrogen receptors during the estrous cycle, however, there is a lack of PR expression in all phases (41).

PR membrane component 1 (PGRMC1) is another potential mediator of P4 action (44). Possibly PGRMC1 mediates anti-apoptotic actions of P4 (45). Western blot analysis has demonstrated the

presence of PGRMC1 in bovine germinal vesicle (GV) and metaphase II (MII) oocytes (46) as well as rat (44), and human oocytes (47). PGRMC1 expression is not only associated with male and female pronuclear formation, it is also highly expressed in blastocysts (46).

Despite high level expression of PGRMC1, little is known about its role on oocyte function, however it may be directly involved in the regulation of meiotic maturation (48).

Despite the lack of PGR (PR) expression in the oocyte, both P4 and estrogen receptor mRNA and proteins have been detected in mature cumulus oocyte complexes (COCs) and embryonic cells from several mammalian species (49-55).

Aparicio and colleagues (39) have shown decreased bovine embryo development as a result of blocking genomic PR and non-genomic PR alpha activity. This result indicates that P4 intracellular signaling is mediated by its interaction with nuclear and membrane PRs and is also important for oocyte developmental competence.

Limited work has been performed on embryo PR expression however there were varied, contradictory results. Expression of PR during the pre-implantation stages of pig and mouse embryos has been shown (48, 49, 51-54, 56, 57). P4 receptor mRNA was present during all stages of bovine embryo development (51). PR mRNA and protein were expressed in pre-implantation pig embryos prior to the fifth cell division but not at later stages through blastocyst formation. P4 receptor mRNA was undetectable until the blastocyst stage (54).

We have located no studies on PR expression during early organogenesis. However, mRNA and protein are expressed in increasing amounts in the female reproductive tract of the rat after organogenesis (58). In the female reproductive system, PR is expressed in the uterus, mammary gland, ovarian tissue, fallopian tubes (57) and placenta (59).

P4 and oocyte maturation

Resumption of meiosis in oocytes is triggered by steroid hormones, specifically P4, in certain species (60). The resumption of meiosis and its progression to MII in several mammalian species such as cows, sheep and pigs is steroid dependent and the inhibition of steroidogenesis in ovine follicles

leads to impairment of resumption of meiosis and progression to MII. According to research, levels of P4 in follicular fluid and its ratio to the estrogen levels are strongly associated with oocyte quality and maturity (61). However, controversy exists regarding the effect of P4 on *in vitro* oocyte maturation (IVM). Our investigations have shown that addition of P4 (10, 38, 50, 100 μ M) to the *in vitro* maturation media of mouse GV oocytes could not improve maturation rates and developmental competence of GV in COCs and cumulus denuded oocytes (CDOs) at any of the tested concentrations when compared to the control groups. When we increased P4 from 10 to 100 μ M in the culture medium, the maturation rate decreased in a dose dependent manner and the GV arrested rates increased. Research has shown that the effect of P4 in inhibition of meiotic resumption was more effective in COC than CDO (62). It seems there were intensive interactions between oocytes and the surrounding cumulus cells. Oocytes could affect cumulus cell functions.

Vanderhyden et al. (63) have shown that mouse oocytes modulate steroid production by the surrounding cumulus cells. These observations have suggested that oocytes secrete a factor (s) which control cumulus cell production of E2 and P4. In contrast, Jamnongjit et al. (64) observed that testosterone or P4 and epidermal growth factor induced meiotic resumption in mouse oocytes during their *in vitro* maturation; the effect of these steroids could have been inhibited by specific receptor antagonists. Fukui et al. (65) demonstrated that P4 supplementation of IVM culture systems decreased the rate of bovine oocyte maturation and that addition of P4 to fertilization culture medium did not improve the number of cleavage stage embryos. Carter et al. (66) have shown that addition of P4 to culture medium did not affect the proportion of *in vitro* matured/*in vitro* fertilized zygotes that developed to the blastocyst stage *in vitro*. There was no effect on conceptus elongation following transfer to synchronized recipient heifers.

Elsewhere, the role of P4 on bovine oocyte developmental competence has been investigated by inhibiting P4 production of cumulus cells. Research has shown that supplementation of oocyte maturation medium with trilostane, an inhibitor of 3 β -hydroxy steroid dehydrogenase, caused a significant decrease in the blastocyst formation rate, which was completely reversed by the addition of P4 or a P4 agonist. This observation might support a positive role for P4 in oocyte quality (39).

Supplementation of canine oocyte culture media with steroid hormones stimulates their nuclear maturation (67, 68). However, this effect of steroid hormones has not been shown in anestrus bitches (69). In rhesus monkeys, the improvement of *in vitro* oocyte development was demonstrated in the presence of P4 and E2 (68).

Overall, these inconsistent results may be due to different experimental strategies that have been used. It seems that the length of time between LH or human chorionic gonadotropin (hCG) stimulation and GV breakdown (GVBD) might explain differences among mammalian species. Possibly, maintaining the healthiness of the oocyte and its ability to mature in species with a long dormant period between LH surge and GVBD requires steroid support.

It has been shown that the P4 antagonist mifepristone (RU486) which occupies PR could not reverse the inhibitory effect of P4 on oocyte maturation (62). Therefore, it has been concluded that the inhibition of mouse oocyte maturation by P4 is not receptor dependent. It appears that P4 could inhibit cAMP phosphodiesterase (PDE) activity through binding to the purine-binding site of this enzyme, which in turn inhibits meiosis by increasing oocyte cAMP levels (70, 71).

P4 and oocyte fertilization

The role of steroids has been shown to be involved in the acquirement of meiotic competence and the ability to undergo normal fertilization and development to the blastocyst stage (33). In humans and rhesus monkeys, high ratios of P4 to E2 in follicular fluid were associated with better embryo development (72).

In this regard we attempted to investigate the effect of P4 in concentrations similar to that of preovulatory follicular fluid (10 and 38 μ M) on developmental competence of mouse GV oocytes and subsequent fertilization potential. Our experiments showed that P4 could not increase the fertilization rate and development of the embryo to the blastocyst stage (62). The result of this experiment was inconsistent with other studies (73-75). Silva and Knight (76) have shown that the addition of P4 to bovine oocyte *in vitro* maturation medium reduced the rate of blastocyst formation.

Mattioli et al. (77) reported that presence of P4 in porcine oocyte maturation medium increased subsequent sperm head decondensation and male pronuclei formation. Zhang and Armstrong (78) reported that the addition of P4 to porcine oocyte maturation medium could increase both fertilization and cleavage rates, whereas E2 could not. P4 had the opposite effect in ovine oocytes (74).

In vivo embryo development and P4

The embryo develops in tubal and uterine microenvironments that are mainly controlled by P4. P4 may act directly as a survival factor or indirectly promote the production and secretion of cytokines which contribute to embryonic survival and development (79). It has been shown that P4 elevation occurs when the embryo does not reach the uterus, thus this finding proposes that the effect of P4 on embryo development is mediated via P4-induced changes in the endometrial transcriptome (80).

Granulocyte macrophage colony stimulating factor is a cytokine secreted by the embryo and endometrium under control of P4 (81) which promotes embryo development (82). Lessey et al. (83) have shown an increase in growth factor production in the stromal cells in response to P4 administration.

Low P4 levels have been linked to early pregnancy failure (84) and poor embryo development (85), while in cattle administration of P4 enhances conceptus development (66, 86).

Some studies have shown that endogenous P4 is a main factor in the preparation of the endometrium for embryo implantation (87, 88). Due to the effects of P4 on improving pregnancy rates of IVF patients in the ART clinic, thus exogenous P4 has been used as luteal support to enhance implantation rates (89). Although P4 is essential for continuation of pregnancy in all mammals, expressions of P4 receptors cease prior to implantation. It seems the loss of P4 receptors is important for maternal recognition and embryo development in early pregnancy (90).

Several studies have described the effect of exogenous P4 supplementation on embryo development with varying results, according to the time

and duration of P4 treatment (91-95).

Initiation of P4 supplementation at the time of onset of the postovulatory rise (between days 4 and 5) resulted in consistent increases in pregnancy rate, however when P4 was administered later there was no improvement in pregnancy rate (92). The results of another study have suggested that the time of/ or strength of the postovulatory P4 rise is critical for embryo development rather than the final concentration of P4 in the luteal phase (91). Use of supraphysiological levels of P4 during early pregnancy in the mouse has resulted in a similar conclusion (96).

In our previous study we compared embryo quality and implantation rate in pregnant mice in superovulated, P4 treated and superovulated-P4 treated groups. Our observation showed a high survival rate of blastocysts (97.68%) and implantation rate (92.06%, $p < 0.001$) in pregnant mice from the P4-treated group compared to the P4 superovulated group, which meant that injections of 1 mg/mouse of progesterone in un-stimulated mice significantly improved implantation rates compared to the control and super ovulated groups (28).

***In vitro* embryo development and P4**

To answer the question of whether P4 directly affects embryo development or there is an indirect effect via changes in the endometrium, some researchers have added P4 to embryo culture medium *in vitro* and examined development to the blastocyst stage, with contradictory results (2, 97-100). These contradictory observations might be attributed to different culture systems which have been used.

In vitro and *in vivo* experiments by Clemente et al. (57) showed that the effects of P4 on conceptus elongation could be due to a direct effect of P4 on the embryo. They demonstrated that a P4 receptor was expressed in all stages of embryo development. These researchers showed the direct effect of P4 on embryo development. Supplementation of simple or co-culture embryo culture systems with P4 did not affect on the embryo development and blastocyst cell number. However, *in vitro*-derived embryo transfer to a recipient treated with P4 resulted in a four-fold increase in conceptus length on day 14. These data confirmed the hypothesis that

conceptus elongation in cattle was related to P4-induced changes in the uterine environment (57). This finding agreed with a study by Geisert et al. (101) who showed that administration of P4 early in the estrous cycle advanced uterine receptivity for the transfer of older asynchronous embryos. Supplementation of embryo culture medium with lipid-soluble P4 resulted in an increase in the numbers of embryos that developed to the blastocyst stage (28, 99, 102-106). However different observations were reported by other studies (76, 99, 107).

Ferguson et al. (97) have demonstrated that addition of physiological concentrations of P4 to embryo culture medium at three days post-insemination benefitted embryo development in several ways. Thus, they have concluded that P4 has a direct positive effect on the developing *in vitro* culture of bovine embryos. P4 supplementation increased the number of *in vitro* culture embryos that developed to the grade 1 blastocyst stages as well as the number of hatched blastocysts.

It was shown that co-culture of an embryo with endometrial tissue cultured in the presence of P4 and E2 benefitted embryo development (108).

Also, our investigations led to similar results in which *in vitro* culture of mouse 2-cell embryos in the presence of 20 ng/ml P4 resulted in a high proportion of embryos that reached the blastocyst stage. In this study, embryo quality was less affected by P4 (28).

Conclusion

These results have shown that P4 could be a factor for embryonic survival and improve *in vivo* embryo development and implantation, both directly and indirectly, on its concentration and the mammalian species. The effect of P4 on oocyte maturation and embryo development may be dependent on its concentration and the mammalian species.

References

1. Tsafiri A, Cao X, Ashkenazi H, Motola S, Popliker M, Pomerantz SH. Resumption of oocyte meiosis in mammals: on models, meiosis activating sterols, steroids and EGF-like factors. *Mol Cell Endocrinol.* 2005; 234 (1-2): 37-45.
2. Lonergan P. Influence of progesterone on oocyte quality and embryo development in cows. *Theriogenology.* 2011; 76(9): 1594-1601.
3. Byskov AG, Yding Andersen C, Hossaini A, Guoliang X. Cumulus cells of oocyte-cumulus complexes secrete a mei-

- osis-activating substance when stimulated with FSH. *Mol Reprod Dev.* 1997; 46(3): 296-305.
4. Gandolfi F. Intra-ovarian regulation of oocyte developmental competence in cattle. *Zygote.* 1996; 4(4): 323-326.
 5. Andersen CY. Levels of steroid-binding proteins and steroids in human preovulatory follicle fluid and serum as predictors of success in in vitro fertilization-embryo transfer treatment. *J Clin Endocrinol Metab.* 1990; 71(5): 1375-1381.
 6. Mendoza C, Ruiz-Requena E, Ortega E, Cremades N, Martinez F, Bernabeu R, et al. Follicular fluid markers of oocyte developmental potential. *Hum Reprod.* 2002; 17(4): 1017-1022.
 7. Flores-Herrera H, Diaz-Cervantes P, De la Mora G, Zaga-Clavellina V, Uribe-Salas F, Castro I. A possible role of progesterone receptor in mouse oocyte in vitro fertilization regulated by norethisterone and its reduced metabolite. *Contraception.* 2008; 78(6): 507-512.
 8. Eppig JJ, Schultz RM, O'Brien M, Chesnel F. Relationship between the developmental programs controlling nuclear and cytoplasmic maturation of mouse oocytes. *Dev Biol.* 1994; 164(1): 1-9.
 9. Wassarman PM, Litscher ES. Mammalian fertilization is dependent on multiple membrane fusion events. *Methods Mol Biol.* 2008; 475: 99-113.
 10. Mahesh VB, Brann DW, Hendry LB. Diverse modes of action of progesterone and its metabolites. *J Steroid Biochem Mol Biol.* 1996; 56 (1-6 Spec No): 209-219.
 11. Golub MS, Kaufman FL, Campbell MA, Li LH, Donald JM. "Natural" progesterone: information on fetal effects. *Birth Defects Res B Dev Reprod Toxicol.* 2006; 77(5): 455-470.
 12. Raber L. Steroid industry honored. *ACS News.* 1999; 77: 78-80.
 13. Jones RE, Lopez KH. The female reproductive system. In: Maragoglio N, editor. *Human reproductive biology.* 3rd ed. USA: Academic Press; 2006; 37.
 14. Simon JA. Micronized progesterone: vaginal and oral uses. *Clin Obstet Gynecol.* 1995; 38(4): 902-914.
 15. Graham JD, Clarke CL. Physiological action of progesterone in target tissues. *Endocr Rev.* 1997; 18(4): 502-519.
 16. Bradshaw FJ, Bradshaw D. Progesterone and reproduction in marsupials: a review. *Gen Comp Endocrinol.* 2011; 170(1): 18-40.
 17. Su YQ, Sugiura K, Woo Y, Wigglesworth K, Kamdar S, Affourtit J, et al. Selective degradation of transcripts during meiotic maturation of mouse oocytes. *Dev Biol.* 2007; 302(1): 104-117.
 18. Letterie GS. A regimen of oral contraceptives restricted to the periovulatory period may permit folliculogenesis but inhibit ovulation. *Contraception.* 1998; 57(1): 39-44.
 19. Barrera D, Avila E, Diaz L. Immunological role of progesterone in the maintenance of pregnancy. *Rev Invest Clin.* 2007; 59(2): 139-145.
 20. Sengupta J, Ghosh D. Role of progesterone on peri-implantation stage endometrium-embryo interaction in the primate. *Steroids.* 2000; 65(10-11): 753-762.
 21. Loutradis D, Beretsos P, Arabatzis E, Anagnostou E, Drakakis P. The role of steroid hormones in ART. *J Steroid Biochem Mol Biol.* 2008; 112 (1-3): 1-4.
 22. Nyboe Andersen A, Popovic-Todorovic B, Schmidt KT, Loft A, Lindhard A, Hojgaard A, et al. Progesterone supplementation during early gestations after IVF or ICSI has no effect on the delivery rates: a randomized controlled trial. *Hum Reprod.* 2002; 17(2): 357-361.
 23. Gillet JY, Andre G, Faguer B, Erny R, Buvat-Herbaut M, Domin MA, et al. Induction of amenorrhea during hormone replacement therapy: optimal micronized progesterone dose. A multicenter study. *Maturitas.* 1994; 19(2): 103-115.
 24. Chhabra S, McCartney CR, Yoo RY, Eagleson CA, Chang RJ, Marshall JC. Progesterone inhibition of the hypothalamic gonadotropin-releasing hormone pulse generator: evidence for varied effects in hyperandrogenemic adolescent girls. *J Clin Endocrinol Metab.* 2005; 90(5): 2810-2815.
 25. Salehnia M. Different pattern of pinopodes expression in stimulated mouse endometrium. *Exp Anim.* 2005; 54(4): 349-352.
 26. Arianmanesh M, Salehnia M, Niknafs B. The effect of progesterone after hyperstimulation on the ultrastructure of mouse endometrium during preimplantation period. *Yakhteh.* 2002; 4(14): 61-67.
 27. Salehnia M. Progesterone shifts the pinopodes expression of mouse endometrium to pre-implantation time after ovarian hyperstimulation. *Iran J Reprod Med.* 2003; 1(1): 20-23.
 28. Ghaemi SR, Salehnia M, Valojerdi MR. The effect of progesterone and exogenous gonadotropin on preimplantation mouse embryo development and implantation. *Exp Anim.* 2008; 57(1): 27-34.
 29. Peyghambari F, Salehnia M, Forouzandeh Moghadam M, Rezaadeh Valoujerdi M, Hajizadeh E. The changes in morphology and morphometrical indexes of endometrium of ovariectomized mice in response to exogenous ovarian hormones. *Iran J Reprod Med.* 2008; 6(3): 125-131.
 30. Peyghambari F, Salehnia M, Forouzandeh Moghadam M, Rezaadeh Valoujerdi M, Hajizadeh E. The correlation between the endometrial integrins and osteopontin expression with pinopodes development in ovariectomized mice in response to exogenous steroids hormones. *Iran Biomed J.* 2010; 14(3): 109-119.
 31. Rekawiecki R, Kowalik MK, Kotwica J. Nuclear progesterone receptor isoforms and their functions in the female reproductive tract. *Pol J Vet Sci.* 2011; 14(1): 149-158.
 32. Dressing GE, Goldberg JE, Charles NJ, Schwertfeger KL, Lange CA. Membrane progesterone receptor expression in mammalian tissues: a review of regulation and physiological implications. *Steroids.* 2011; 76(1-2): 11-17.
 33. Tsafiri A, Motola S. Are steroids dispensable for meiotic resumption in mammals? *Trends Endocrinol Metab.* 2007; 18(8): 321-327.
 34. Franco HL, Jeong JW, Tsai SY, Lydon JP, DeMayo FJ. In vivo analysis of progesterone receptor action in the uterus during embryo implantation. *Semin Cell Dev Biol.* 2008; 19(2): 178-186.
 35. Hild-Petito S, Stouffer RL, Brenner RM. Immunocytochemical localization of estradiol and progesterone receptors in the monkey ovary throughout the menstrual cycle. *Endocrinology.* 1988; 123(6): 2896-2905.
 36. Gava N, Clarke CL, Byth K, Arnett-Mansfield RL, deFazio A. Expression of progesterone receptors A and B in the mouse ovary during the estrous cycle. *Endocrinology.* 2004; 145(7): 3487-3494.
 37. Teilmann SC, Clement CA, Thorup J, Byskov AG, Christensen ST. Expression and localization of the progesterone receptor in mouse and human reproductive organs. *J Endocrinol.* 2006; 191(3): 525-535.
 38. Conneely OM, Mulac-Jericevic B, Lydon JP. Progesterone-dependent regulation of female reproductive activity by two distinct progesterone receptor isoforms. *Steroids.* 2003; 68(10-13): 771-778.
 39. Aparicio IM, Garcia-Herreros M, O'shea LC, Hensey C, Lonergan P, Fair T. Expression, regulation, and function of progesterone receptors in bovine cumulus oocyte complexes during in vitro maturation. *Biol Reprod.* 2011; 84(5): 910-921.
 40. Shimada M, Yamashita Y, Ito J, Okazaki T, Kawahata K, Nishibori M. Expression of two progesterone receptor isoforms in cumulus cells and their roles during meiotic resumption of porcine oocytes. *J Mol Endocrinol.* 2004; 33(1): 209-225.

41. Goncalves JS, Vannucchi CI, Braga FC, Paula-Lopes FF, Milazzotto MP, Assumpcao ME, et al. Oestrogen and progesterone receptor gene expression in canine oocytes and cumulus cells throughout the oestrous cycle. *Reprod Domest Anim.* 2009; 44 Suppl 2: 239-242.
42. Vermeirsch H, Simoens P, Coryn M, Van den Broeck W. Immunolocalization of progesterone receptors in the canine ovary and their relation to sex steroid hormone concentrations. *Reproduction.* 2001; 122(1): 73-83.
43. Bayaa M, Booth RA, Sheng Y, Liu XJ. The classical progesterone receptor mediates *Xenopus* oocyte maturation through a nongenomic mechanism. *Proc Natl Acad Sci USA.* 2000; 97(23): 12607-12612.
44. Peluso JJ, Pappalardo A, Losel R, Wehling M. Progesterone membrane receptor component 1 expression in the immature rat ovary and its role in mediating progesterone's antiapoptotic action. *Endocrinology.* 2006; 147(6): 3133-3140.
45. Engmann L, Losel R, Wehling M, Peluso JJ. Progesterone regulation of human granulosa/luteal cell viability by an RU486-independent mechanism. *J Clin Endocrinol Metab.* 2006; 91(12): 4962-4968.
46. Luciano AM, Lodde V, Franciosi F, Cecilian F, Peluso JJ. Progesterone receptor membrane component 1 expression and putative function in bovine oocyte maturation, fertilization, and early embryonic development. *Reproduction.* 2010; 140(5): 663-672.
47. Wood JR, Dumesic DA, Abbott DH, Strauss JF3rd. Molecular abnormalities in oocytes from women with polycystic ovary syndrome revealed by microarray analysis. *J Clin Endocrinol Metab.* 2007; 92(2): 705-713.
48. Nousiainen M, Sillje HH, Sauer G, Nigg EA, Korner R. Phosphoproteome analysis of the human mitotic spindle. *Proc Natl Acad Sci USA.* 2006; 103(14): 5391-5396.
49. Rambags BP, van Tol HT, van den Eng MM, Colenbrander B, Stout TA. Expression of progesterone and oestrogen receptors by early intrauterine equine conceptuses. *Theriogenology.* 2008; 69(3): 366-375.
50. Hasegawa J, Yanaiharu A, Iwasaki S, Otsuka Y, Negishi M, Akahane T, et al. Reduction of progesterone receptor expression in human cumulus cells at the time of oocyte collection during IVF is associated with good embryo quality. *Hum Reprod.* 2005; 20(8): 2194-2200.
51. Hou Q, Gorski J. Estrogen receptor and progesterone receptor genes are expressed differentially in mouse embryos during preimplantation development. *Proc Natl Acad Sci USA.* 1993; 90(20): 9460-9464.
52. Hou Q, Paria BC, Mui C, Dey SK, Gorski J. Immunolocalization of estrogen receptor protein in the mouse blastocyst during normal and delayed implantation. *Proc Natl Acad Sci USA.* 1996; 93(6): 2376-2381.
53. Hong SH, Nah HY, Lee YJ, Lee JW, Park JH, Kim SJ, et al. Expression of estrogen receptor-alpha and -beta, glucocorticoid receptor, and progesterone receptor genes in human embryonic stem cells and embryoid bodies. *Mol Cells.* 2004; 18(3): 320-325.
54. Ying C, Yang YC, Hong WF, Cheng WT, Hsu WL. Progesterone receptor gene expression in preimplantation pig embryos. *Eur J Endocrinol.* 2000; 143(5): 697-703.
55. Kowalski AA, Graddy LG, Vale-Cruz DS, Choi I, Katzenellenbogen BS, Simmen FA, et al. Molecular cloning of porcine estrogen receptor-beta complementary DNAs and developmental expression in periimplantation embryos. *Biol Reprod.* 2002; 66(3): 760-769.
56. Bagchi IC, Li Q, Cheon YP, Mantena SR, Kannan A, Bagchi MK. Use of the progesterone receptor antagonist RU 486 to identify novel progesterone receptor-regulated pathways in implantation. *Semin Reprod Med.* 2005; 23(1): 38-45.
57. Clemente M, de La Fuente J, Fair T, Al Naib A, Gutierrez-Adan A, Roche JF, et al. Progesterone and conceptus elongation in cattle: a direct effect on the embryo or an indirect effect via the endometrium? *Reproduction.* 2009; 138(3): 507-517.
58. Okada A, Ohta Y, Buchanan D, Sato T, Iguchi T. Effect of estrogens on ontogenetic expression of progesterone receptor in the fetal female rat reproductive tract. *Mol Cell Endocrinol.* 2002; 195(1-2): 55-64.
59. Shanker YG, Sharma SC, Rao AJ. Expression of progesterone receptor mRNA in the first trimester human placenta. *Biochem Mol Biol Int.* 1997; 42(6): 1235-1240.
60. Morrison T, Waggoner L, Whitworth-Langley L, Stith BJ. Nongenomic action of progesterone: activation of *Xenopus* oocyte phospholipase C through a plasma membrane-associated tyrosine kinase. *Endocrinology.* 2000; 141(6): 2145-2152.
61. Moor RM, Polge C, Willadsen SM. Effect of follicular steroids on the maturation and fertilization of mammalian oocytes. *J Embryol Exp Morphol.* 1980; 56: 319-335.
62. Zavareh S, Saberivand A, Salehnia M. The effect of progesterone on the in vitro maturation and developmental competence of mouse germinal vesicle oocytes. *Int J Fertil Steril.* 2009; 3(1): 21-28.
63. Vanderhyden BC, Tonary AM. Differential regulation of progesterone and estradiol production by mouse cumulus and mural granulosa cells by A factor(s) secreted by the oocyte. *Biol Reprod.* 1995; 53(6): 1243-1250.
64. Jamnongjit M, Gill A, Hammes SR. Epidermal growth factor receptor signaling is required for normal ovarian steroidogenesis and oocyte maturation. *Proc Natl Acad Sci USA.* 2005; 102(45): 16257-16262.
65. Fukui Y, Fukushima M, Terawaki Y, Ono H. Effect of gonadotropins, steroids and culture media on bovine oocyte maturation in vitro. *Theriogenology.* 1982; 18(2): 161-175.
66. Carter F, Rings F, Mamo S, Holker M, Kuzmany A, Besenfelder U, et al. Effect of elevated circulating progesterone concentration on bovine blastocyst development and global transcriptome following endoscopic transfer of in vitro produced embryos to the bovine oviduct. *Biol Reprod.* 2010; 83(5): 707-719.
67. Willingham-Rocky LA, Hinrichs K, Westhusin ME, Kraemer DC. Effects of stage of oestrous cycle and progesterone supplementation during culture on maturation of canine oocytes in vitro. *Reproduction.* 2003; 126(4): 501-508.
68. Vannucchi CI, de Oliveira CM, Marques MG, Assumpcao ME, Visintin JA. In vitro canine oocyte nuclear maturation in homologous oviductal cell co-culture with hormone-supplemented media. *Theriogenology.* 2006; 66(6-7): 1677-1681.
69. Hewitt DA, England GC. Effect of preovulatory endocrine events upon maturation of oocytes of domestic bitches. *J Reprod Fertil Suppl.* 1997; 51: 83-91.
70. Duval D, Durant S, Homo-Delarche F. Non-genomic effects of steroids. Interactions of steroid molecules with membrane structures and functions. *Biochim Biophys Acta.* 1983; 737(3-4): 409-442.
71. Kaji E, Bornslaeger EA, Schultz RM. Inhibition of mouse oocyte cyclic AMP phosphodiesterase by steroid hormones: a possible mechanism for steroid hormone inhibition of oocyte maturation. *J Exp Zool.* 1987; 243(3): 489-493.
72. Morgan PM, Boatman DE, Bavister BD. Relationships between follicular fluid steroid hormone concentrations, oocyte maturity, in vitro fertilization and embryonic development in the rhesus monkey. *Mol Reprod Dev.* 1990; 27(2): 145-151.
73. Karlach V. The effect of FSH, LH, oestradiol-17 beta, and progesterone on cytoplasmic maturation of bovine follicular oocytes in vitro. *Folia Biol (Praha).* 1987; 33(4): 258-265.
74. Osborn JC, Moor RM, Crosby IM. Effect of alterations in follicular steroidogenesis on the nuclear and cytoplasmic

- maturation of ovine oocytes. *J Embryol Exp Morphol.* 1986; 98: 187-208.
75. Zheng P, Si W, Bavister BD, Yang J, Ding C, Ji W. 17Beta-estradiol and progesterone improve in-vitro cytoplasmic maturation of oocytes from unstimulated prepubertal and adult rhesus monkeys. *Hum Reprod.* 2003; 18(10): 2137-2144.
 76. Silva CC, Knight PG. Effects of androgens, progesterone and their antagonists on the developmental competence of in vitro matured bovine oocytes. *J Reprod Fertil.* 2000; 119(2): 261-269.
 77. Mattioli M, Galeati G, Bacci ML, Seren E. Follicular factors influence oocyte fertilizability by modulating the intercellular cooperation between cumulus cells and oocyte. *Gamete Res.* 1988; 21(3): 223-232.
 78. Zhang X, Armstrong DT. Effects of follicle-stimulating hormone and ovarian steroids during in vitro meiotic maturation on fertilization of rat oocytes. *Gamete Res.* 1989; 23(3): 267-277.
 79. Martal J, Chene N, Camous S, Huynh L, Lantier F, Hermier P, et al. Recent developments and potentialities for reducing embryo mortality in ruminants: the role of IFN-tau and other cytokines in early pregnancy. *Reprod Fertil Dev.* 1997; 9(3): 355-380.
 80. Tian J, Kim S, Heilig E, Ruderman JV. Identification of XPR-1, a progesterone receptor required for *Xenopus* oocyte activation. *Proc Natl Acad Sci USA.* 2000; 97(26): 14358-14363.
 81. Robertson SA. GM-CSF regulation of embryo development and pregnancy. *Cytokine Growth Factor Rev.* 2007; 18(3-4): 287-298.
 82. Sheikholslami B, Salehnia M, Valojerdi MR, Ramezanzadeh M. Developmental potential of isolated blastomeres from early mouse embryos in the presence and absence of LIF and GM-CSF. *J Assist Reprod Genet.* 2008; 25(1): 7-12.
 83. Lessey BA, Ilesanmi AO, Castelbaum AJ, Yuan L, Somkuti SG, Chwalisz K, et al. Characterization of the functional progesterone receptor in an endometrial adenocarcinoma cell line (Ishikawa): progesterone-induced expression of the alpha1 integrin. *J Steroid Biochem Mol Biol.* 1996; 59(1): 31-39.
 84. Mann GE, Lamming GE, Robinson RS, Wathes DC. The regulation of interferon-tau production and uterine hormone receptors during early pregnancy. *J Reprod Fertil Suppl.* 1999; 54: 317-328.
 85. Walton JS, Halbert GW, Robinson NA, Leslie KE. Effects of progesterone and human chorionic gonadotrophin administration five days postinsemination on plasma and milk concentrations of progesterone and pregnancy rates of normal and repeat breeder dairy cows. *Can J Vet Res.* 1990; 54(3): 305-308.
 86. Garrett JE, Geisert RD, Zavy MT, Gries LK, Wettemann RP, Buchanan DS. Effect of exogenous progesterone on prostaglandin F2 alpha release and the interestrus interval in the bovine. *Prostaglandins.* 1988; 36(1): 85-96.
 87. Bazer FW, Spencer TE, Johnson GA, Burghardt RC. Uterine receptivity to implantation of blastocysts in mammals. *Front Biosci (Schol Ed).* 2011; 3: 745-767.
 88. Szekeres-Bartho J, Halasz M, Palkovics T. Progesterone in pregnancy; receptor-ligand interaction and signaling pathways. *J Reprod Immunol.* 2009; 83(1-2): 60-64.
 89. Daya S. Luteal support: progestogens for pregnancy protection. *Maturitas.* 2009; 65 Suppl 1: S29-34.
 90. Bazer FW, Spencer TE, Johnson GA, Burghardt RC, Wu G. Comparative aspects of implantation. *Reproduction.* 2009; 138(2): 195-209.
 91. Mann GE, Fray MD, Lamming GE. Effects of time of progesterone supplementation on embryo development and interferon-tau production in the cow. *Vet J.* 2006; 171(3): 500-503.
 92. Mann GE, Lamming GE. Relationship between maternal endocrine environment, early embryo development and inhibition of the luteolytic mechanism in cows. *Reproduction.* 2001; 121(1): 175-180.
 93. Sohn SH, Penzias AS, Emmi AM, Dubey AK, Layman LC, Reindollar RH, et al. Administration of progesterone before oocyte retrieval negatively affects the implantation rate. *Fertil Steril.* 1999; 71(1): 11-14.
 94. Song H, Han K, Lim H. Progesterone supplementation extends uterine receptivity for blastocyst implantation in mice. *Reproduction.* 2007; 133(2): 487-493.
 95. Beltman ME, Lonergan P, Diskin MG, Roche JF, Crowe MA. Effect of progesterone supplementation in the first week post conception on embryo survival in beef heifers. *Theriogenology.* 2009; 71(7): 1173-1179.
 96. Harini C, Sainath SB, Reddy PS. Progesterone administration induces preimplantation embryonic loss in mice. *Fertil Steril.* 2009; 91(5 Suppl): 2137-2141.
 97. Ferguson CE, Davidson TR, Mello MR, Lima AS, Kesler DJ, Wheeler MB, et al. Evidence of a direct effect of P4 on IVF-derived bovine 8-cell embryos. *Reprod Fertil Dev.* 2005; 17(2): 219.
 98. Merlo B, Iacono E, Mari G. Effect of progesterone and epidermal growth factor on in vitro-produced eight-cell bovine embryos in serum-free culture medium. *Reprod Fertil Dev.* 2007; 19(1): 211.
 99. Goff AK, Smith LC. Effect of steroid treatment of endometrial cells on blastocyst development during co-culture. *Theriogenology.* 1998; 49(5): 1021-1030.
 100. Reggio BC, Lynn JW, Godke RA. The effect of progesterone on the development of IVF-derived bovine embryos cultured in a semi-defined culture medium. *Theriogenology.* 1997; 47(1): 284.
 101. Geisert RD, Fox TC, Morgan GL, Wells ME, Wettemann RP, Zavy MT. Survival of bovine embryos transferred to progesterone-treated asynchronous recipients. *J Reprod Fertil.* 1991; 92(2): 475-482.
 102. Shimada M, Kawano N, Terada T. Delay of nuclear maturation and reduction in developmental competence of pig oocytes after mineral oil overlay of in vitro maturation media. *Reproduction.* 2002; 124(4): 557-564.
 103. Forde N, Beltman ME, Duffy GB, Duffy P, Mehta JP, O'Gaora P, et al. Changes in the endometrial transcriptome during the bovine estrous cycle: effect of low circulating progesterone and consequences for conceptus elongation. *Biol Reprod.* 2011; 84(2): 266-278.
 104. Diskin MG, Murphy JJ, Sreenan JM. Embryo survival in dairy cows managed under pastoral conditions. *Anim Reprod Sci.* 2006; 96(3-4): 297-311.
 105. Lamb GC, Dahlen CR, Larson JE, Marquezini G, Stevenson JS. Control of the estrous cycle to improve fertility for fixed-time artificial insemination in beef cattle: a review. *J Anim Sci.* 2010; 88 Suppl 13: E181-192.
 106. Ryan GJ, Waddington D, Campbell KHS. Addition of progesterone during bovine oocyte maturation in the presence of gonadotrophins improves developmental competence. *Theriogenology.* 1999; 51(1): 392.
 107. Wiemer KE, Amborski GF, Denniston RS, White KL, Godke RA. Use of a hormone-treated fetal uterine fibroblast monolayer system for in vitro culture of bovine embryos. *Theriogenology.* 1987; 27(1): 294.
 108. Lavranos TC, Seamark RF. Addition of steroids to embryoterpine monolayer co-culture enhances embryo survival and implantation in vitro. *Reprod Fertil Dev.* 1989; 1(1): 41-46.