

*Review Article*

Production of Lactic Acid from Various Feed Stocks: A Review on Studies and Research

Sunil J. Kulkarni

Datta Meghe College of Engineering, Airoli, Navi Mumbai, Maharashtra, India.

*Received: 05/12/2015**Revised: 19/12/2015**Accepted: 19/12/2015*

ABSTRACT

In the present scenario of increasing population and advancements in technologies, reduction in the production cost has become important to the chemical manufacturing sector. The use of biological methods and application of biotechnology can be very effective way to produce some chemicals. Production of chemicals such as Lactic acid, citric acid, acetic acid, and ethanol can be done by using biological processes such as fermentation. The current review focuses on production of Lactic acid. Various investigators have carried out research by using different raw materials and bacteria. The present review summarizes research carried out by various investigators for production of Lactic acid by using various biological species and raw materials.

Key words: Biomass, concentration, raw material, nutrients, substrates.

INTRODUCTION

In the era of cutting edge technology, production of various chemical by using low cost and environment friendly technology is the need of the society for sustainable development. Process intensification methods such as hydrotropy, extractive distillation solar distillation can be used to optimize and economize the process. [1-3] Two main aspects of environmental friendly process are temperature and effluent emission. The effluent can be treated by various chemical, biological and physical methods. [4-7] Manufacturing of the chemicals by using biological pathways has advantages such as high specificity, moderate temperature and low energy requirement. Biocatalysts have been used effectively for production of various chemicals. [8-11] The production of chemicals such as acetic acid, ethanol and

starch has been reported by various investigators. [12-15]

RESEARCH AND STUDIES ON PRODUCTION OF LACTIC ACID

Hosamane et.al carried out research on Lactic acid production from cane sugar molasses. [16] They employed the response surface methodology to make mathematical modeling and optimization of the temperature and shaking time. They observed that two variables had mutually dependent influence on the % of Lactic acid recovered. They also found that maximum % of Lactic acid recovered was 0.65. Goranov et.al investigated the Lactic acid production by encapsulated *Lactobacillus casei* ssp. *rhamnosus* ATCC 11979 cells in alginate/chitosan complexes with solid and liquid core. [17] They compared the Lactic acid biosynthesis with immobilized cell. The free and the immobilized cells produced maximal

Lactic acid quantities at pH 6.5. In their investigation they also determined dynamics of substrate uptake, accumulation of Lactic acid, pH and redox potential of the fermentation medium. They observed that the temperature increase above 42°C led to a decrease in the cells' catalytic activity.

Bayitse reviewed prospect for bioresidue utilization in Ghana for Lactic Acid production from biomass. [18] They studied current research in Lactic acid fermentation processes, bio-residue availability in Ghana and potential utilization for Lactic acid production. They found that both starchy and lignocellulosic biomass has been extensively used. Materials such as maize cobs, millet stalk, sorghum stalk and rice straw were found to be excellent feed sticks for Lactic acid production.

Mudaliyar and Kulkarni studied the development of a bioprocess to produce Lactic acid using dry grass, coconut husk, sugarcane waste and wood chips as substrate. [19] They found that by using *Rhizopus* genus, many renewable resources including refined sugars, molasses, raw starch materials and even lignocellulose can be converted to Lactic acid. Wee et.al carried out a review on Lactic acid-producing microorganisms, raw materials for Lactic acid production, fermentation approaches for Lactic acid production. [20] They also studied various applications of Lactic acid. According to their estimates, the worldwide consumption of Lactic acid is estimated to be 130 000-150 000 (metric) tonnes per year. According to them, there are still several issues that need to be addressed in order to produce Lactic acid biotechnologically within the targeted cost. They concluded that the biotechnological processes for the production of Lactic acid from cheap raw materials should be improved further to make them competitive with the chemically-derived one.

Farooq et.al carried out investigation on production of Lactic acid by *Lactobacillus delbrueckii* using food wastes as substrates. [21] They observed that the productivity was affected by fermentation time, temperature and the level of substrate. They achieved maximum Lactic acid production after 7 days of fermentation in media containing 18% substrate level. The maximum recovery obtained was 78-80 percent. Corn by-product raw materials were used for production of Lactic acid by Minh et.al [22] They studied various factors affecting Lactic acid and optimized them. They obtained optimum parameter values as 5% glucose concentration, pH value 6, time 84 hours. At these conditions, they obtained highest concentration of Lactic acid 19.07 (g/l). Their research was mainly focused on biotransformation of xenuloza in raw materials. They concluded that it was necessary to perform further studies to not only reduce sugar (calculated by glucose) but also sugar5-carbon and 6-carbon.

Jahromi et.al carried out investigation on Lactic acid fermentation. [23] They used Different *Lactobacillus* Species using sorghum seed extract as a carbon source. They observed that, among the selected bacterial species, *Lactobacillus delbrueckii* was having unique characteristics. It could tolerate different physico-chemical conditions. They concluded that hydrolyzed sorghum seed extract has enough nutrients to support the efficient growth of Lactic acid bacteria and production of Lactic acid under favorable conditions.

Munyon and Merchant investigated the relation between glucose utilization and Lactic acid production. [24] They also studied utilization and the growth cycle of I strain fibroblasts. In their research, they attempted to follow carbohydrate metabolism by measurements of glucose and lactate levels at successive periods of growth. They observed that the pH of the medium decreased from 7.3 to 6.8 during

the period of linear increase in cell numbers. Also they observed that there was no constant relation between the rate of glucose uptake with growth. Zhang and Jin carried out investigation on alternative method for L (+)-Lactic acid production using sugarcane molasses and waste potato starch. [25] According to them, the main disadvantage of using bacteria is that expensive nutrients such as yeast extract and peptone are required. They examined the production of Lactic acid from sugarcane molasses and waste potato starch by the fungus *Rhizopus arrhizus*. They found that adding extra nutrients into molasses did not improve Lactic acid yield.

Xue et.al carried out investigation on production of Lactic acid by thermophilic *Bacillus* sp. strain XZL4 from corn stover hydrolyzate. [26] They obtained the highest Lactic acid concentration (81.0 g L⁻¹) and yield (0.98 g g⁻¹ total reducing sugars) from corn stover hydrolyzate. They also found that *Bacillus* sp. XZL4 was a promising polymer-grade L-Lactic-acid producer from cellulosic biomass. Fermentation processes using Lactic acid bacteria (LAB) producing bacteriocins for preservation were carried out by Juodeikiene et.al. [27] According to them, the use of bacteriocins and/or bacteriocin-producing strains of LAB are of great interest. They are generally recognized as safe organisms and their antimicrobial products as biopreservatives. They emphasized that there is a need for practical technology controlled from an economical point of view.

Cell-recycle repeated-batch bioreactor was used for Lactic acid production by Oh et.al. [28] They investigated the effect of various nitrogen sources on cell growth and Lactic acid production. They found that increasing the yeast extract content raises the total production cost of Lactic acid. They tried to find the optimum yeast extract dosage

for a repeated-batch operation with cell recycling. Narayanan et.al carried out investigation on L (+) Lactic acid fermentation. [29] They also studied its product polymerization. They reviewed the microorganisms being used for Lactic acid fermentation, the raw materials reported, the various novel fermentation processes and its processing methods. They found that varieties of raw materials can be used for the Lactic acid production.

Sobrun et.al carried out investigation on isolation of Lactic acid bacteria from sugar cane juice. [30] During their studies, they observed that several isolates exhibited a clear zone and growth on de Man, Rogosa, Sharpe (MRS) agar supplemented with sodium azide, bromocresol purple and sucrose. They isolated a total of 112 mutants. Out of these 9 homofermentative isolates were further investigated for their ability to produce Lactic acid. They observed that all mutant isolates produced Lactic acid as the sole fermentation product. Shiphrah et.al carried out research on Lactic acid production from whey water. [31] They used a by-product after the precipitation of cheese as a substrate. They observed that the various milk products and milk itself has the maximum content of Lactic acid bacteria. They concluded that the potential of using this isolate for bioremediation of dairy waste (whey water) could be explored and exploited in future.

Wang et.al carried out studies on corncob molasses containing a high content of xylose for Lactic acid production. [32] They obtained highest Lactic acid concentration when initial corncob molasses concentration was 150 g/l. Thereafter sharp decrease was observed in Lactic acid production. According to them, the production of Lactic acid from various by-products or agricultural residues has gained considerable interest because of the increasing interest in producing biotechnological products from low-cost

and renewable biomass. Kivanc et.al isolated a total of 45 Lactic acid bacteria from 10 different boza samples consumed in Turkey. [33] They tested isolates for inhibitory activity against food-borne bacterial pathogens. They determined antimicrobial effects of these Lactic acid bacteria. They observed that Lactic acid bacteria strains examined in this study were rather moderate acid producers.

Hadadji and Bensoltane carried out an investigation on bifidobacterium longum and Lactobacillus acidophilus in goat's milk for growth and Lactic acid production. [34] They compared the production of Lactic acid with pure and mixed cultures. They observed that the maximum rate of Lactic acid production was obtained with mixed culture at 45°C. Maas et.al carried out investigation on Lactic acid production from xylose. [35] They used the fungus Rhizopus oryzae bacteria for the purpose. Their investigation resulted in the conversion of xylose in synthetic media into Lactic acid by ten R. oryzae strains resulting in yields between 0.41 and 0.71 g g⁻¹. Diauxic growth was observed in case of mixed substrates. Afolabi demonstrated a fermentation process, which would economically produce Lactic acid from inexpensive carbon source. [36] First, they inoculated the spores of lactobacillus casei to the growth medium and then changed the medium to production medium. They observed that with glucose concentration, The Lactic acid production increased. Hundred percent conversions could not be achieved as a result of Lactic acid inhibition. They also found that the inhibitory effect can be minimized by using certain additives such as ammonium sulphate and manganese sulphate. Ray et.al carried out research on use of fibrous residue obtained during extraction of starch from cassava for Lactic acid production. [37] They carried out semi-solid state fermentation using Mann Rogassa Sharpe medium in lieu of glucose

[2 % (wv⁻¹)] as the carbon source. They used response surface methodology (RSM) to evaluate the effect of main variables. The important variable includes incubation period, temperature and pH. The optimum values of these variables were obtained to be 120 hr, 35°C and 6.5, respectively during their investigation.

CONCLUSION

The review suggests that starchy and lignocellulosic biomass has been extensively used for Lactic acid production. Various researchers have reported Lactic acid production from various feed stocks such as agricultural waste, corncob, dairy industry waste, dry grass, coconut husk, sugarcane waste and wood chips. In order to make the Lactic acid fermentation more attractive more research on affecting parameters, bacteria and feed stocks is required.

REFERENCES

1. Sunil Jayant Kulkarni, Process Intensification and Nanomaterials: A Short Review, International Journal of Research, 2014, 1(9), 392-394.
2. Sunil Jayant Kulkarni, Ajaygiri Kamalgiri Goswami, Research on Application of Hydrotropy: A Review, International Journal of Science, Engineering and Technology Research (IJSETR), 2014, 3(10) 2617-2619.
3. Sunil Jayant Kulkarni, A Review on Research and Advancements in Extractive Distillation, International Journal of Research (IJR), 2015, 2(1), 306-309.
4. Sunil J. Kulkarni, Modeling for Adsorption Columns for Wastewater Treatment: a Review, International Journal of Innovative Research in Engineering & Multidisciplinary Physical Sciences, 2014, 2(2), 7-11.
5. Sunil Jayant Kulkarni, A Review on Packed Bed Removal of Organic Matter from Wastewater, Int. Journal on Scientific Research in Science, Engineering and Technology, 2015, 1 (2), 27-30.
6. Sunil J. Kulkarni, Removal Of Organic Matter From Domestic Waste Water By

- Adsorption, International Journal Of Science, Engineering And Technology Research (IJSETR), 2013, 2(10), 1836-1839.
7. Sunil J. Kulkarni, Ajaygiri K. Goswami, Applications and Advancements in Treatment of Waste Water by Membrane Technology- A Review, International Journal Of Engineering Sciences & Research Technology, 2014, 3(9), 446-450.
 8. Sunil Jayant Kulkarni, Research on Biocatalysts: A Review, International Journal of Research (IJR), 2015, 2(4), 784-788.
 9. Martin H. Fechter and Herfried Griengl, Hydroxynitrile Lyases: Biological Sources and Application as Biocatalysts, Food Technol., 2004, 42(2), 287-294.
 10. Paul A. Bird, John M. Woodley and David C.A. Sharp, Using Biocatalysts at Useful Reactant and Product Concentrations Monitoring and Controlling Biocatalytic Processes, BioPharm International, 2002, 14-21.
 11. Monika Gorak, Ewa Zymanczyk-Duda, Application of cyanobacteria as biocatalysts for the reduction of diethyl 2-oxopropylphosphonate, Chemik, 2014, 68(2), 123-128.
 12. Sunil J. Kulkarni, Nilesh L. Shinde, Ajaygiri K. Goswami, A Review on Ethanol Production from Agricultural Waste Raw Material, International Journal of Scientific Research in Science, Engineering and Technology, 2015, 1(4), 231-233.
 13. Veena Ramachandran, Nisha Pujari, Tanmay Matey, Sunil Kulkarni, Enzymatic Hydrolysis for Glucose-A Review, International Journal of Science, Engineering and Technology Research (IJSETR), 2013, 2(10), 1937-1942.
 14. Sunil J. Kulkarni, Use of Biotechnology for Synthesis of Various Products from Different Feedstocks -A Review, International Journal of Advanced Research in Bio-Technology, 2014, 2(2), 1-3.
 15. Veena Ramachandran, Nisha Pujari, Tanmay Matey, Sunil Kulkarni, Enzymatic Hydrolysis of Cassava using wheat Seedlings, International Journal of Science, Engineering and Technology Research (IJSETR), 2014, 3(5), 1216-1219.
 16. Sateesh N Hosamane, Shrikrishna H Gurlhosur And Shridhar S Bagali, Optimization Of Lactic Acid Production From Cane Sugar Molasses, Research Journal Of Pharmaceutical, Biological And Chemical Sciences, 2013, 4(1), 872-876.
 17. B. Goranov, D. Blazheva, G. Kostov, Z. Denkova And Y. Germanova, Lactic Acid Fermentation With Encapsulated Lactobacillus Casei Ssp. Rhamnosus Atcc 11979 (NBIMCC 1013) In Alginate/ Chitosan Matrices, Bulgarian Journal Of Agricultural Science, 2013, 19 (2), 101-104.
 18. Richard Bayitse, Lactic Acid Production From Biomass: Prospect For Bioresidue Utilization In Ghana: Technological Review, International Journal Of Applied Science And Technology, 2015, 5(1), 164-174.
 19. Pushparani Mudaliyar And Chandrashekar Kulkarni, Screening Of Novel Substrates For Lactic Acid Production By Rhizopus Oryzae, Int. Journal Of Life Science And Pharma Research, 2012, 2(1), 122-127.
 20. Young-Jung Wee, Jin-Nam Kim And Hwa-Won Ryu, Biotechnological Production Of Lactic Acid And Its Recent Applications, Food Technol. Biotechnol., 2006, 44(2), 163-172.
 21. Umar Farooq, Faqir Muhammad Anjum, Tahir Zahoor, Sajjad-Ur-Rahman, Muhammd Atif Randhawa, Anwaar Ahmed And Kashif Akram, Optimization Of Lactic Acid Production From Cheap Raw Material: Sugarcane Molasses, Pak. J. Bot., 2012, 44(1), 333-338.
 22. Nguyen Phuoc Minh, Investigation Of Lactic Acid Fermentation From Corn By-Product Using L. Casei And L. Plantarum Strain, International Journal Of Multidisciplinary Research And Development, 2014, 1(3), 92-100.
 23. M. A. Farboodniay Jahromi, M.S. Hatampour, R. Shafiei, Lactic Acid Fermentation By Different Lactobacillus Species Using Sorghum Seed Extract As Carbon Source, Iranian Journal Of Chemical Engineering, 2013, 10(2), 1-12.
 24. W. H. Munyon And D. J. Merchant, The Relation Between Glucose Utilization,

- Lactic Acid Production And Utilization And The Growth Cycle Of L Strain Fibroblasts, *Experimental Cell Research*, 1959, 17, 490-498.
25. Z Zhang, B Jin, L(+)-Lactic Acid Production Using Sugarcane Molasses And Waste Potato Starch:An Alternative Approach, *Proc Aust Soc Sugar Cane Technol.*, 2009, 31, 372-381.
 26. Zhangwei Xue, Limin Wang, Jiansong Ju, Bo Yu, Ping Xu And Yanhe Ma, Efficient Production Of Polymer-Grade L-Lactic Acid From Corn Stover Hydrolyzate By Thermophilic *Bacillus Sp.* Strain XZL4, *Springerplus*, 2012, 1(43),1-7.
 27. Grazina Juodeikiene, Elena Bartkiene, Pranas Viskelis, Dalia Urbonaviciene, Dalia Eidukonyte And Ceslovas Bobinas, Fermentation Processes Using Lactic Acid Bacteria Producing Bacteriocins For Preservation And Improving Functional Properties Of Food Products, *Advances In Applied Biotechnology*, [www.Intechopen.com](http://www.intechopen.com), 64-103,
 28. Hurok Oh, Young-Jung Wee, Jong-Sun Yun, And Hwa-Won Ryu, Lactic Acid Production Through Cell-Recycle Repeated-Batch Bioreactor, *Applied Biochemistry And Biotechnology*, 2003, vol.105-108, 603-613.
 29. Niju Narayanan, Pradip K. Roychoudhury, Aradhana Srivastava, L (+) Lactic Acid Fermentation And Its Product Polymerization, *Electronic Journal Of Biotechnology*, 2004, 7(2), 167-179.
 30. Yoganand Sobrun, Archana Bhaw-Luximon, Dhanjay Jhurry, Daneshwar Puchooa, Isolation Of Lactic Acid Bacteria From Sugar Cane Juice And Production Of Lactic Acid From Selected Improved Strains, *Advances In Bioscience And Biotechnology*, 2012, 3, 398-407.
 31. Vethakanraj Helen Shiphrah, Sayanti Sahu, Ashoke Ranjan Thakur And Shaon Ray Chaudhuri, Screening Of Bacteria For Lactic Acid Production From Whey Water, *American Journal Of Biochemistry And Biotechnology*, 2013, 9 (2),118-123.
 32. Limin Wang, Bo Zhao, Bo Liu, Bo Yu, Cuiqing Ma, Fei Su, Dongliang Hua, Qinggang Li, Yanhe Ma, Ping Xu, Efficient Production Of L-Lactic Acid From Corn cob Molasses, A Waste By-Product In Xylitol Production, By A Newly Isolated Xylose Utilizing *Bacillus Sp.* Strain, *Bioresource Technology*, 2010, 101(20), 1-8.
 33. Merih Kivanc, Meral Yilmaz, Erdoğan Cakir, Isolation and identification of Lactic acid bacteria from boza, and their microbial activity against several reporter strains, *Turk J Biol.*, 2011, 35, 313-324.
 34. Miloud Hadadji and Ahmed Bensoltane, Growth and Lactic acid production by *Bifidobacterium longum* and *Lactobacillus acidophilus* in goat's milk, *African Journal of Biotechnology*, 2006, 5(6), 505-509.
 35. Ronald H., W. Maas, Robert R. Bakker, Gerrit Eggink, Ruud A. Weusthuis, Lactic Acid Production From Xylose By The Fungus *Rhizopus Oryzae*, *Appl Microbiol. Biotechnol.*, 2006, 72, 861-868.
 36. Ayo S. Afolabi, Production of Lactic Acid from Cassava Starch Hydrolysate using Immobilized *Lactobacillus Casei* in a Fibrous Bed Bioreactor, *Proceedings of the World Congress on Engineering 2012 Vol III, WCE 2012*, July 4 - 6, 2012, London, U.K., 1-4.
 37. Ramesh Chandra Ray, Piyush Sharma and Smita Hasini Panda, Lactic acid production from cassava fibrous residue using *Lactobacillus plantarum* MTCC 1407, *Journal of Environmental Biology*, 2009, 30(5), 847-852.

How to cite this article: Kulkarni SJ. Production of lactic acid from various feed stocks: a review on studies and research. *Int J Res Rev.* 2015; 2(12):758-763.
