The role of soil's particle-size fractions in the adsorption of heavy metals

Saglara Mandzhieva [a,*], Tatiana Minkina [a], David Pinskiy [b],
Tatiana Bauer [a], Svetlana Sushkova [a]

[a] Southern Federal University, Rostov-on-Don, Russia
[b] Institute of Physicochemistry and Biological Problems of Soil Sciences, Moscow oblast, Pushino, Russia

Abstract

The parameters of adsorption of Cu^{2+}, Pb^{2+}, and Zn^{2+} cations by southern chernozem and their particle-size fractions were studied. The adsorption of metals by soils and the strength of the fixation on the surface of soil particles under both mono- and poly-element contamination decreased with the decreasing proportion of fine fractions in the soil. The aim of this work was to study the effect of the particle-size distribution and the silt and physical clay fractions on the adsorption of copper, lead, and zinc by chernozems. The objects of study included the upp humus horizons of different southern chernozems of the Rostov oblast. To study the ion-exchange adsorption of the Cu^{2+}, Pb^{2+}, and Zn^{2+} cations, the soil in the natural ionic form was

Article Info

Received : 15.05.2014
Accepted : 14.10.2014

Table 2. Parameters of the Cu^{2+}, Pb^{2+}, and Zn^{2+} adsorption by southern chernozems with different particle-size distributions at the mono- (above the line) and polyelement (under the line) contamination of soils

Adsorption parameters	Cu^{2+}	Pb^{2+}	Zn^{2+}
Clay loamy southern chernozem			
Cmax, mM/kg	tuáxx¬uáwv suáw,{19s	tsásw¬wá{r stázz¬sáxt	suávz¬rávz váuu¬ráut
k, l/mM	wzátw¬suátr sswá{r¬uwáys	vyásu¬syáss wrárz¬záyu	vár{¬ráuw ssáuv¬táyy
R^2	0,99 0,93	0,96 0,99	0,99 0,97
Loamy southern chernozem			
Cmax, mM/kg	tráw{¬{ásx suárw¬sá{y	sxáwy¬wáu{ stázt¬sátt	stáww¬ráw{ 4,1{¬rátx
k, l/mM	wváwv¬tsá{t zváxw¬syárr	uvázr¬stáyu vrász¬záyr	uá{w¬rátw {á{v¬sá{{
R^2	0,98 0,95	0,88 0,98	0,99 0,98
Loamy sandy southern chernozem			
Cmax, mM/kg	s{árs¬sárt stáuw¬sáxx	sváxw¬sás{ ssáxr¬sáwv	ssárw¬rá{u uáyz¬rász
k, l/mM	twá{r¬táux vzávr-48sá	tsázu¬uávv usáus¬yá{v	táxu¬ráuv zásty¬sávu
R^2	0,99 0,95	0,99 0,97	0,99 0,99

The isotherms had a form described by the following Langmuir equation:

$$C_{ads} = C_{max} kC/(1 + kC), \quad (1)$$

where C_{ads} is the content of the adsorbed cations, C_{max} is the maximum adsorption of the HM, mM/100 g of soil; k is the constant of the affinity; and C is the HM concentration in the equilibrium solution, mM/l.

The analysis of the obtained data showed for the southern chernozems that a tendency toward a decrease in the maximum adsorption C_{max} and the metal adsorption constant k was observed when going from the clay loamy to loamy sandy soils (Figure 1). The tendency was more pronounced for the adsorption of Cu^{2+} and Pb^{2+} ions. The decrease in C_{max} at the change of the particle-size distribution in similar soils is related to the known relationship between the specific surface and the adsorption capacity of the soils. For the southern chernozems, it varied in the following sequence: clay loamy > loamy > loamy sandy.

The tendencies toward regular changes in the parameters of the ion exchange adsorption of the HM cations by the soils with different particle-size distributions can be due to the actual differences in the chemistry and mineralogy of the corresponding fractions of the studied soils and the ability of HMs to specifically interact with specific groups of exchangeable sites (Pinsky et al., 2013). Wong et al. (2007) studied the sorption of zinc by soils differing in acidity and particle-size. By the ability to sorb zinc cations, they formed the following sequence: calcareous clayey soil > calcareous sandy soil > acid sandy laterite soil. This sequence indicated an important role of the particle-•‹œ‡ †‹•–”‹„——–‹‘• ƒ•† –Š‡ •‘‹Ž •‘Ž——–‹‘•ï• ’ metals.

The important role of organic matter in metal adsorption was noted in the works of Karpukhin and Sychev (2003) and Minkina et al. (2006). Data obtained by Plyaskina and Ladonin (2005) indicated that more than 50% of the copper and zinc in all the particle-size fractions of a leached chernozem was bound to organic matter, and the remaining portion was strongly bound to the mineral soil components, including iron minerals. For zinc, the interaction with organic matter was less typical (Putilina et al., 2009; Samokhin et al., 2003). At the same time, a significant part of the surface of chernozem particles is covered with humus films. They have a complex effect on the adsorption capacity of the soil particles. They can hamper the contact of HM ions with reactive sites located on the surfaces of mineral soil components and in their medium and small pores and hydrophobize some surface areas, which resulted in a decrease in the exchange capacity of soils (Pinskii and Kurochkina, 2006). At the same time, new reactive sites can appear due to the functional groups of adsorbed organic molecules. However, the ratio between these two processes in real soils is almost not studied, and no definite quantitative conclusions can be drawn about their effect on the ion exchange adsorption capacity of soils.

The reliable decrease in the value of k with the decreasing content of the fine fractions in soils cannot be explained only by the change in the specific surface of the soil particles. The constants characterize the energy of the cations interaction with the active sites on the surfaces and are mainly related to the qualitative composition of the particles. Thus, the change in the adsorption constants with the variation in the particle-size distribution clearly indicates a difference in the qualitative composition of the fine fractions of the soils studied. Moreover, the k values depend more on the content of the fine fractions than the C_{max} values. The polyelement adsorption of HMs by soils is characterized by their mutual competition for the exchangeable sites. This competition changes not only the shape of the isotherms but also the adsorption parameters (Figure 1 and Table 2).

In general, the isotherms of the polyelement adsorption of HMs by the studied soils follow the Langmuir equation. The mutual competition of ions eliminates the differences manifested at the monoelement adsorption. According to the constants of the adsorption on the clay loamy and loamy southern chernozems, the cations of the studied HMs form the following sequence: $Cu^{2+} > Pb^{2+} >> Zn^{2+}$. On loamy sandy southern chernozem, they form the following sequence: Cu^{2+} [illegible]$^{2+}$ > Zn^{2+}. Hence, the relative affinities of all the soils for Cu^{2+} and Pb^{2+} ions are again significantly higher than for Zn^{2+}, as is confirmed by the isotherm shapes.

When the studied soils are ranked according to the HM adsorption parameters, the following sequence is observed for the adsorption constants: clay loamy southern chernozem > loamy southern chernozem > loamy sandy southern chernozem. The values of C_{max} for the adsorption of copper and lead little vary with the changes in the particle-size distribution. However, the tendency toward a decrease in this parameter with decreasing contents of physical clay and silt is clearly traced. Thus, in this case too, the extensive adsorption parameter - the maximum adsorption - is less sensitive to changes in the particle-size distribution than the intensive parameter of the process - the constant of the adsorption equilibrium.

The simultaneous participation of three HM cations in the process resulted in their redistribution among the sorption sites with the highest relative affinities for these cations because of the cations mutual competition (Ladonin and Plyaskina, 2003). Correspondingly, the individual values of the maximum adsorption for each of them (Cu^{2+}, Pb^{2+}, and Zn^{2+}) decreased, and the constants characterizing the energy of their interaction with the sorption sites on the surface of the soil particles increased (Table 2). These changes were most noticeable and statistically reliable for zinc, because it was the least competitive element compared to copper and lead (Minkina et al., 2014; Pinskii et al., 2013).

The physical clay, which includes the silt and fine dust ([illegible] fractions), is the main carrier of [illegible] adsorption properties; therefore, given the quantitative and qualitative compositions of the soil, the further behavior of the HMs can be predicted. Therefore, the direct study of the HM adsorption by the separate soil particle-size fractions is of significant interest.

The separation of the particle-size fractions by the pipette method after the pyrophosphate treatment of the samples disturbs the natural composition of the exchangeable cations; therefore, the samples should be converted into the same state. For this purpose, the soil and its particle-size fractions were converted into the monoionic Ca form. The original soil sieved through a 1 mm sieve and the separated particle-size fractions were converted into the monoionic Ca forms by tenfold treatment with a 0.25 M $Ca(NO_3)_2$ solution. Then, the soil was washed from excess salt with distilled water, air dried, and homogenized. Samples of the soils and fractions were treated with tenfold volumes of solutions containing different amounts of HM nitrates with the addition of calcium nitrate to maintain a constant ionic strength of 0.01 mol/l. The further procedures were similar to those described above for the study of the HM adsorption by soils.

The general shapes of the isotherms of the Cu^{2+}, Pb^{2+}, and Zn^{2+} adsorption by the silt and physical clay fractions from the southern chernozem were analogous to those of the HM adsorption by the entire soil: the adsorption was of limited character and followed the Langmuir equation (Figure 2). Copper and lead were more intensively adsorbed by the particle-size fractions than the zinc.



Figure 2. Isotherms of adsorption of (1) Cu^{2+}, (2) Pb^{2+}, and (3) Zn^{2+} by the (a) silt and (b) physical clay fractions of the clay loamy southern chernozem under monoelement contamination.

In the particle-size fractions separated from the soils, the concentrations of copper, lead, and zinc decreased with the decreasing particle-size. The values of k and C_{max} characterizing the adsorption of HMs by the southern chernozem and its particle-size fractions formed the following sequence: silt > physical clay > entire soil (Table 3).

Table 3. Parameters of the Cu^{2+}, Pb^{2+}, and Zn^{2+} adsorption by a clay loamy southern chernozem and its particle-size fractions

Adsorption parameters	Cu^{2+}	Pb^{2+}	Zn^{2+}
Silt			
Cmax, mM/kg	tzävw ¬ rävx	twätr ¬ räw{	syä{r ¬ rävy
k, l/mM	zrätr ¬ trät{	xwä{r ¬ sxäsv	szäxw ¬ uärr
R^2	0.99	0.99	0.99
Physical clay			
Cmax, mM/kg	ttäsw ¬ sätt	trävr ¬ täsw	stäwr ¬ sä{x
k, l/mM	wzätr ¬ sväwv	v{ätx ¬ suäuw	stäry ¬ uäst
R^2	0.92	¬ rä{z	¬ rä{z
Soil			
Cmax, mM/kg	syäwz ¬ uäru	sväwv ¬ tä{y	zä{{ ¬ sä{r
k, l/mM	uzäzr ¬ stäuu	urävw ¬ ssä{x	vä{v ¬ räxu
R^2	0.95	¬ rä{x ¬	0.94

The works of some authors (Baidina, 2001; Derkham, 2009; Zhuravleva, 1985; Zyrin et al., 1986; Zyrin and Chebotareva, 1989; Panin and Siromlya, 2005; Plyaskina and Ladonin, 2005; Sadovnikova and Ladonin, 2000; Titova et al., 1996; Travnikova et al., 2000) carried out with different soil types showed that the saturation of the fine HM fractions decreased in the following sequence: silt > fine dust > medium dust. The dust fraction contained 14.6% Cu^{2+}, 16.6% Pb^{2+}, and 12.6% Zn^{2+} on the average for all the soils (Zyrin and Chebotareva, 1989). On the average for all the soils but the krasnozem, the concentrations of Cu^{2+}, Pb^{2+}, and Zn^{2+} in the silt fractions were found to be 52, 38, and 162 mg/kg, respectively. Higher contents of HMs in these fractions were reported by Titova et al. (1996). It was noted that the dust and silt fractions of the soils accumulated 25 65% of the Cu^{2+}, 35 95% of the Pb^{2+}, and 53 89% of the Zn^{2+}. The maximum concentration of Cu^{2+} was found in the dust fractions lower than 1.8 g/cm^3 in density; that of Pb^{2+} was most frequently observed in the silt.

The differences in the parameters of the Cu^{2+}, Pb^{2+}, and Zn^{2+} adsorption by the studied soils and their particle-size fractions are determined not only by the specific surfaces of the adsorbents but also by their composition and properties. The silt fraction mainly consists of secondary clay minerals, coagels of iron hydroxides and silicic acid, crystallized iron hydroxides, amorphous compounds of different compositions, and humic substances of lower degrees of condensation. These compounds usually have large exchange capacity and high affinity for HMs. The fine dust fraction, which also enters in the physical clay fraction, contains more primary minerals (quartz and amphiboles), as well as mica and iron hydroxides.

The data obtained by Kryshchenko and Kuznetsov (2003) showed that the silt fractions of the southern chernozem contain four groups of clay minerals: kaolinite, chlorite, mica, and smectite. In the dust fraction of the soils, the content of hydromicas increases by 10-19% compared to the silt faction, and that of smectites decreases. The content of humus in the dust fraction increases by the same factor. The content of clay minerals in the silt and dust fractions is higher than that in the entire soil. On the other hand, the higher the content of secondary minerals in the fraction, the higher its adsorption capacity.

The role of organic matter in the adsorption of HMs by the soils and the particle-size fractions is ambiguous. The soil organic matter has a high exchange capacity. However, in the soils and especially in the fine fractions, it is usually bound to the mineral soil components. Highly condensed humus blocks the medium and fine pores, which significantly reduces the adsorption capacity. Organic molecules are strongly sorbed at these positions because of their high activity (Kaiser and Guggenberger, 2003). Kurochkina and Pinskii (2002) showed that carboxyls containing organic molecules are most strongly sorbed on positively charged surface ites, primarily on the apices and edges of crystals and various surface defects, including medium and fine pore throats.

To assess the role of humus in the adsorption of HM cations by the soils and their particle-size fractions, the coefficients of correlation (R) between the content of the sorbed Cu^{2+}, Pb^{2+}, and Zn^{2+}; the contents of the silt and physical clay fractions in the soils; and the humus under mono- and polyelement contamination were calculated. The closest correlations were found between the content of the adsorbed HMs and the contents of the silt and physical clay in the soils under the monoelement contamination. A looser correlation was observed between the content of the adsorbed HMs and the content of humus in the studied soils.

Conclusion

- The increase in the degree of dispersion of the particle-size fractions in similar soils resulted not only in an increase in the content of adsorbed HMs but also in an enhancement of their fixation on the surface of the fine particles. Therefore, the adsorption capacity of the Lower Don southern chernozems for Cu^{2+}, Pb^{2+}, and Zn^{2+} decreased in the following sequence: clay loamy southern chernozem > loamy southern chernozem > loamy sandy southern chernozem.
- The isotherms of the Cu^{2+}, Pb^{2+}, and Zn^{2+} adsorption were described by the Langmuir equation in all the cases. In the case of the polylement adsorption of HMs on the soils and their particle-size fractions, the •—––—ƒŽ …‘•’‡–‹–‹‘• ‘ˆ –Š‡ …ƒ–‹‘•• ˆ‘” –Š‡ ‡š…Šƒ•‰‡ƒ„Ž‡ •‹–‡• ƒ adsorption parameters. The Cmax values of the individual ions decreased and the k values increased compared to the corresponding data for the monoelement adsorption. This indicated an increase in the adsorption selectivity (specificity) of each competing cation. The changes in the adsorption parameters were the most pronounced for zinc and less pronounced (manifested as tendencies) for lead.
- According to the values of the adsorption parameters (C_{max} and k) for the different particle-size fractions, the HM cations formed a sequence analogous to that obtained for the entire soils: Cu^{2+} · „$^{2+}$ > Zn^{2+}.
- The parameters of the HM adsorption by the same particle-size fractions isolated from the different soils decreased in the following sequence: clay loamy chernozem > loamy chernozem > loamy sandy chernozem. This was related to the qualitative differences in the mineralogy and chemistry of the separated fractions and the significant effect of their composition and properties on the parameters of the HM adsorption.
- The analysis of the changes in the parameters of the Cu^{2+}, Pb^{2+}, and Zn^{2+} adsorption by the studied soils and their particle-size fractions showed that the extensive characteristic - the maximum adsorption (C_{max}) - is a less sensitive parameter characterizing the adsorption capacity of the soil than the intensive characteristic of the process - the adsorption equilibrium constant (k).

Acknowledgements

This research was supported by projects of the Ministry of Education and Science of Russia, no. wäzzwätrsv ™á –Š‡ —••‹ƒ• ‘–•‡†ƒ–‹‘• ‘ˆ ƒ•‹… esearch, no. 14-05-00586-a, grant of the President of —••‹ƒ• ‡†‡”ƒ–‹‘• -6448.2014.4, Russian Foundation of Basic Research no. 14-04-00034, and the Leading Scientific School, no. 5548.2014.5. Analytical work was carried out on the equipment of Centers for collective use of Southern Federal University "High Technology", "Biotechnology, Biomedical and Environmental Monitoring".

References

Agrochemical Methods of Soil Studies, 1975. Akad. Nauk SSSR, Moscow [in Russian].

Aslanov, N.N., Ryzhov, S.N., 1969. Composition and Properties of Particle-Size Fractions of Sierozems. Fan, Tashkent [in Russian].

Baidina, N.L., 2001. On the Contents of Heavy Metals in Particle-Size Fractions of a Soil in Novosibirsk. Agrokhimiya. 3, 69 74 [in Russian]

Vadyunina, A.F., Korchagina, Z.A., 1986. Methods for Studying Soil Physical Properties. Agropromizdat, Moscow [in Russian].

‡”•Š ƒ•á Šä äá trr{ä š–‡•†‡† „•–”ƒ…– ‘ˆ ƒ•†‹†ƒ–‡ï• ‹••‡”–ƒ–‹‘• ‹• ‹‘Ž‘‰›ä

Zhuravleva, E.G., 1985. Microelements in Soils and Modern Methods for Their Study. Moscow [in Russian].

Zyrin, N.G., Serdyukova, A.V., Sokolova, T.A., 1986. Sorption of Lead and the State of Adsorbed Lead in Soils and Soil Components. Pochvovedenie. 4, 39 44 [in Russian]

Zyrin, N.G., Chebotareva, N.A., 1979. The Contents and Forms of Microelement Compounds in Soils. Moscow [in Russian].

Karpukhin, A.I., Sychev, V.G., 2005. Complex Compounds of Soil Organic Substances and Metal Ions. Moscow [in Russian].

Kryshchenko, V.S., Golozubov, O.M., Kolesov, V.V., Ryabyets, T.V., 2008. Data Bases on the Composition and Properties of Soils. Rostov-on-Don [in Russian].

Kryshchenko, V.S., Kuznetsov, R.V., 2003. Clay Minerals in Soils of the Lower Don and Northern Caucasus. Izv. Vuzov. Severo-Kavkazskii Region. Estestv. Nauki. 3, 86 92 [in Russian]

Kurochkina, G.N., Pinskii, D.L., 2002. Mechanism of Adsorption of High-Molecular Surfactants on Synthetic Analogues of Soil Aluminosilicates. Eurasian Soil Science 35 (10), 1046 1057.

Ladonin, D.V., 2003. The Effect of Iron and Clay Minerals on the Adsorption of Copper, Zinc, Lead, and Cadmium in the Nodular Horizon of Podzolic Soil. Eurasian Soil Science 36 (10), 1065 1073.

Ladonina, N.N., Ladonin, D.V., 2000. Soil Contamination with Copper and Zinc in the Southeastern Administrative District of Moscow. Ekologiya. 1, 61 64 [in Russian]

Ladonin, D.V., Plyaskina, O.V., 2004. Mechanisms of Cu (II), Zn (II), and Pb (II) Sorption by Soddy-Podzolic Soil. Eurasian Soil Science 37 (5), 460 468.

Ladonin, D.V., Plyaskina, O.V., 2003. Fractional Composition of Copper, Zinc, Cadmium, and Lead Compounds in Some Soil Types under Polyelemental Contamination. Moscow University Soil Science Bulletin 1, 8 15.

Minkina, T.M., Motuzova, G.V., Nazarenko, O.G., 2006. Interaction of Heavy Metals with Organic Matter of an Ordinary Chernozem. Eurasian Soil Science 39 (7), 720 726.

Minkina, T.M., Pinskiy, D.L., Bauer, T.V., Mandzhieva, S.S., Belyaeva, O.N., Kalinichenko, V.P., Endovitsky, A.P., 2014. Effect of associated anions on zinc adsorption and transformation in chernozems. Journal of Geochemical Exploration 144, 226 229

Samokhin, A.P., Minkina, T.M., Statovoi, A.A., 2003. Comparison anion effect on zink and lead adsorption by chernozem and its pH value. Proceeding of 7th International Conference on the Biochemistry of Trace Elements. Uppsala. V.4, pp. 82-83.

Panin, M.S., Siromlya, T.I., 2005. Adsorption of Copper by Soils of the Irtysh River Region, Semipalatinsk Oblast. Eurasian Soil Science 38 (4), 364 378.

Perelomov, L.V., 2001. Extended Abs–”ƒ…– ‘ˆ ƒ•†‹†ƒ–‡ï• ‹••‡”–ƒ–‹‘• ‹• ‹‘Ž‘‰›ä ‘•…‘™ ‹•

Perelomov, L.V., Pinskii, D.L., 2003. Mn, Pb, and Zn Compounds in Gray Forest Soils of the Central Russian Upland. Eurasian Soil Science 36 (6), 610 618.

Pinskii, D.L., Kurochkina, G.N., 2006. Evolution of the Concepts of the Physicochemical Adsorption in Soils. Nauka, Moscow [in Russian].

Pinsky D.L., Minkina T.M. 2013. Regularities of Cu, Pb and Zn adsorption by chernozems of the South of Russia. Eurasian Journal of Soil Science 2, 59 68.

Pinskii D. L., Minkina T. M., Mandzhieva S. S., Fedorov U.A., Nevidomskaya D.G, Bauer T. V. 2014. Adsorption features of Cu(II), Pb(II), and Zn(II) by an Ordinary Chernozem from Nitrate, Chloride, Acetate, and Sulfate Solutions. Eurasian Soil Science 47 (1): 10 17

Plyaskina, O.V., Ladonin, D.V., 2005. Heavy Metal Compounds in Particle-Size Fractions of Some Soil Types. Vestn. Mosk. Univ., Ser. 17: Pochvovedenie 4, 36-43.

Putilina, V.S., Galitskaya, I.V., Yuganova, T.I., 2009. Adsorption of Heavy Metals by Soils and Rocks. Characteristics of Sorbents and Conditions, Parameters, and Mechanisms of Adsorption: Analytic Review. GPNTB SO RAN, Novosibirsk [in Russian].

Sadovnikova, L.K., Ladonin, D.V., 2000. Sorption of Copper and Zinc by Soddy-Podzolic Soils under Different Levels of Technogenic Pollution. Vestn. Mosk. Univ., Ser. 17: Pochvoved. 3, 33-39

Titova, N.A., Travnikova, L.S., Kakhnovich, Z.N., Sorokin, S.Ye., Schulz, E., Korschens, M., 1996. Heavy Metal Content in Various Particle-Size and Density Fractions of Soils. Eurasian Soil Science 29 (7), 820 830.

” ƒ ˜ • ‹ • ‘ ˜ ƒ á ä ä á ƒ • Š • ‘ ˜ ‹ … Š á ä ä á ‘ Ž ï • Š ƒ • ‘ ˜ á ä ä á ‘ ‰ — – á ä ä á ‘ ” ‘ • ‹ • á Significance of Organomineral Fractions for Evaluation of Heavy Metal Contamination of Soddy-Podzolic Soil. Eurasian Soil Science 33 (1), 81 88.

Shaimukhametov, M.Sh., 1993. On the Method of Determination of Adsorbed Ca and Mg in Chernozemic Soils. Pochvovedenie. 12, 105 111.

McBride, M.B., 1989. Reactions Controlling Heavy Metals Solubility in Soils. Advances in Soil Science 10, 2 47.

Kaiser. K., Guggenberger, G., 2003. Mineral Surface and Soil Organic Matter. Eurasian Soil Science 54, 219 236.

Wong, J.W.C., Li, K.L., Zhou, L.X., Selvam, A., 2007. Sorption of Cd and Zn by Different Soils in the Presence of Dissolved Organic Matter from Sludge. Geoderma 137, 310 317.