

SPECTRUM OF FNAC IN PALPABLE HEAD AND NECK LESIONS IN A TERTIARY CARE HOSPITAL IN INDIA- A 3 YEARS STUDY

Suryawanshi Kishor H.^{1,*}, Damle Rajshri P.², Dravid Nandkumar V.³, Tayde Yogesh⁴

¹Associate Professor, ^{2,4}Assistant Professor, ³Professor & Head, Department Of Pathology
A. C.P.M. Medical College, Dhule, Maharashtra. India

***Corresponding Author:**

Email: ompathologylab@gmail.com

ABSTRACT

Background: Head and neck lesions encompass a multitude of congenital, inflammatory or neoplastic lesions including several anatomic sites and originating in different tissues and organs. Fine needle aspiration cytology (FNAC) is a simple, quick, feasible, cost effective and repeatable outpatient procedure with minimal risk of complication.

Aim:

1. To evaluate the role of FNAC and its utility in diagnosis of palpable head neck masses.
2. To study the spectrum of head neck lesions in rural population.
3. To study diagnostic accuracy of FNAC by histopathological correlation wherever possible.

Material and methods: The study included 288 patients presented with palpable head and neck swelling in a tertiary care hospital from January 2011 to December 2014. Detailed clinical history of patient was noted. Aspirations were done by using 10 ml syringe and 22/23 gauge needles. Smears were stained with PAP, Haematoxylin and eosin and Leishman stain. Cytomorphological diagnosis was given. Cyto-histopathological correlations were done wherever possible.

Results: Out of 288 patients of head and neck lesions studied, lymph node (39.58%) was the predominant site aspirated with tubercular lymphadenitis being the commonest lesion. Thyroid lesions constituted 31.25 % followed by salivary gland (18.75%), soft tissue and miscellaneous (7.29%). FNAC was inconclusive in 3.12 % cases. Overall accuracy rate of FNAC was 93.02% with sensitivity of 81.81 % and specificity of 96.87 %.

Conclusion: Though excisional biopsy is the gold standard for diagnosis of head and neck neoplastic lesion. FNAC is a rapid, cheap diagnostic tool now-a-days with overall accuracy rate more than 90 %.

Key words: head and neck, FNAC, rapid, diagnostic

INTRODUCTION

Palpable head and neck swellings include various non-neoplastic and neoplastic lesions of lymph node, salivary gland, thyroid gland and soft tissue. Proximity of tissues of various types and wide range of primary and metastatic neoplasms are responsible for this site being the most common in FNAC diagnosis.^[1]

FNAC is relatively painless, produces speedy results and cheap. FNA is highly suitable for debilitated patients, is repeatable, useful for multiple lesions and has low risk of complications. FNAC is applicable to easily palpable lesions of thyroid, breast, salivary glands, superficial lymph nodes, superficial growth of skin & soft tissue. With the help of newer

radiological techniques FNAC of deeper structures is easily possible.^[1] Head and neck neoplasms is a major form of cancer in India accounting for 23 % of all cancer in males and 6 % in females.^[2,3] FNAC of head and neck region is a generally well accepted technique with high specificity.^[4] Psychological and economical advantage of an immediate diagnosis in outpatient clinic is obvious. Ancillary techniques done on cytology like flow cytometry, cytogenetic, electron microscopy, cell block preparation, immunocytochemistry have further added a tool in diagnosis.

FNAC is a prerequisite for various neck swellings as procedure is non-traumatic, easily accessible, inexpensive, excellent compliance, avoids anaesthetic complications and requirement of open

surgical biopsy. FNAC differentiates non neoplastic lesions from neoplastic lesions thus eliminating need of surgical intervention in these lesions which can be treated conservatively.^[5]

MATERIAL AND METHODS

The present study was conducted in Department of Pathology from January 2011 to December 2014 and included 288 patients with palpable head and neck swellings. Outdoor as well as indoor patients with palpable head and neck swellings were referred to cytology department. Detail clinical history and significant findings were noted. After explanation of procedure and taking informed consent of patient, FNAC was done using 10 cc disposable syringe and 22/23 gauge needle taking all aseptic precautions. Both aspiration and non-aspiration technique was used wherever required. Three or four smears were prepared by cytopathologist following standard guidelines. Wet fixed smears in 95% alcohol were stained with PAP and Haematoxylin-Eosin stain while air dried smears were stained with Leishman stain. Zeihl-Neelsen staining for acid fast bacilli was done in suspected tubercular lesions. Aspirations taken from various sites include lymph node, thyroid, salivary gland and soft tissue. Cytomorphological diagnosis was given depending upon the pathology. Excisional biopsy specimens were fixed in 10% neutral buffered formalin processed by paraffin embedding and stained with haematoxylin and eosin stain. Cyto-histopathological correlation was done in those cases.

RESULTS

The present study included 288 cases palpable head and neck lesions from various departments as an OPD as well as indoor patient. Age group of patients ranged from 1 year to 75 years. Out of total 2067 cytology patients over a period of 3 years FNAC of head and neck lesions constituted 288(13.93%) of cases. Maximum no. of patients were in the age group of 21-30 years(32.63%) followed by 31-40 years(20.83%) and least no. of patients were seen in age group of above 70 years. Out of

288 patients 168(58.33%) were females and 120(41.67%) were males. Site wise distribution of head and neck FNAC [Table-1] shows lymph nodes lesion as the predominant site of FNAC followed by thyroid lesions (31.25%), salivary glands (18.75%) and soft tissue (7.29%). FNAC was inconclusive in 9 (3.12%) cases. Out of 114(39.58%) cases of lymph node lesions, tubercular lymphadenitis (47.36%) was the predominant cause of lymphadenopathy followed by reactive lymphadenitis in 40(35.08%) cases. In malignant lesions, metastatic epithelial malignancy was the predominant finding (3.50%) and one case (0.87%) of lymphoma was found.[Table-2] Amongst 90 (31.25%) cases of thyroid lesions, inflammatory lesions including Hashimoto's thyroiditis, chronic lymphocytic thyroiditis were found in 28.88 % of cases. Benign neoplastic lesions constituted 69.97% of cases. In malignant lesions one case (1.11%) of papillary carcinoma was observed.[Table-3] In salivary glands lesions, sialadenitis was observed in 66.66% of cases. Benign neoplasm included 13(24.07%) cases of pleomorphic adenoma and two cases (3.70%) of benign lymphoepithelial cyst.3 cases of malignant neoplasms were reported including one case of each of carcinoma ex pleomorphic adenoma, mucoepidermoid carcinoma and anaplastic carcinoma.[Table-4] FNAC of soft tissue and miscellaneous constituted 21 cases (7.29%) with varied pathological lesions like lipoma (23.80%), epidermal cyst (57.14%), benign adnexal tumor (4.67%) and malignant neoplasms including squamous cell carcinoma and metastatic epithelial neoplasm.[Table-5] Histo-cytopathological correlation was possible in 43 cases only which included 33 benign lesions and 10 malignant lesions. In 31 cases of benign lesions cytological diagnosis was consistent with histopathological diagnosis while two cases showed false negative result. In 9 cases of malignant lesions cytological diagnosis was consistent with histopathological examination and one case showed false positive result. Overall accuracy rate was 93.02% with sensitivity, specificity, positive predictive value and negative predictive value of 81.81%, 96.87%, 90.0 % and 93.93% respectively.

Table-1: Distribution of Head & Neck Lesions [Cases=288]

Site	No. of Cases	%
Lymph Node	114	39.58
Thyroid gland	90	31.25
Salivary gland	54	18.75
Soft tissue & miscellaneous	21	7.29

Table-2: Distribution of Various Lymph Node Lesions [N=114]

Lesions	No. of cases	%	
Reactive lymphadenitis	40	35.08	
Inflammatory	Nonspecific	14	12.80
	Tuberculosis	54	47.36
Malignant	Lymphoma	1	0.87
	Metastasis	4	3.50
Total	114	100	

Table-3: Distribution of various Thyroid lesions [n= 90]

Thyroid lesions	No. of cases	%
Inflammatory	26	28.88
Benign		
1.Colloid goitre	43	47.77
2.Hurthle cell neoplasm	8	8.88
3.Thyroglossal cyst	6	6.66
4.Follicular neoplasm	6	6.66
Malignant		
1.Papillary carcinoma	1	1.11
Total	90	100

Table-4: Distribution of various Salivary gland lesions [n= 54]

Salivary gland lesions	No. of cases	%
Inflammatory		
1.Acute sialadenitis	12	22.22
2. Chronic sialadenitis	24	44.44
Benign		
1.Pleomorphic adenoma	13	24.07
2.Benign lymphoepithelial lesion	2	3.70
Malignant		
1.Ca ex pleomorphic adenoma	1	1.85
2.Mucoepidermoid carcinoma	1	1.85
3.Anaplastic carcinoma	1	1.85
Total	54	100

Table-5: Distribution of soft tissue & miscellaneous [n=21]

	No. of cases	%
Benign		
1.Lipoma	5	23.80
2.Epidermal cyst	12	57.14
3.Benign adnexal tumor	1	4.67
Malignant		
1.Squamous cell carcinoma	2	9.52
2.Metastatic epithelial tumor	1	4.67
Total	21	100

DISCUSSION

In 1930, Martin and Ellis described and first introduced the technique of FNAC for diagnosis of organ lesion.^[6] The two fundamental requirements on which success of FNA depends are representative sample and high quality of preparation. These two prerequisites will always remain a sine qua non, no matter how sophisticated supplementary techniques.^[1] Head and neck neoplasm constitute a major form of cancer in India accounting for 23% of all cancer in males and 6% in females^[2,3] and approximately 5% all childhood neoplasms.^[7] Increased prevalence of malignancies may be due to use of various forms tobacco in our country. Palpable lesions of head and neck include variety of

developmental, inflammatory and neoplastic lesions.

The present study was carried out over a period of 3 years in a tertiary care hospital to find out frequency of variety of pathologies and to find out accuracy of FNAC as a rapid diagnostic tool in outdoor patients. The present study also compares its findings with various national and international studies published in the literature.[Table-6] The study included patients from all age groups. Majority of patients were females with male to female ratio of 0.71:1. Similar results of female preponderance were also reported by Muddegowda et al^[8], Sharma et al^[11] and Ahmad T et al.^[13]

Table-6: Comparison of Results of Various National and International Studies

	Our study 2014	Muddegowda et al ^[8] [2014]		Bhagat VM et al ^[9] [2013]	Mohmed MH ^[10] [2013]	Sharma R ^[11] [2012]	Rathod G ^[12] [2012]	Ahmad T ^[13] [2008]	El Hag ^[4] [2003]	Kamal F ^[14] [1996]
Location	India	India Wyanand	India Salem	India	Malasiya	India	India	Peshawar	Saudi Arabia	Lahore
Duration of study	3 years	8 months	8 months	1 year	1 year 5 months	---	1 year 4 months	1 year	5 years	1 year
No.Of Patients	288	100	100	701	37	125	200	50	225	847
M:F ratio	0.71:1	0.29:1	0.53:1	-----	---	0.64:1	1.43:1	0.47:1	---	-----
Predominant site	Lymph Node	Thyroid	Thyroid	Lymph Node	Lymph Node	Lymph Node	Thyroid	Lymph Node	Lymph Node	Lymph Node
Reactive/nonspecific LN (%)	18.75	17	13	11.98	8	25.6	9.5	18	33	--
TB LN (%)	18.75	04	12	35.66	4	20.8	12	36	21	13
Malignant Neoplasm (%)	4.16	07	16	20.68	5	16	15	14	13	11
yst/goiter (%)	20.83	60	46	---	9	11.2	35	10	11	3.6
Benign neoplasm (%)	12.84	8	9	6.56	6	12	12.5	8	9	1
Sialadenitis (%)	9.02	2	1	0.71	5	6.4	4.5	6	5	0.6
Inconclusive	3.12	2	3	10	0	3.2	4.5	8	8	1.3

Predominant site of FNAC was lymph node lesions (39.58%) followed by thyroid gland. Similar result reported by various studies. [Table-6] In lymph node lesions tubercular lymphadenitis was the most common pathological findings followed by reactive lymphadenitis which is in concordance with Bhagat et al^[9], Sharma et al^[11], Ahmad T et al^[13] and El Haq et al^[4]. In malignant neoplasms, epithelial metastasis was found in 4 cases (3.50 %) and one case of lymphoma was found. All cases showed metastatic squamous cell carcinoma. This is

due to consumption of tobacco in various forms in our area leading to high incidence of malignancy in aerodigestive tract.

FNAC of thyroid lesions was the next common site in our study. Colloid goiter (47.77%) was the predominant finding in benign lesion followed by inflammatory lesions consisting of Hashimoto's thyroiditis, chronic lymphocytic thyroiditis. One case of papillary carcinoma was detected which was confirmed on histopathology. Muddegowda et al^[8] and Rathod et al^[12] also found

thyroid lesions as the predominant site of FNAC in their study with colloid goiter as the predominant finding. Female preponderance was observed in FNAC of thyroid lesion in our study with similar findings reported by Rathod et al [12] and Muddegowda et al. [8] In salivary gland lesions acute and chronic sialadenitis together comprised 66.66% followed by pleomorphic adenoma in 13 cases (24.07%) and benign lymphoepithelial cyst in two cases (3.70%). Three cases of malignant neoplasms were observed including one case of each Ca-ex-pleomorphic adenoma, mucoepidermoid carcinoma and anaplastic carcinoma. Sharma et al [11] and Rathod et al [12] found inflammatory lesions as the commonest findings followed by benign neoplasms including pleomorphic adenoma while Bhagat et al [9] found benign pleomorphic adenoma as the predominant salivary gland lesion in his study.

In soft tissue and miscellaneous lesions benign lesions were commonest finding including 12 cases (57.14%) of epidermal cyst, 5 cases (23.80%) of lipoma and one case (4.07%) of benign adnexal tumor. Three cases of malignant neoplasms comprising of 2 cases of squamous cell carcinoma and one case of metastatic epithelial tumor to scalp was reported. Bhagat et al [9] reported neoplastic lesions in 63% cases with lipoma as the predominant benign tumor and squamous cell carcinoma as the commonest malignant neoplasm. In 3.12 % of cases cytology was inconclusive. The causes of unsatisfactory aspirates were smaller lesions, poor handling of material, inadequate aspirate. Incidence of inadequate reports ranged from 0 to 10 % in various studies in the literature. Cyto-histopathological correlation was possible in only 43 cases (14.93 %). Out of 266 benign lesions in 33 cases only cyto-histopathological correlation was possible. Cytomorphological diagnosis was in concordance with histopathological diagnosis in 31 cases (93.93%) while 2 cases showed false negative results. One case reported on cytology as reactive lymphadenitis turned out to be tubercular lymphadenitis on histopathological examination. The probable reason may be focal epithelioid granuloma was missed

during aspiration hence multiple passes in various directions is necessary to avoid false negative result. Other case reported as a follicular neoplasm of thyroid gland turned out to be follicular carcinoma on histopathological examination. This is one of the known limitations of FNAC in thyroid lesions.

Out of 10 malignant cases reported on cytology, in 9 cases cytological diagnosis was consistent histopathological diagnosis. One case reported as mucoepidermoid carcinoma on cytology turned out to be pleomorphic adenoma with squamous metaplasia on histopathological examination. In this study overall accuracy rate of FNAC was 93.02% with sensitivity of 81.81%, specificity of 96.87%, and positive predictive value of 90% and negative predictive value of 93.93%. Cyto-histopathological correlation was done by Sharma et al [11] in 71 cases out of 125 cases with sensitivity of 89.28%, specificity of 90.69%, positive predictive value of 85% and negative predictive value of 90.14%. Tilak et al [15] studied cyto-histopathological correlation in 55 out of cases 154 with overall diagnostic accuracy rate of 92.75%, sensitivity of 90.91% and specificity of 93.18%. In most of the national studies including our study inflammatory and non-neoplastic lesions were the predominant cause of head and neck masses while neoplastic lesion were the commonest finding in various international studies.

To avoid or minimize false positive results various factors including regenerative changes, metaplasia and others should be taken into considerations while reporting. False negative results may be due to cystic change, necrotic and hemorrhagic areas revealing no diagnostic cellular yield. Repeat FNAC of solid areas or adjacent area may minimize false negative results in such cases. In this study we observed that FNAC is a rapid, cost effective, highly accurate and feasible first line diagnostic tool in management of palpable head and neck swellings. Though FNAC has certain limitations and pitfalls diagnostic accuracy rate in various studies including our study was over 90%.




Figure 1: FNAC of lymph node showing necrotic background and epithelioid cell granuloma [1a (Pap stain, x100) and 1b (Leishman Stain, x400)] and metastasis of squamous cell carcinoma [1c (Leishman Stain, x100) and 1d (Leishman Stain, x400)].




Figure 2: FNAC of Thyroid gland showing follicular adenoma [2a (Haematoxylin and Eosin, x100)] and intranuclear inclusions in papillary carcinoma [2b (Leishman Stain, x400)].




Figure 3: FNAC of salivary gland showing acute sialadenitis [3a (Leishman Stain, x100)], benign pleomorphic adenoma [3b(Leishman Stain,x100)] and Carcinoma Ex Pleomorphic adenoma [3c(Pap stain, x100) and 3d (Haematoxylin and Eosin, x100)].

CONCLUSION

Though excisional biopsy is the gold standard for diagnosis of head and neck neoplastic lesion FNAC is a rapid, cheap

diagnostic tool now-a-days with overall accuracy rate more than 90 % and diagnostic accuracy in differentiating non-neoplastic lesions from neoplastic lesions is well established.

REFERENCES:

- Orell SR, Sterrett GF, Walters MN-I, Whitaker D: Manual and Atlas of Fine Needle Aspiration Cytology. 2nd edn. New York: Churchill Livingstone; 1992.p. 2-36.
- Ahluwalia H, Gupta SC, Singh M, Gupta SC, Mishra V, Singh PA. Spectrum of head and neck cancers at Allahabad. J Otolaryngol Head Neck Surg 2001;53:16-20.
- Mehrotra R, Singh M, Gupta RK, Singh M, Kapoor AK. Trends of prevalence and pathological spectrum of head and neck cancers in North India. Indian J Cancer 2005;42:89-93.
- El Hag IA, Chiedozi LC, al Reyees FA, Kollur SM. Fine needle aspiration cytology of head and neck masses. Seven years' experience in a secondary care hospital. Acta Cytol 2003;47:387-92.
- Klijanienko J. Head and Neck and Salivary gland. In: Layfield LJ, editor. Atlas of Fine Needle Aspiration Cytology, 1st edn. New Delhi: Jaypee Publishers; 2014.p.11.
- Martin HE, Ellis EB. Biopsy of needle puncture and aspiration. Ann Surg 1930; 92:169-81.
- Ponder TB, Smith D, Ramzy I. Lymphadenopathy in children and adolescents: role of fine-needle aspiration in management. Cancer Detect Prev. 2000; 24:228-33.
- Muddegowda PH, Srinivasan S, Lingegowda JB, Ramkumar KR, Murthy KS. Spectrum of Cytology of Neck Lesions: Comparative Study from Two Centers. Journal of Clinical and Diagnostic Research 2014 ;Vol-8(3): 44-45 .
- Bhagat VM, Tailor HJ, Saini PK, Dudhat RB, Makawana GR, Unjiya RM. Fine Needle Aspiration Cytology In Nonthyroidal Head And Neck Masses-A Descriptive Study In Tertiary Care Hospital. National Journal Of Medical Research 2013; volume 3(3):273-76.
- Mohamed MH, Hitam S, Brito-Mutunayagam S, Yunus MRM. Role of FNAC in evaluation of neck masses. J Curr Surg 2013;3(1):19-23.