




Contents lists available at ScienceDirect

## Asian Pacific Journal of Tropical Biomedicine

journal homepage: www.elsevier.com/locate/apjtb



Document heading

doi:

© 2013 by the Asian Pacific Journal of Tropical Biomedicine. All rights reserved.

Antioxidant and cytotoxic activity of *Acanthus ilicifolius* flower

Muhamad Firdaus, Asep Awaludin Prihanto\*, Rahmi Nurdiani

Laboratory of Biochemistry, Faculty of Fisheries and Marine Science, Brawijaya University, Malang–65145, Indonesia

## PEER REVIEW

## Peer reviewer

Nashi Widodo, Associate Professor, Department of Biology, Faculty of Mathematic and Natural Sciences, Brawijaya University, Malang East Java, Indonesia.

Tel: +62 341 575841

Fax: +62 341 575841

E-mail: widodo@ub.ac.id

## Comments

The article is explained the best method to extract the active compounds from *A. ilicifolius* that have high antioxidant activity. Moreover the extracts have anti-cancer activity with low dose that warrant cancer therapeutic. This finding is really important for community, scientist and pharmaceutical company as frontier information to explore *A. ilicifolius* as source of anti-cancer material.

(Details on Page 20)

## ABSTRACT

**Objective:** To investigate the antioxidant and cytotoxic activity of the flower of *Acanthus ilicifolius* (*A. ilicifolius*). **Methods:** Antioxidant activity was determined as antiradical efficiency with diphenyl picrylhydrazil (DPPH) method and cytotoxic assay was undertaken using brine shrimp lethal toxicity test. **Results:** *A. ilicifolius* flower contained terpenoid, phenolic compounds, and alkaloid. The methanol extract of *A. ilicifolius* flower showed the highest antiradical efficiency ( $AE=1.41 \times 10^{-3}$ ) against DPPH radicals and the highest cytotoxicity ( $LC_{50}=22 \mu\text{g/mL}$ ) against brine shrimp nauplii. **Conclusions:** It is suggested that active compounds of *A. ilicifolius* flower solved in methanol play a role to inhibit free radical activity and kill *Artemia salina* nauplii. The substances can be considered as potential antioxidant and cytotoxic agents as well as imminent candidate for cancer therapy.

## KEYWORDS

*Acanthus ilicifolius*, Antioxidant, Cytotoxic, Flower

## 1. Introduction

Cancer is a notorious disease that now becomes the major cause of human mortality in the world. Almost half of the incidence and mortality happen in Asia, with lung and bronchus, breast, and colorectal cancers in women to be the most common fatal cancers[1]. The aetiology of cancer is primarily from unhealthy lifestyle and pollution. Carcinogenesis or production of cancer involves mainly three steps, namely initiation, promotion, and progression. The implication of free radicals in different

steps of carcinogenesis is well documented. Some of the reactive oxygen and nitrogen species and their pathways could facilitate cancer development by DNA or other biomolecules damaging[2].

Chemotherapeutic drugs are still considered as the most important treatments for cancer, however, this kind of treatments trigger enormous side effects. Regarding this dilemma, ongoing research on natural medicine sources in form of functional foods or nutraceuticals has been attracting many scientists. Phytochemicals containing antioxidant properties showed capacity to

\*Corresponding author: Asep Awaludin Prihanto, Laboratory of Biochemistry, Faculty of Fisheries and Marine Science, Brawijaya University, Malang–65145, Indonesia.

Tel: 0341 553512

Fax: 0341 557837

E-mail: asep\_awa@ub.ac.id

Fundation Project: Supported by Directorate of Higher Education, Ministry of National Education, Republic of Indonesia (DIPA Brawijaya University No. 190/SK/2010).

Article history:

Received 8 Oct 2012

Received in revised form 28 Oct, 2nd revised form 20 Nov, 3rd revised form 2 Dec 2012

Accepted on 16 Dec 2012

Available online 28 Jan 2013

inhibit carcinogenesis[3]. Several studies discovered these antioxidant compounds are related to other bioactivities. For instances, *in-vitro* cytotoxicity in tumor cells[4], *in-vivo* cytotoxicity in experimental animals and anti-carcinogenesis[5,6].

There are numerous mechanisms of antioxidant substances as anticancer such as: by scavenging free radical[7], by inducing antioxidant enzymes[8], by modulating protein kinase and lipid kinase signaling pathway[9], by inhibiting cyclooxygenase-2 (COX-2) enzymatic activity[10], by affecting phytoestrogenic[11], by influencing nuclear transcription factor NF-κB[12], by inducing phase I or phase II metabolizing enzymes[13], inducing cell cycle arrest[14], and by inhibiting matrix metalloproteinases (MMP)-2 and MMP-9[15].

Based on World Health Organization data, more than 80% of world inhabitants depend on using plant for their medicine and mangroves have been widely used for that purpose[16,17]. *Acanthus ilicifolius* (*A. ilicifolius*) is a mangrove species that has been utilized traditionally for human remedies. Fruit, leaves, bark, and root of *A. ilicifolius* have been used for asthma, diabetes, hepatitis, inflammation, and rheumatoid treatments[17,20]. This mangrove species has been known to contain bioactive compounds *e.g.* triterpenoids, alkaloids, phenolic compounds, lignan, flavonoid, steroids, and terpenoids[21]. The leaf, root, stem, and bark of *A. ilicifolius* have been reported to be able to prevent tumor growth and cancer progression[22,23]. In order to intensively explore the potency of *A. ilicifolius* for cancer medication, the study of antioxidant and cytotoxic activity of *A. ilicifolius* flower was undertaken.

## 2. Materials and methods

### 2.1. Collection of plant samples

*A. ilicifolius* KÖENIG flowers were collected in November 2010 from Tempurejo village in Surabaya District of East Java, Indonesia and authenticated by the botanist from Department of Botany, Brawijaya University.

### 2.2. Phytochemical screening

Fresh *A. ilicifolius* flower were air dried and then homogenized into fine powder and stored in airtight bottles at 4 °C. Qualitative phytochemical analysis of the flower powder was done based on Farnsworth *et al.* method[24].

### 2.3. Preparation of flower extract of *A. ilicifolius*

The flower extract of *A. ilicifolius* was prepared by maceration. Around 100 g of fresh flower samples were macerated in 300 mL of acetone *p.a.*, methanol *p.a.*,

acetone 70%, methanol 80%, and aquades, respectively. The maceration procedure was repeated three times for 24 h. The extract was then filtered using Whatman filter paper. The filtrate was vacuum-evaporated and freeze dried before it was used for antioxidant and cytotoxic assay.

### 2.4. Antioxidant analysis

The antioxidant activity of flower extract was examined using diphenyl picryl hydrazil (DPPH) radical[25]. The 0.5 mmol DPPH was obtained by dissolving it with methanol. One mL of serial concentration of extract diluted in methanol was mixed in 3 mL of DPPH solution and then read their absorbance on spectrophotometer at 515 nm. Effective concentration (EC<sub>50</sub>) and time of effective concentration (T<sub>EC50</sub>) were analysed using statistic software of GraphPad Prism 5.0 and Excel 4.0. The antiradical efficiency was determined using the following equation:

$$AE = \frac{1}{EC_{50} \times T_{EC50}}$$

where: EC<sub>50</sub>: concentration to reduction 50 percent of free radical, T<sub>EC50</sub>: time needed to reach the EC<sub>50</sub>

### 2.5. Cytotoxic analysis

The cytotoxicity was conducted using brine shrimp lethality test[26]. The brine shrimp eggs were placed in 1 L of sea water, aerated for 48 h at 37 °C to hatch become nauplii. After 48 h, ten brine shrimp nauplii were placed in a small container filled with sea water. *A. ilicifolius* flower extract serially diluted with sea water were then added to the container. The lethality of brine shrimp was observed after 24 h of treatment was given. Probit analysis was used to determine lethal concentration (LC<sub>50</sub>) of *A. ilicifolius* flower methanol extract on nauplii.

### 2.6. Statistical analysis

Statistical analysis was performed using One-way analysis of variance (ANOVA) and followed by least square difference. Results were expressed as mean±SD from three replications. The *P* values < 0.01 were considered significant.

## 3. Results

The phytochemical screening test showed that the flower of *A. ilicifolius* contained several active compounds and dominated polar compounds (Table 1). The antiradical efficiency of *A. ilicifolius* flower extract was presented in Table 2. The percentage scavenging activity of the DPPH by 50% has been used to measure antioxidant activity. A comparable scavenging activity was observed among the

extract of *A. ilicifolius* flower, but the highest antiradical efficiency was obtained from the methanol extract.

**Table 1**  
Phytochemicals of *A. ilicifolius* flower.

Phytochemical	Inference
Triterpenoids	+
Saponin	+
Alkaloids	+
Phenolics	+
Flavonoids	+
Tannins	+
Steroids	–

+: Presence, -: Absence.

Table 2 shows the cytotoxic activity of *A. ilicifolius* flower extracts against brine shrimp nauplii. The cytotoxic activity of extracts of flower *A. ilicifolius* was statistically significant and the strongest activity was exhibited by methanol extract.

**Table 2**  
Antioxidant and cytotoxic activity of *A. ilicifolius* flower extracts (mean  $\pm$ SD).

Extract	Antioxidant (AE= $\times 10^{-5}$ )	Cytotoxic (LC <sub>50</sub> = $\mu$ g/mL)
Acetone	6.95 $\pm$ 0.15 <sup>b</sup>	1411 $\pm$ 27 <sup>b</sup>
Methanol	141.30 $\pm$ 2.10 <sup>d</sup>	22 $\pm$ 40 <sup>a</sup>
Acetone 70%	12.90 $\pm$ 0.90 <sup>c</sup>	2674 $\pm$ 66 <sup>c</sup>
Methanol 80%	10.90 $\pm$ 0.40 <sup>c</sup>	1348 $\pm$ 18 <sup>b</sup>
Water	0.0037 $\pm$ 0.0002 <sup>a</sup>	10610 $\pm$ 19 <sup>d</sup>

Different letters in the same column indicate significant statistically ( $P < 0.01$ ).

#### 4. Discussion

Bandaranayake stated that mangrove is one of active compounds source in the nature[17–19]. *A. ilicifolius* is clustered as true mangrove. The aerial part of *A. ilicifolius* has been extensively investigated for its secondary metabolite content and a large number of compounds have been structurally elucidated[21]. The flower of *A. ilicifolius* inferred positive triterpenoid and this compound in the leaf of this plant have been obtained. For instance lupeol,  $\alpha$  amyirin, olcanolic, and ursolic acids[21]. The flower of *A. ilicifolius* contained saponin and triterpenoidal saponin and these compounds have been found in the root of *A. ilicifolius*[27–31]. Tiwara *et al*[32] and Huo *et al*[33] reported that *A. ilicifolius* mangrove contained alkaloids *i.e.* acanthicifoline and benzoxazinium compounds. A wide range of phenolic compounds has been identified in *A. ilicifolius e.g.* acanfolioside, ilicifolioside, acteoside, verbascoside, and apigenin derivatives. Steroids were not screened in our samples while it have been detected in the leaf of *A. ilicifolius* in form of stigmasterol, campesterol, and sitosterol[21].

The methanol extract of *A. ilicifolius* contains phenolic substances clustered as antioxidant compounds[34]. The

antioxidant activity of *A. ilicifolius* extract is related to its capability as radical scavenger by transferring proton to free radical, nevertheless the antioxidant capacity of this mangrove species was weaker than ascorbic acid[25]. The capacity of antiradical efficiency of *A. ilicifolius* flower can be classified as medium. The extract was not in pure form; however, it can be categorized as a good and potential antioxidant agent.

The LC<sub>50</sub> of extracts or pure compounds on brine shrimp or cell line less than 100  $\mu$ g/mL is categorized as a potential cytotoxic and toxic substance[26,34]. The ethanol leaves extract of *A. ilicifolius* was found to be cytotoxic towards lung fibroblast (L–929) cells in 72 h MTT assay and the concentration required for 50% cell death was 18  $\mu$ g/mL[23], meanwhile the methanol extract of this plant was cytotoxic to Hela and  $\kappa$ B cell line[35]. Wöstmann and Liebezeit[21] reported that this mangrove contained antioxidant substances.

The highest antioxidant and cytotoxic activity were found on methanol extract. Methanol has been known more effective to dissolve active compounds in cells. Hence, it was easier to penetrate the cellular membrane to extract the intracellular ingredients from plant materials. Tiwari *et al.* stated that several active compounds will be obtained if methanol used as solvent in the extraction technique *i.e.* anthocyanins, saponins, tannins, flavones, and polyphenols[36]. These compounds have known as free radical scavenger, reactive species quencher, hydrogen donor, antioxidant enzymes activator, detoxification inducer, normal cell differentiation promoter, tumor production and proliferation cell inhibitor, and apoptosis inducer[23,37–41].

The methanol extract of *A. ilicifolius* flower exhibited both the highest antioxidant and anticancer activities in compare to other extracts. The cytotoxicity of methanol extract can be related to the antioxidant activity and synergism effect of multi–component in extract. Several active compounds contained in the flower of this mangrove have been revealed as a scavenging radical which may be able to inhibit carcinogenesis. Triterpenoid saponin showed its cytotoxicity in HeLa cells through both mitochondrial dysfunction and ER stress cell death pathways, while saponin suppressed tumor invasive and migration by inhibiting MMP–2 and MMP–9 activation[42]. Imai *et al.* determined that flavonoid effectively suppressed the proliferation of a human colon carcinoma cell line, COLO 201, through apoptosis induction while phenolics showed anticancer activity on cancer colon cell by arresting the cell cycle[43–45].

In conclusion, the methanol extract of *A. ilicifolius* flower is potential as antioxidant substances and cytotoxic compounds. Further studies are needed to identify the unknown antioxidant components to establish their pharmacological properties.

## Conflict of interest statement

We declare that we have no conflict of interest.

## Acknowledgement

This work was financially supported by Directorate of Higher Education, Ministry of National Education, Republic of Indonesia (DIPA Brawijaya University No. 190/SK/2010).

## Comments

### Background

The background has demonstrated linkage of the material that will be discussed. It also suggests that a study conducted by the author is still not done by other authors that make it feasible for journal of APJTB.

### Research frontiers

Studies on evaluation of active compound from *A. illicifolius* to cure several diseases have been reported. Hence the extract has not been evaluated on cancer cell toxicity yet. Therefore this article elucidated the best method to extract of antioxidant from the plant that has anti-cancer activity. This finding is novel and will inspire for further research.

### Related reports

Potent biological application of *A. illicifolius* was reported. However, in the next the biological properties of *A. illicifolius* should be revealed. From this preliminary report regarding *A. illicifolius* flower can enrich the biological application of mangrove.

### Innovations and breakthroughs

This paper showed the best method to extract active compound from *A. illicifolius* by using methanol. Furthermore, the author also confirmed that the extract has activity as anti-cancer with low dose that warrant for cancer therapeutic.

### Applications

The mangrove plant (*A. illicifolius*) contains a lot of active compounds that have activity as antioxidant. Moreover best method to extract the antioxidant is by using methanol extraction. The mangrove also important to remediate coastal area, so the conservation of the plant will have mutual benefit for environmental and also human health.

### Peer review

The article is explained the best method to extract

the active compounds from *A. illicifolius* that have high antioxidant activity. Moreover the extracts have anti-cancer activity with low dose that warrant cancer therapeutic. This finding is really important for community, scientist and pharmaceutical company as frontier information to explore *A. illicifolius* as source of anti-cancer material.

## References

- [1] Parkin DM, Bray F, Ferlay J, Pisani P. Global cancer statistics, 2002. *Cancer J Clin* 2005; **55**: 74–108.
- [2] Halliwell B. Oxidative stress and cancer. *Biochem J* 2007; **401**: 1–11.
- [3] Wang S, Meckling KA, Marccone MF, Kakuda Y, Tsao R. Can phytochemical antioxidant rich foods act as anti-cancer agents? *Food Res Int* 2011; **44**: 2545–2554.
- [4] Kalaivani T, Rajasekaran C, Mathew L. Free radical scavenging, cytotoxic, and hemolytic activities of an active antioxidant compound ethyl gallate from leaves of *Acacia nilotica* (L.) wild. Ex. Delile subsp. Indica (Benth.) Brenan. *J Food Sci* 2011; **76**: 144–149.
- [5] Murugan RS, Priyadarsini RV, Ramalingam K, Hara Y, Karunakaran D, Nagini S. Intrinsic apoptosis and NF- $\kappa$ B signaling are potential molecular targets for chemoprevention by black tea polyphenols in HepG2 cells *in vitro* and in a rat hepatocarcinogenesis model *in vivo*. *Food Chem Toxicol* 2010; **48**: 3281–3287.
- [6] Sukanuma M, Saha A, Fujiki H. New cancer treatment strategy using combination of green tea catechins and anticancer drugs. *Cancer Sci* 2011; **102**: 317–323.
- [7] Jia N, Xiong YL, Kong B, Liu Q, Xia X. Radical scavenging activity of black currant (*Ribes nigrum* L.) extract and its inhibitory effect on gastric cancer cell proliferation via induction of apoptosis. *J Funct Foods* 2012; doi:10.1016/j.jff.2012.01.009
- [8] Nirmala P, Ramanathan M. Effect of kaempferol on lipid peroxidation and antioxidant status in 1,2-dimethyl hydrazine induced colorectal carcinoma in rats. *Eur J Pharmacol* 2011; **654**: 75–79.
- [9] Liu JF, Ma Y, Wang Y, Du ZY, Shen JK, Peng HL. Reduction of lipid accumulation in HepG2 Cells by luteolin is associated with activation of AMPK and mitigation of oxidative stress. *Phytother Res* 2011; **25**: 588–596.
- [10] Suh Y, Afaq F, Johnson JJ, Mukhtar H. A plant flavonoid fisetin induces apoptosis in colon cancer cells by inhibition of COX2 and Wnt/EGFR/NF $\kappa$ B signaling pathways. *Carcinogenesis* 2009; **30**: 300–307.
- [11] Stokes Z, Chan S. Principles of cancer treatment by hormone therapy. *Surgery* 2009; **27**: 165–168.
- [12] Moon DO, Kim MO, Kang SH, Choi YH, Kim GY. Sulforaphane suppresses TNF- $\alpha$ -mediated activation of NF- $\kappa$ B and induces apoptosis through activation of reactive oxygen species-dependent caspase-3. *Cancer Lett* 2009; **274**: 132–142.

- [13] Androutsopoulos VP, Papakyriakou A, Vourloumis D, Tsatsakis AM, Spandidos DA. Dietary flavonoids in cancer therapy and prevention: Substrates and inhibitors of cytochrome P450 CYP1 enzymes. *Pharmacol Ther* 2010; **126**: 9–20.
- [14] Man S, Gao W, Zhang Y, Huang L, Liu C. Chemical study and medical application of saponins as anti-cancer agents. *Fitoterapia* 2010; **81**: 703–714.
- [15] Weng CJ, Yen GC. Chemopreventive effects of dietary phytochemicals against cancer invasion and metastasis: Phenolic acids, monophenol, polyphenol, and their derivatives. *Cancer Treat Rev* 2012; **38**: 76–87.
- [16] Firdaus M. Cytotoxic, antioxidant and antimicrobial activity of bark (*Xylocarpus granatum*) methanol extract. In: *Proceedings of International Conference on Basic Science*. Sulistyarti M, O'Neill M, Solich P, Paski E, Malherbe F, Coo L, et al.(eds). Malang: Galaxy Science Publisher; 2011, p. 62–64.
- [17] Bandaranayake WM. Traditional and medicinal uses of mangroves. *Mangroves Salt Marshes* 1998; **2**: 133–148.
- [18] Govindasamy C, Kannan R. Pharmacognosy of mangrove plants in the system of unani medicine. *Asian Pac J Trop Dis* 2012; **2**(Suppl 1): S38–S41.
- [19] Sundaram R, Ganesan R, Murugesan G. *In vitro* antiplasmodial activity of spiro benzofuran compound from mangrove plant of Southern India. *Asian Pac J Trop Med* 2011; **5**(5): 358–361.
- [20] Kumar KTMS, Gorain B, Roy DK, Zothanpuia, Samanta SK, Pal M, et al. Anti-inflammatory activity of *Acanthus ilicifolius*. *J Ethnopharmacol* 2008; **120**: 7–12.
- [21] Wöstmann R, Liebezeit G. Chemical composition of the mangrove holly *Acanthus ilicifolius* (Acanthaceae). *Senckenbergiana Mar* 2008; **38**: 31–37.
- [22] Graham JG, Quinn ML, Fabricant DS, Farnsworth NR. Plants used against cancer. *J Ethnopharmacol* 2000; **73**: 347–377.
- [23] Babu BH, Shylesh BS, Padikkala J. Tumour reducing and anticarcinogenic activity of *Acanthus ilicifolius* in mice. *J Ethnopharmacol* 2002; **79**: 27–33.
- [24] Farnsworth NR. Biological and phytochemical screening of plant. *J Pharm Sci* 1996; **55**: 225–265.
- [25] Sanchez-Moreno C, Larrauri JA, Saura-Calixto F. A procedure to measure the antiradical efficiency of polyphenols. *J Sci Food Agric* 1998; **76**: 270–276.
- [26] Ara J, Sultana V, Ehteshamul-Haque S, Qasim R, Uddin V. Cytotoxic activity of marine macro-algae on *Artemia salina* (Brine shrimp). *Phytother Res* 1999; **13**: 304–307.
- [27] Kokpol U, Chittawong V, Miles DH. Chemical constituents of the roots of *Acanthus ilicifolius*. *J Nat Prod* 1986; **49**: 355–356.
- [28] MA Hossain, MD Shah, C Gnanaraj, M Iqbal. *In vitro* total phenolics, flavonoids contents and antioxidant activity of essential oil, various organic extracts from the leaves of tropical medicinal plant *Tetrastigma* from Sabah. *Asian Pac J Trop Med* 2011; **4**(9): 717–721.
- [29] Ramachandran S, Rajasekaran S, Kumar KTM. Antidiabetic, antihyperlipidemic and antioxidant potential of methanol extract of *Tectona grandis* flowers in streptozotocin induced diabetic rats. *Asian Pac J Trop Med* 2011; **4**(8): 624–631.
- [30] Visweswara Rao P, Sujana P, Vijayakanth T, Dhananjaya Naidu M. *Rhinacanthus nasutus*—Its protective role in oxidative stress and antioxidant status in streptozotocin induced diabetic rats. *Asian Pac J Trop Dis* 2012; **2**(4): 327–330.
- [31] Kumar S, Narwal S, Kumar D, Singh G, Narwal S, Arya R. Evaluation of antihyperglycemic and antioxidant activities of *Saraca asoca* (Roxb.) De Wild leaves in streptozotocin induced diabetic mice. *Asian Pac J Trop Dis* 2012; **2**(3): 170–176.
- [32] Tiwara KP, Minocha PK, Masood M. Acanthicifoline – a new alkaloid from *Acanthus ilicifolius*. *Polish J Chem* 1980; **54**: 857–858.
- [33] Huo CH, Wang B, Lin WH, Zhao YY. Benzoxazinones from *Acanthus ilicifolius*. *Biochem Syst Ecol* 2005; **33**: 643–645.
- [34] Spavieri J, Allmendinger A, Kaiser M, Casey R, Hingley-Wilson S, Lalvani A, et al. Antimycobacterial, antiprotozoal, and cytotoxic potential of twenty-one brown algae (Phaeophyceae) from British and Irish waters. *Phytother Res* 2010; **24**: 1724–1729.
- [35] Khajure PV, Rathod JL. Potential anticancer activity of *Acanthus ilicifolius* extracted from the mangroves forest of Karwar, West Coast of India. *World J Technol* 2011; **1**: 1–6.
- [36] Tiwari P, Kumar B, Kaur M, Kaur G, Kaur H. Phytochemical screening and extraction. *Int Pharm Sci* 2011; **1**: 98–106.
- [37] Breton F, Cérantola S, Gall EA. Distribution and radical scavenging activity of phenols in *Ascophyllum nodosum* (Phaeophyceae). *J Exp Mar Biol Ecol* 2011; **399**: 167–172.
- [38] Nielsen MF, Ingold KU. Kinetic solvent effects on proton and hydrogen atom transfers from phenols. Similarities and differences. *J Am Chem Soc* 2006; **128**: 1172–1182.
- [39] Fernandez-Pachon MS, Berna G, Otaolauruchi E, Troncoso AM, Martin F, Garcia-Parrilla MC. Changes in antioxidant endogenous enzymes (activity and gene expression levels) after repeated red wine intake. *J Agric Food Chem* 2009; **57**: 65780–65783.
- [40] Hsu YW, Tsai CF, Chen WK, Huang CF, Yen CC. A subacute toxicity evaluation of green tea (*Camellia sinensis*) extract in mice. *Food Chem Toxicol* 2011; **49**: 2624–2630.
- [41] Chahar MK, Sharma N, Dobhal MP, Joshi Y. Flavonoids: A versatile source of anticancer drugs. *Pharmacog Rev* 2011; **5**: 1–12.
- [42] Man S, Gao W, Zhang Y, Huang L, Liu C. Chemical study and medical application of saponins as anti-cancer agents. *Fitoterapia* 2010; **81**: 703–714.
- [43] Imai M, Kikuchi H, Denda T, Ohyama K, Hirobe C, Toyoda H. Cytotoxic effects of flavonoids against a human colon cancer derived cell line, COLO 201: A potential natural anti-cancer substance. *Cancer Lett* 2009; **276**: 74–80.
- [44] Gonzalez-Sarrias A, Li L, Seeram NP. Anticancer effects of maple syrup phenolics and extracts on proliferation, apoptosis, and cell cycle arrest of human colon cells. *J Funct Foods* 2011; doi:10.1016/j.jff.2011.10.004.
- [39] Gonzalez-Sarriás A, Espín JC, Tomas-Barberán FA, García-Conesa MT. Gene expression, cell cycle arrest, and MAPK signalling regulation in Caco-2 cells exposed to ellagic and its metabolites, urolithins. *Mol Nutr Food Res* 2009; **53**: 686–698.