



Contents lists available at [ScienceDirect](http://www.sciencedirect.com)

Asian Pacific Journal of Tropical Biomedicine

journal homepage: www.elsevier.com/locate/apjtb



Document heading doi:10.1016/S2221-1691(11)60003-8 © 2011 by the Asian Pacific Journal of Tropical Biomedicine. All rights reserved.

ETM study of electroporation influence on cell morphology in human malignant melanoma and human primary gingival fibroblast cells

Nina Skolucka¹, Malgorzata Daczewska², Jolanta Saczko¹, Agnieszka Chwilkowska¹, Anna Choromanska¹, Malgorzata Kotulska³, Iwona Kamińska¹, Julita Kulbacka^{1*}

¹Department of Medical Biochemistry, Medical University, Chalubinskiego 10 St., 50–368 Wrocław, Poland

²Department of General Zoology, Zoological Institute, University of Wrocław, Sienkiewicza 21 St., 50–335 Wrocław, Poland

³Institute of Biomedical Engineering and Instrumentation, Wrocław University of Technology, ul. Wybrzeże Wyspińskiego 27, 50–370 Wrocław, Poland

ARTICLE INFO

Article history:

Received 6 February 2011

Received in revised form 22 February 2011

Accepted 13 March 2011

Available online 1 April 2011

Keywords:

Electroporation

Electron microscopy

Human melanoma cells

Primary fibroblasts

Cell morphology

Transmission electron microscope

Malignant cell

Normal cell

Intracellular effect

Electric field

Proliferation

Ultrastructural analysis

ABSTRACT

Objective: To estimate electroporation (EP) influence on malignant and normal cells. **Methods:** Two cell lines including human malignant melanoma (Me-45) and normal human gingival fibroblast (HGFs) were used. EP parameters were the following: 250, 1 000, 1 750, 2 500 V/cm; 50 μ s by 5 impulses for every case. The viability of cells after EP was estimated by MTT assay. The ultrastructural analysis was observed by transmission electron microscope (Zeiss EM 900). **Results:** In the current study we observed the intracellular effect following EP on Me-45 and HGF cells. At the conditions applied, we did not observe any significant damage of mitochondrial activity in both cell lines treated by EP. Conversely, we showed that EP in some conditions can stimulate cells to proliferation. Some changes induced by EP were only visible in electron microscopy. In fibroblast cells we observed significant changes in lower parameters of EP (250 and 1 000 V/cm). After applying higher electric field intensities (2 500 V/cm) we detected many vacuoles, myelin-like bodies and swallowed endoplasmic reticulum. In melanoma cells such strong pathological modifications after EP were not observed, in comparison with control cells. The ultrastructure of both treated cell lines was changed according to the applied parameters of EP. **Conclusions:** We can claim that EP conditions are cell line dependent. In terms of the intracellular morphology, human fibroblasts are more sensitive to electric field as compared with melanoma cells. Optimal conditions should be determined for each cell line. Summarizing our study, we can conclude that EP is not an invasive method for human normal and malignant cells. This technique can be safely applied in chemotherapy for delivering drugs into tumor cells.

1. Introduction

Cell membranes are sensitive to physicochemical conditions, such as biochemical environment, temperature, mechanical stress, and electro-magnetic field. It was discovered that under diverse conditions plasma membrane of a cell loses its stability. Lipids in membrane change their conformation after application of electric field with proper parameters, thus they form non-stable hydrophilic

pores[1–3]. This event was observed in 1958 for the first time. It was registered as unexplained, sudden increase of ionic conductivity through lipid membrane, and it is called electroporation (EP)[4–6]. The mechanism of EP is still not fully understood. Currently EP is used in cell culture as the purest available method of gene transfection[7]. First experiments with EP on animal models took place in the end of the 80s of the 20th century. Effectiveness of these attempts triggered research on the human organism, which began in the 90s[8–10], which resulted in the most important recent EP application—electrochemotherapy (ECT), accepted for treatment in EU in 2005. Electrochemotherapy uses the EP effect to facilitate transport of cytotoxic drugs inside malignant cell across cell membrane. Intentionally, the selected cytostatics were very poorly transported inside cells and demonstrated higher affinity to cancer cells. The experiments showed very significantly increased

*Corresponding author: Julita Kulbacka, Department of Medical Biochemistry, Medical University, Chalubinskiego 10 St., 50–368 Wrocław, Poland.

E-mail: jkulbacka@gmail.com

Foundation Project: Supported by statutory funds of Medical Wrocław University and research fellowship within "Development Program of Wrocław Medical University" funded from European Social Fund, Human Capital, National Cohesion Strategy (Contract No. UDA-POKL.04.01.01-00-010/08-00).

intracellular drug concentration limited to cancer cells exposed to the electric impulses. ECT can be applied locally in case of accessible cancers or tumors with superficial localization and internal organs. Impulses of the electric field in ECT are typically 100 μ s long, with the intensity of 1 kV/cm. Cisplatin and bleomycin are used very often in this local treatment. Its effectiveness increases from 2–8 times in case of cisplatin, 500 times for bleomycin, and 3–5 times in case of actinomycin-D^[11,12]. Current research shows very promising medical application of EP, such as immunogenetherapy^[10] and ECT with nanoimpulses^[9].

Electrochemotherapy has been proved to be an excellent technique in melanoma treatment raising the complete response (CR) factor from 5% (standard chemotherapy) to 45% (ECT with bleomycin systemically), 77% (bleomycin locally), and 74% (cisplatin locally)^[13–15]. Even new multidrug chemotherapy reaches CR not higher than 25% (for example Cisplatin CDDP +dacarbazine DITC, interleukin 2 (IL-2)+ interferon 22b (INF-22b))^[16]. Except for melanoma, ECT was mainly applied in basal breast cancer non-responsive to conventional treatment or various superficial neck and head cancers. CR in treatment of these cancers by ECT was about 55%^[13]. Enhancing currently used therapies by EP open new perspectives in anticancer treatment, especially in case of multidrug resistance^[5,9]. Application of combined therapy can be more efficient than usual chemotherapy and less radical than surgical intervention.

Although EP is effective in facilitating molecular transport, the intracellular effects of its application have not been sufficiently studied. The main intention of the presented study was examination of EP influence on cell morphology in two cell lines (normal and malignant) by electron transmission microscopy. The present paper shows unique studies of the EP influence on cell ultrastructure. The analysis of cell ultrastructure and determination of cells viability appeared proper techniques to study the EP impact.

2. Material and methods

2.1. Cell culture

We used human melanoma cells (Me-45) and human gingival fibroblasts (HGFs) from primary culture derived from a 30-year-old patient. Fibroblasts were isolated from soft tissue of gingival. The cells were isolated from the healthy gingival tissues according to the procedure described by Saczko *et al*^[17]. Both cell lines were harvested in Dulbecco medium (Sigma) with 10% of FBS and addition of antibiotics. The cells were maintained in a humidified atmosphere at 37 °C in 5% CO₂.

2.2. Electroporation (EP)

The EP was carried out by use of ECM 830 (BTX) square-wave electroporator. At the high voltage mode, the electroporator provides electrical pulses with the magnitude 505–3 000 V/cm, 10–600 μ s long, in the series

of 1–99 impulses separated by the time interval of 100 ms–10 s. Different parameters of EP applied in our study were: 250, 1 000, 2 500, 3 250 and 4 000 V/cm; 50 ms in 5 impulses for every case. As electrodes we used two thin aluminum parallel plates, 4 mm apart. They were connected to the voltage generator and produced a uniform electric field in the cuvette (Cuvettes Plus 640, 800 μ L). Cells in suspension were centrifuged for 5 min at 1 500 rpm and resuspended in the EP buffer with low electrical conductivity (10 mM phosphate, 1 mM MgCl₂, 250 mM sucrose, pH 7.4). Cells for examination by transmission electron microscopy were left for 5 min after pulsation, then cells were fixed for 30 min in 2.5% (vol/vol) glutaraldehyde and 0.1 M phosphate buffer (pH 7.4). Cells for MTT assay were left for 20 min with addition of 1 600 μ L DMEM after pulsation, then washed and centrifuged twice with DMEM.

2.3. MTT assay

The 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) test was used for the assessment of cell viability. After EP experiment, the cells were seeded into 96-well microculture plates (Nunc, Nunclon™ Surface) at the concentration of 5×10^3 cells/well and incubated for 24 h at 37 °C and 5% CO₂. Then the medium of each well was replaced with 10 μ L of 0.5 mg/mL MTT stock solution [3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide; Sigma Saint Louis, MO; *In Vitro* Toxicology Assay] diluted in 90 μ L PBS. After 2 h of incubation, isopropanol with 0.04 M HCl was added (100 μ L/well). The absorbance was determined using a multiwell scanning spectrophotometer at 570 nm (Labsystem Multiscan MS type 352, Helsinki, Finland). Mitochondrial function was expressed as a percentage of viable cells under treatment relative to untreated control cells (the growth fraction of non-treated cells equals 1); the outstanding results not taken into account.

2.4. Transmission electron microscopy

Morphological assessment of cells after and before EP was examined by transmission electron microscope Zeiss EM 900. The cells were prepared according to the standard procedure. Cells were fixed for 30 min in 2.5% (vol/vol) glutaraldehyde and 0.1 M phosphate buffer (pH 7.4). After postfixation in 1% (wt/vol) osmium tetroxide, cells were dehydrated through a graded series of alcohol and propylene oxide and embedded in Epon.

2.5. Statistical analysis

The significance of the difference between mean values of different groups of cells was assessed by Student's *t*-test with *P*-value of *P*=0.05, to show the statistical significance. Results were analyzed by commercial software Statistica 9.0.

3. Results

The sensitivity of both cell lines to EP was dependent on the applied pulse parameters. However, EP did not affect cells viability significantly. Cells viability after EP (250, 1 000 and 2 500 V/cm) was shown in Figure 1 (A and B). Since pulse length in these experiments is 50 μ s, the energy provided

in EP here corresponds to twice lower field intensities as compared with experiments at pulses of 100 μ s, which are more typically used in ECT. Hence, the intensity of 2 500 V/cm, at 50 μ s, represents standard conditions of ECT. The mitochondrial activity in fibroblast cells was on the control




Figure 1. The cells viability after electroporation at pulse length of 50 μ s.

A: human fibroblasts; B: human melanoma; Error bars shown are means \pm SD, $n=4$; $*P \leq 0.05$.




Figure 2. The cells ultrastructure in fibroblast cells.

A: control cells; B: cells EP at 250 V/cm; C: cells EP at 1 000 V/cm; D: cells EP at 2 500 V/cm.

GA: Golgi apparatus; RER: rough endoplasmic reticulum; N: nucleus; IF: intermediate filament; L: lysosomes V: vacuole; RB: residual body; M: mitochondria.

level (about 100% of cells) and only increased at 2 500 V/cm. In case of Me-45 cells we observed higher mitochondrial activity, which reached more than 100% of cells and decreased with increasing voltage.

In Figure 2 and 3 ultrastructural morphology of human fibroblast and melanoma cells was presented. In the control fibroblast cells (Figure 2A), no significant pathological changes were observed. A lot of vacuoles containing a flocculent and granular material were found. The Golgi apparatus and mitochondria established regular morphological structures. We could also observe joined primary lysosomes and heterochromatin on the left side. After application of EP of 250 V/cm, (Figure 2B) we detected the abnormal, folded nuclear membrane and a lot of vacuoles with fibrous and flocculent material. The characteristic for these cells vimentin intermediate filaments forming regular clusters is also visible. When we applied electric field of higher intensities (1 000 V/cm, Figure 2C) the swollen rough endoplasmic reticulum was observed. There are numerous vacuoles with flocculent material. The vacuoles with matrix

appeared and also the characteristic vimentin intermediate filaments were found. There are also visible lysosomes (black color). After EP at 2 500 V/cm (Figure 2D) extended rough endoplasmic reticulum was observed. The abnormal folded nuclear membrane appeared. There are almost no vacuoles. We noticed a lot of translucent mitochondrial matrix. The shape of mitochondria was changed.

In case of melanoma control cells (Figure 3A) also insignificant pathological changes were observed. Only the folded nuclear shape (right side of the figure) appeared and plenty of melanin granules were found. After application of EP at 250 V/cm (Figure 3B), the strongly folded nuclear membrane appeared. The fragment of cell external membrane was observed. The mitochondria had translucent matrix; a few of crista and the decreased number of residual bodies was observed. Stronger EP at 2 500 V/cm (Figure 3C) generated abnormal mitochondrial structures with translucent matrix and also a few crista. A lot of myelin bodies could be noticed. The folded nuclear membrane and residual bodies appeared.




Figure 3. The cells ultrastructure in Me-45 cells.

A: control cells; B: cells EP at 250 V/cm; C: cells EP at 2 500 V/cm.

EXC: extruded from the cell a part of cytoplasm; hM: human melanin; GA: Golgi apparatus; N: nucleus; L: lysosomes; V: vacuole; RB: residual body; M: mitochondria.

4. Discussion

EP is a widely applicable method of introducing molecules into cells. It is based on transient permeabilization of the cell membrane by applying electric field pulses of high intensity. Although the technique is sufficiently effective, the morphological side-effects, which could be field dependent, have not been sufficiently studied. In the current study we reported the intracellular effect following EP on Me-45 and HGF cells. At the conditions applied, we did not observe any significant damage of mitochondrial activity in both cell lines treated by EP. Conversely, we showed that EP in some conditions can stimulate cells to proliferation. Some changes induced by EP were only visible in electron microscopy. In fibroblast cells we observed significant changes in lower parameters of EP (250 and 1000 V/cm). After applying higher electric field intensities (2500 V/cm) we

detected many vacuoles, myelin-like bodies and swallowed endoplasmic reticulum (ER). In melanoma cells such strong pathological modifications after EP were not observed, in comparison with control cells. The ultrastructure of both treated cell lines was changed according to the applied parameters of EP.

In the present study we showed the difference of the influence of EP on morphology of Me-45 and HGF cells. Ghosh *et al* also examined human fibroblast cell line (WI-38/VA-130), however derived from lung. They applied current-voltage measurements to detect pore formation and monitor pore resealing in the plasma membrane of adherent mammalian fibroblastic cells. The authors observed that the response of the cells to pulsation varied with the degree of confluency of the cell layer on the electrode^[18]. Other authors examined melanoma xenografts grown in nude mice. Spugini *et al* applied eight biphasic pulses using an

electric field of 1 250 or 2 450 V/cm. In electron microscopy they detected defects in the dynamic assembly of lipids and proteins, which generated “areas with rough structure” and intensive clustering of intramembrane proteins^[19].

According to the results obtained in this study, we can claim that EP conditions are cell line dependent. In terms of the intracellular morphology, human fibroblasts are more sensitive to electric field as compared with melanoma cells. Optimal conditions should be determined for each cell line. Summarizing our study, we can conclude that EP is not an invasive method for human normal and malignant cells. This technique can be safely applied in chemotherapy for delivering drugs into tumor cells.

Conflict of interest statement

We declare that we have no conflict of interest.

Acknowledgments

This study was supported by statutory funds of Medical Wrocław University and research fellowship within “Development program of Wrocław Medical University” funded from European Social Fund, Human Capital, National Cohesion Strategy” (Contract No. UDA–POKL.04.01.01–00–010/08–00).

References

- [1] Kotulska M, Koronkiewicz S, Kalinowski S. Self-similar processes and flicker noise from a fluctuating nanopore in a lipid membrane. *Phys Rev E Stat Nonlin Soft Matter Phys* 2004; **69**: 031920.
- [2] Kotulska M. Natural fluctuations of an electropore show fractional Lévy stable motion. *Biophys J* 2007; **92**(7): 2412–2421.
- [3] Dyrka W, Augusti AT, Kotulska M, Ion flux through membrane channels—an enhanced algorithm for the poisson–nernst–planck model. *J Comput Chem* 2008; **29**: 1876–1888.
- [4] Böckmann RA, de Groot BL, Kakorin S, Neumann E, Grubmüller H. Kinetics, statistics, and energetics of lipid membrane electroporation studied by molecular dynamics simulations. *Biophys J* 2008; **95**(4): 1837–1850.
- [5] Skořucka N, Sączko J, Kotulska M, Kulbacka J, Choromańska A. Electroporation and its application. *Pol Merk Lek* 2010; **168**: 501.
- [6] Stämpfli R. Reversible electrical breakdown of the excitable membrane of a ranvier node. *Ann Acad Bras Ciens* 1958; **30**: 57–63.
- [7] Neumann E, Schaefer–Ridder M, Wang Y, Hofschneider PH. Gene transfer into mouse lyoma cells by electroporation in high electric fields. *EMBO J* 1982; **1**(7): 841–845.
- [8] Mir LM, Belehradec M, Domenge C, Orlowski S, Poddevin B, Belehradec J, et al. Electrochemotherapy, a new antitumor treatment: first clinical trial. *C R Acad Sci III* 1991; **313**(13): 613–618.
- [9] Belehradec M, Domenge C, Luboinski B, Orlowski S, Belehradec J, Mir LM. Electrochemotherapy, a new antitumor treatment. First clinical phase I–II trial. *Cancer* 1993; **72**(12): 3694–3700.
- [10] Daud AI, Deconti RC, Andrews S, Urbas P, Riker AI, Sondak VK, et al. Phase I trial of interleukin–12 plasmid electroporation in patients with metastatic melanoma. *J Clin Oncol* 2008; **26**(36): 5896–5903.
- [11] Mir LM, Gehl J, Sersa G, Collins CG, Garbay JR, Billard V, et al. Standard operating procedures of the electrochemotherapy: Instructions for the use of bleomycin or cisplatin administered either systemically or locally and electric pulses delivered by the Cliniporator™ by means of invasive or non-invasive electrodes. *Eur J Cancer* 2006; **S4**: 14–25.
- [12] Jaroszeski MJ, Coppola D, Nesmith G, Pottinger C, Hyacinthe M, Benson K, et al. Effects of electrochemotherapy with bleomycin on normal liver tissue in a rat model. *Eur J Cancer* 2001; **37**(3): 414–421.
- [13] Sersa G. The state-of-the-art of electrochemotherapy before the ESOPE study: advantages and clinical uses. *Eur J Cancer* 2006; **S4**(11): 52–59.
- [14] Ives NJ, Stowe RL, Lorigan P, Wheatley K. Chemotherapy compared with biochemotherapy for the treatment of metastatic melanoma: a meta-analysis of 18 trials involving 2 621 patients. *J Clin Oncol* 2007; **25**(34): 5426–5434.
- [15] Kranjc S, Cemazar M, Grosel A, Sentjurs M, Sersa G. Radiosensitising effect of electrochemotherapy with bleomycin in LPB sarcoma cells and tumors in mice. *BMC Cancer* 2005; **16**(5): 115.
- [16] Bar J, Yerushalmi R, Shapira–Frummer R, Kutchuk I, Sulkes A, Gutman H, et al. Concurrent chemobiotherapy with cisplatin, dacarbazine, decrescendo interleukin–2 and interferon alpha2b in patients with metastatic melanoma. *Oncol Rep* 2008; **20**(6): 1533–1538.
- [17] Sączko J, Dominiak M, Kulbacka J, Chwiłkowska A, Krawczykowska H. A simple and established method of tissue culture of human gingival fibroblasts for gingival augmentation. *Folia Histochem Cytobiol* 2008; **46**: 117–119.
- [18] Ghosh PM, Keese CR, Giaever I. Monitoring electroporation in the plasma membrane of adherent mammalian cells. *Biophys J* 1993; **64**: 1602–1609.
- [19] Spugnini EP, Arancia G, Porrello A, Colone M, Formisano G. Ultrastructural modifications of cell membranes induced by “electroporation” on melanoma xenografts. *Microsc Res Tech* 2007; **70**(12): 1041–1045.