



Document heading doi:10.12980/APJTB.4.2014APJTB-2014-0077 © 2014 by the Asian Pacific Journal of Tropical Biomedicine. All rights reserved.

Insecticidal activity of the essential oil of *Thymus transcaspicus* against *Anopheles stephensi*

Leila Dargahi¹, Kamal Razavi–Azarkhiavi², Mohammad Ramezani³, Mohammad Reza Abae⁴, Javad Behravan^{5*}

¹Neuroscience Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

²Department of Pharmacodynamics and Toxicology, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran

³Pharmaceutical Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

⁴Department of Medical Entomology, School of Public Health and Institute of Health Research, Tehran University of Medical Sciences, Tehran, Iran

⁵Biotechnology Research Center, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran

ARTICLE INFO

Article history:

Received 11 Feb 2014

Received in revised form 5 May 2014

Accepted 8 Jun 2014

Available online 3 Jul 2014

Keywords:

Thymus transcaspicus

Anopheles stephensi

Insecticidal activity

Essential oil

ABSTRACT

Objective: To investigate the insecticidal activity of the essential oil of *Thymus transcaspicus* (*T. transcaspicus*) against *Anopheles stephensi* (*An. stephensi*).

Methods: *An. stephensi* were exposed to 31, 63, 125 and 250 µg/L of essential oil of *T. transcaspicus* for 24 h.

Results: The most toxicity was observed at 250 µg/L of essential oil with the LC₅₀ values of 134.1 µg/L after 24 h.

Conclusions: The essential oil of *T. transcaspicus* exhibited strong insecticidal activity against *An. stephensi* which can be attributed to its constituent especially carvacrol and thymol phenols.

1. Introduction

Control of the insects usually necessitates the use of synthetic insecticides, whereas these chemical-based insecticides lead to serious concerns. Expensiveness, being environmentally hazardous, growing incidence of insect resistance, bioaccumulation through food chains and toxic effects on human health, encouraged researchers to find naturally occurring, reasonable and more environmentally friendly agents^[1,2]. Therefore, several plants have been studied as insecticide, larvicidal and repellent agents^[3–6].

Thymus is a genus of plant which contains different species. About 350 species have been reported from this genus, over 14 of which are identified in Iranian flora^[7]. The antibacterial, insecticidal and antifungal properties of *Thymus* species have been reported in various studies^[8].

Thymus transcaspicus (*T. transcaspicus*) (Khorasan thyme) is an aromatic plant which is native to northern regions of Iran^[9]. It has been illustrated that carvacrol and thymol are the major constituents of this plant^[10].

The insects play an important role as vectors for transmission of a variety of diseases including malaria, filariasis, yellow fever, dengue fever and Japanese encephalitis between human populations^[11]. More than 300 million people are infected with malaria every year worldwide^[3]. *Anopheles stephensi* (*An. stephensi*) is the main vector of human malaria in the Middle East and in the Persian Gulf area^[12,13]. It has been also demonstrated that *An. stephensi* is the most prevalent anopheline species in the southern areas of Iran^[13].

In light of the above knowledge, the present study has been designed to investigate the insecticidal activity of the essential oil of *T. transcaspicus* against *An. stephensi*.

2. Material and methods

2.1. Preparation of essential oil

T. transcaspicus was collected from the Tandoureh National Park (Khorasan Province, Iran), identified by the

*Corresponding author: Javad Behravan, Ph.D., Biotechnology Research Center, Mashhad University of Medical Sciences, Mashhad, P. O. Box, 91775–1365, I.R. Iran.

Tel: +98–511–882–3255

Fax: +98–511–882–3251

E-mail: behravanj@mums.ac.ir

Foundation Project: Supported by Faculty of Medicine, Mashhad University of Medical Sciences with the Grant No. 1035.

herbarium of School of Pharmacy, Mashhad University of Medical Sciences and a voucher sample was deposited (Voucher No. 153–2020–18).

The plants were cleaned; the leaves, flowers and stems were separated and dried in the dark. The dried plants were then powdered with a mechanical grinder. Batches of 75 g powder were subjected to hydrodistillation for 4 h, using a Clevenger–type apparatus^[14].

2.2. *An. stephensi* cultures

An. stephensi was obtained from insectarium of the School of Public Health, Tehran University of Medical Sciences. The larvae were fed with tetramin and yeast powder.

2.3. Insecticidal activity of the essential oil on *An. stephensi*

The larvicidal activity of essential oil was examined according to the standard method suggested by World Health Organization^[15]. Different concentrations of essential oil of *T. transcaspicus* (31, 63, 125, 250 µg/L) in acetone were prepared. Dechlorinated tap water was used for all the experimental groups. Twenty five early fourth instar larvae were exposed to different concentrations in a working volume of 250 mL in 400–mL glass beaker. Two replicates for each concentration and controls (acetone) were included for larval bioefficacy. The larval mortality for each concentration was recorded after 24 h of exposure.

2.4. Data analysis

LC₅₀ (lethal concentration to cause 50% mortality in the population) and LC₉₀ (lethal concentration to cause 90% mortality in the population) values were calculated by probit analyses using SPSS (2001). The percentage mortality was corrected using Abbott's formula.

3. Results

The final yield of the hydrodistillation of *T. transcaspicus* was 0.7% (v/w). The influence of different concentrations of *T. transcaspicus* essential oil against *An. stephensi* after 24 h is shown in Figure 1.




Figure 1. Percentage mortality response of *An. stephensi* exposed to different concentrations (µg/L) of *T. transcaspicus* for 24 h.

The concentration–dependent lethal toxicity was occurred due to essential oil treatment. The most toxicity was observed at 250 µg/L. Probit analysis indicated that LC₅₀ and LC₉₀ values of the essential oils against *An. stephensi* were 134.1 (52.8–328.1) and 248.0 (159.0–457.4) µg/L.

4. Discussion

Insects are responsible for considerable damage to human environment and health. About 15% of the world's crop losses are occurred due to insects^[16]. Moreover, insects play an important role as vectors in transmission of different diseases among human. These diseases include malaria, yellow fever, dengue fever, Japanese encephalitis and filariasis^[11]. It has been suggested that the best way for protecting human health against the vector–borne diseases is vector control^[17]. The extensive use of synthetic insecticides have led to disturbing consequences such as environmental hazards, bioaccumulation of toxic agents in food, incidence of insect resistance and many human health problems^[2]. Botanical insecticides have been applied in control of insects for at least two thousand years in Asia^[18]. The considerable interest for new botanical insecticides is due to their physiological activity, biodegradability and bioefficiency^[19]. The genus *Thymus* L. (Lamiaceae) comprises more than 250 herbaceous perennial and sub–shrubs throughout the world. Over 14 species of this genus such as *T. transcaspicus* are represented in the flora of Iran^[20]. It has been illustrated that *Thymus* L. possesses various biological properties such as antifungal, antimicrobial and insecticidal^[8,21].

In the present study, the insecticidal potential of *T. transcaspicus* (Khorasan thyme) against malaria vector (*An. stephensi*) was investigated. The essential oil of *T. transcaspicus* was prepared by hydrodistillation using a Clevenger–type apparatus.

The insecticidal activity of the *T. transcaspicus* essential oil is significant when considering that more than 90% of *An. stephensi* larvae were killed due to the exposure to 250 µg/L of the essential oil (LC₅₀: 134.1 µg/L). *An. stephensi* is the major vector of malaria in the southern area of Iran, with an expansion through the Middle East into the Indian subcontinent^[22]. The widespread resistance from this species has been reported against synthetic insecticides including malathion, benzene hexachloride and DDT^[23].

It has been shown that *Thymus* species have strong insecticidal activity against various insects such as *Aedes aegypti* (L)^[24], *Tribolium castaneum* and *Callosobruchus maculatus*^[25], *Aedes albopictus*^[26] and *Sitophilus oryzae*^[21]. The insecticidal activity of *T. transcaspicus* could be attributed to its chemical composition. Thymol (56.4%), γ–terpinene (7.7%), carvacrol (7.6%) and *p*–cymene (6.3%) are the main compounds of *T. transcaspicus* essential oil^[10]. Carvacrol and thymol are the monoterpene phenols which have broad toxicity against insects. Carvacrol [2–methyl–5–(1–methylethyl) phenol], is a constituent of essential oils produced by various aromatic plants and species such as *Thymus* species^[27]. It has been proved that carvacrol is effective against different insect pests like *Leucania separata*^[28]. Thymol which is another major monoterpene phenol was reported to have insecticidal activity against

An. stephensi^[29].

The results obtained indicate that the essential oil of *T. transcaspicus* has a significant insecticidal activity against malaria vector (*An. stephensi*). This toxicity against insects can be attributed to its constituent especially carvacrol and thymol phenols.

Conflict of interest statement

We declare that we have no conflict of interest.

Acknowledgements

The authors are thankful to the Vice Chancellor of Research, Mashhad University of Medical Sciences for financial support. This research was funded by Faculty of Medicine, Mashhad University of Medical Sciences with the Grant No. 1035. The results described in this paper are part of the Pharm. D. thesis of L. Dargahi.

References

- [1] Majumder P, Mondal HA, Das S. Insecticidal activity of *Arum maculatum* tuber lectin and its binding to the glycosylated insect gut receptors. *J Agric Food Chem* 2005; **53**: 6725–6729.
- [2] Klein W. *The future for insecticides, need and prospects*. In: Metcalf RL, Mckelvey JJ, editors. New York, USA: Wiley Interscience; 1976, p. 65.
- [3] Ahmad N, Fazal H, Abbasi BH, Iqbal M. *In vitro* larvicidal potential against *Anopheles stephensi* and antioxidative enzyme activities of *Ginkgo biloba*, *Stevia rebaudiana* and *Parthenium hysterophorous*. *Asian Pac J Trop Med* 2011; **4**: 169–175.
- [4] Kweka EJ, Nyindo M, Mosha F, Silva AG. Insecticidal activity of the essential oil from fruits and seeds of *Schinus terebinthifolia* Raddi against African malaria vectors. *Parasit Vectors* 2011; **4**: 129.
- [5] Rajashekar Y, Gunasekaran N, Shivanandappa T. Insecticidal activity of the root extract of *Decalepis hamiltonii* against stored-product insect pests and its application in grain protection. *J Food Sci Technol* 2010; **47**: 310–314.
- [6] Shahi M, Hanafi-Bojd AA, Iranshahi M, Vatandoost H, Hanafi-Bojd MY. Larvicidal efficacy of latex and extract of *Calotropis procera* (Gentianales: Aselepiadaceae) against *Culex quinquefasciatus* and *Anopheles stephensi* (Diptera: Culicidae). *J Vector Borne Dis* 2010; **47**: 185–188.
- [7] Tabrizi L, Koocheki A, Moghaddam PR, Mahallati MN. Chemical composition of the essential oils from *Thymus transcaspicus* in natural habitats. *Chem Nat Compd* 2010; **46**: 121–124.
- [8] Imelouane B, Amhamdi H, Wathelet JP, Ankit M, Khedid K, El Bachiri A. Chemical composition and antimicrobial activity of the essential oil of thyme (*Thymus vulgaris*) from Eastern Morocco. *Int J Agric Biol* 2009; **11**: 205–208.
- [9] Zamani N, Mianabadi M, Abdolzadeh A. Changes in anti-oxidant activity of *Thymus transcaspicus* (Klokov) during growth and developmental stages. *J Cell Mol Res* 2011; **3**(1): 12–18.
- [10] Miri R, Ramezani M, Javidnia K, Ahmadi L. Composition of the volatile oil of *Thymus transcaspicus* Klokov from Iran. *Flavour Fragr J* 2002; **17**: 245–246.
- [11] Service M. *Medical entomology for students*. Cambridge, UK: Cambridge University Press; 2004.
- [12] Dash AP, Adak T, Raghavendra K, Singh OP. The biology and control of malaria vectors in India. *Curr Sci* 2007; **92**: 1571–1578.
- [13] Hanafi-Bojd AA, Vatandoost H, Oshaghi MA, Haghdoost AA, Shahi M, Sedaghat MM, et al. Entomological and epidemiological attributes for malaria transmission and implementation of vector control in southern Iran. *Acta Trop* 2012; **121**: 85–92.
- [14] Clevenger JF. Apparatus for the determination of volatile oil. *J Am Pharm Assoc* 1928; **17**: 345–349.
- [15] World Health Organization. Instructions for determining susceptibility or resistance of mosquito larvae to insecticides. Geneva: World Health Organization; 1981. [Online] Available from: http://apps.who.int/iris/bitstream/10665/69615/1/WHO_VBC_81.807_eng.pdf?ua=1 [Accessed on 4th February, 2014]
- [16] Schultz JC. Biochemical ecology: how plants fight dirty. *Nature* 2002; **416**: 267.
- [17] Sharma VP. Health hazards of mosquito repellents and safe alternatives. *Curr Sci* 2001; **80**: 341–343.
- [18] Thacker JRM. *An introduction to arthropod pest control*. Cambridge, UK: Cambridge University Press; 2002, p. 343.
- [19] Isman MB, Miresmailli S, Machial C. Commercial opportunities for pesticides based on plant essential oils in agriculture, industry and consumer products. *Phytochem Rev* 2011; **10**: 197–204.
- [20] Hamideh J, Hesamzadeh Hejazi SM, Babayav MS. Karyotypic studies of three *Thymus* (Lamiaceae) species and populations in Iran. *Caryologia* 2009; **62**: 316–325.
- [21] Saroukolai AT, Moharramipour S, Meshkatalasadat MH. Insecticidal properties of *Thymus persicus* essential oil against *Tribolium castaneum* and *Sitophilus oryzae*. *J Pest Sci* 2010; **83**: 3–8.
- [22] Oshaghi MA, Yaaghoobi F, Vatandoost H, Abaei MR, Akbarzadeh K. *Anopheles stephensi* biological forms, geographical distribution and malaria transmission in malarious regions of Iran. *Pak J Biol Sci* 2006; **9**(2): 294–298.
- [23] Kumari R, Thapar BR, Das Gupta R, Kaul SM, Lal S. Susceptibility status of malaria vectors to insecticides in India. *J Commun Dis* 1998; **30**: 179–185.
- [24] de la Torre Rodriguez YC, Martínez Estrada FR, Flores Suarez AE, Waksman de Torres N, Salazar Aranda R. Larvicidal and cytotoxic activities of extracts from 11 native plants from northeastern Mexico. *J Med Entomol* 2013; **50**(2): 310–313.
- [25] Moharramipour S, Taghizadeh A, Meshkatalasadat MH, Talebi AA, Fathipour Y. Repellent and fumigant toxicity of essential oil from *Thymus persicus* against *Tribolium castaneum* and *Callosobruchus maculatus*. *Commun Agric Appl Biol Sci* 2008; **73**(3): 639–642.
- [26] Park YU, Koo HN, Kim GH. Chemical composition, larvicidal action, and adult repellency of *Thymus magnus* against *Aedes albopictus*. *J Am Mosq Control Assoc* 2012; **28**(3): 192–198.
- [27] Pete UD, Zade CM, Bhosale JD, Tupe SG, Chaudhary PM, Dikundwar AG, et al. Hybrid molecules of carvacrol and benzoyl urea/thiourea with potential applications in agriculture and medicine. *Bioorg Med Chem Lett* 2012; **22**(17): 5550–5554.
- [28] Tang X, Chen S, Wang L. Purification and identification of carvacrol from the root of *Stellera chamaejasme* and research on its insecticidal activity. *Nat Prod Res* 2011; **25**: 320–325.
- [29] Pandey SK, Upadhyay S, Tripathi AK. Insecticidal and repellent activities of thymol from the essential oil of *Trachyspermum ammi* (Linn) Sprague seeds against *Anopheles stephensi*. *Parasitol Res* 2009; **105**(2): 507–512.