



Contents lists available at ScienceDirect

## Asian Pacific Journal of Tropical Biomedicine

journal homepage: www.elsevier.com/locate/apjtb



Document heading

doi: 10.12980/APJTB.4.2014C77

© 2014 by the Asian Pacific Journal of Tropical Biomedicine. All rights reserved.

# Chemistry and bioactivities of essential oils of some *Ocimum* species: an overview

Abhay Kumar Pandey, Pooja Singh\*, Nijendra Nath Tripathi

Bacteriology and Natural Pesticide Laboratory, Department of Botany, DDU Gorakhpur University, Gorakhpur–273009, India

## ARTICLE INFO

## Article history:

Received 23 Nov 2013

Received in revised form 13 May 2014

Accepted 28 Jun 2014

Available online 24 Jul 2014

## Keywords:

*Ocimum* species

Essential oil

Chemistry

Biological properties

## ABSTRACT

Essential oils of different species of the genus *Ocimum* are natural flavouring materials of commercial importance. The data given in current literature are pertaining to the chemical composition of essential oils of different *Ocimum* species viz., *Ocimum basilicum* Linn. (alt. *Ocimum basilicum* var. *minimum*, *Ocimum basilicum* var. *purpurience*), *Ocimum campechianum* Mill., *Ocimum canum* Sims. (*Ocimum americanum*), *Ocimum citriodorum*, *Ocimum gratissimum* Linn., *Ocimum kilimandscharicum* Linn., *Ocimum micranthum* Willd., *Ocimum sanctum* Linn., (alt. *Ocimum tenuiflorum* Linn.), *Ocimum selloi* Benth., *Ocimum trichodon*, *Ocimum utricifolium* from different geographical regions. A considerable difference in chemical composition of a particular species is found, which may be due to their occurrence in different eco-climatic zones and changes in edaphic factors. Attention is also focused on the biological properties of *Ocimum* oils which are related to their various interesting applications as antimicrobial, antioxidant, repellent, insecticidal, larvicidal, nematocidal and therapeutic (anti-inflammatory, antinociceptive, antipyretic, antiulcer, analgesic, anthelmintic, anticarcinogenic, skin permeation enhancer, immunomodulatory, cardio-protective, antilipidemic) agents.

## 1. Introduction

Worldwide demand of essential oils has increased during the past few years. Essential oils are reported in aromatic plants which are distributed in Mediterranean and tropical countries across the world where they are esteemed as an imperative component of the native medicine systems. Almost all plant organs (flowers, buds, stems, leaves, fruits, seeds and roots) of aromatic plants contain essential oils. These are accumulated in secretory cells, cavities, channels, and epidermic cells[1]. Lamiaceae (syn. Labiatae) herb family includes one of the richest essential oil bearing plant family, consists of more than 252 genera and 7000 species in the vicinity of plant kingdom[2]. Lamiaceae family is known for the wealth of species with medicinal properties, which have been used since early times and many of these species are common in Mediterranean region. Many species of

Lamiaceae have long history of uses in culinary spices and folk medicine. For example, oregano, rosemary, sage and thyme are typical seasonings in the Mediterranean region, and especially oregano is consumed in larger quantities all around the world as part of pizza seasoning mix[3]. Although there have been toxic components described for some Lamiaceae plants, the importance of many family members to the culinary and essential oil industries has been commercially explored for more than 75 years[4–6]. Essential oils recovered from the Lamiaceae species have been used against different diseases like intestinal disorder and bronchitis[7].

The genus *Ocimum*, comprising of more than 150 species grows widely and is distributed throughout temperate regions of the world[6,8,9]. *Ocimum basilicum* (*O. basilicum*), *Ocimum gratissimum* (*O. gratissimum*) and *Ocimum sanctum* (*O. sanctum*), commonly known as holy basil, clove basil (wild basil/East India basil) and sweet basil, respectively, are frequently cultivated in several countries of East Asia, Europe, America and Australia for the production of essential oils[10,11]. *Ocimum americanum* (*O. americanum*) formerly known as *Ocimum canum* (*O. canum*), includes wild species in India, but is cultivated

\*Corresponding author: Dr. Pooja Singh, Assistant Professor, Bacteriology & Natural Pesticide Lab, Department of Botany, DDU Gorakhpur University, Gorakhpur–273009, India.

E-mail: pooja.ddu@gmail.com

Tel: +919415083883

Foundation Project: Supported by CST UP Lucknow (Grant No. CST/AAS, D–09, April 3, 2007).

in Indonesia for its essential oil for commercial purposes. *Ocimum kilimandscharicum* (*O. kilimandscharicum*), a variety species of *O. basilicum*, commonly called African blue basil is known for its camphor like scent of its essential oil. Similarly, *Ocimum minimum* and *Ocimum citriodorum* also include variety species of *O. basilicum* popular in Indonesia, Mexico and Africa for their naturally occurring essential oils as in perfumery and cosmetic applications. Likewise, *Ocimum tenuiflorum* (*O. tenuiflorum*) includes variety species of *O. sanctum*. Traditionally, these *Ocimum* species have been extensively utilized in food and perfumery industries<sup>[12]</sup>. The aerial parts of the plants are considered as antispasmodic, stomachic and carminative in native medicine<sup>[13]</sup>.

A high degree of polymorphism in the genus *Ocimum* determines a large number of subspecies, different varieties and forms producing essential oils with varying chemical composition offering variable level of medicinal potential<sup>[14]</sup>. Essential oils extracted from *Ocimum* plants have been reported to possess interesting biological properties. These volatile oils have been applied in perfumery, to inhibit growth of microorganisms, in food preservation and in aromatherapy. The potential uses of *O. basilicum*, *O. canum*, *O. gratissimum* and *O. sanctum* essential oils, particularly as antioxidant and antimicrobial agents have also been explored<sup>[15–18]</sup>. Recently, Mondal *et al.* reviewed the antimicrobial, adaptogenic, antidiabetic, hepato-protective, anti-inflammatory, anti-carcinogenic, radioprotective, immunomodulatory, neuro-protective, cardio-protective and mosquito repellent properties of *O. sanctum*<sup>[19]</sup>. In order to investigate the structural activity relationship, it is necessary to know the chemical differentiation among volatiles of various *Ocimum* species. Keeping this in view we have reviewed here the chemistry of various samples of *Ocimum* oils from different geographical regions. Further, the objective of our ongoing investigation is to establish biological evaluation of oils as potential source for beneficial uses.

## 2. Chemical investigation on essential oils

Researches on chemical composition of most of the essential oils were undertaken by various researchers using GC and GC-MS techniques. The other techniques used are Kovat's index from packed capillary column, retention time data from GC, HPLC, GC-MS, capillary GC, HRGG-FID, HRGG-MS and IR, C13, NMR. The constituents were identified by calculation of their retention indices under temperature programmed condition for *n*-alkanes (C8–C20). Identification of individual component was assigned by retention time comparison with authentic components and oil of known composition and by mass spectra with those obtained from Wiley/NIST/Pfleger library spectra as well as with literature data.

## 3. Chemical composition

Essential oils are very complex natural mixtures which contain about 20–60 components at quite different concentrations. They are characterized by two or three major components at fairly high concentration (20%–70%) as compared to other components present in trace amount. Essential oils are complex mixtures of natural organic compounds which are predominantly composed of terpenic hydrocarbons (myrcene, pinene, terpinene, limonene, *p*-cymene,  $\alpha$ - and  $\beta$ - phellandrene) and terpenoids (oxygen containing hydrocarbons) like acyclic monoterpene alcohols (geraniol, linalool), monocyclic alcohols (menthol, 4-carvomenthol, terpineol, carveol, borneol), aliphatic aldehydes (citral, citronellal, perillaldehyde), aromatic phenols (carvacrol, thymol, safrol, eugenol), bicyclic alcohol (verbenol), monocyclic ketones (menthone, pulegone, carvone), bicyclic monoterpene ketones (thujone, verbenone, fenchone), acids (citronellic acid, cinnamic acid) and esters (linalyl acetate)<sup>[20]</sup>. Mono- and sesquiterpenoidal essential oil constituents are formed by the condensation of isopentenyl pyrophosphate units. Diterpenes usually do not occur in essential oils but are sometimes encountered as by-products.

Lawrence classified the basil oils into three large groups' European type, exotic or reunion type and African type according to their chemical composition and geographical origin<sup>[21]</sup>. He established four essential oil chemotypes (methyl chavicol, linalool, methyl eugenol and methyl cinnamate) and also numerous subtypes of oils extracted from *O. basilicum*<sup>[22]</sup>. According to the biosynthetic origin of major compounds, he classified them as chemotypes with single or dual biosynthetic pathways. From time to time, other chemotypes *viz.*, eugenol-rich and thymol-rich in *O. gratissimum*<sup>[23]</sup>, sesquiterpenes-rich in *O. canum* and *O. sanctum* and terpinene-4-ol-rich<sup>[22,24]</sup> have also been reported from different places. The *Ocimum* species which are reviewed here for their essential oil chemistry, have been collected from the different countries and regions of the world such as Bangladesh, Brazil, Cameroon, Egypt, Europe, Guinea, India, Iran, Italy, Mali, Nigeria, Pakistan, Rwanda, Thailand, Togo, Turkey, Um Ruaba and Yaounde. Literature survey revealed that chemical constituents and their respective percentage of several *Ocimum* species varied depending upon the origins and cultivars (Table 1).

It can be described from Table 1 that the major constituents which have been isolated from different *Ocimum* oils include 1,8-cineol, linalool, pinene, eugenol, camphor, methyl chavicol, ocimene, terpinene, limonene *etc.* Table 1 also demonstrates that within the *Ocimum* species, there is a clear variation in their composition in terms of type of constituents and their composition. The chemical structures of these compounds are depicted in Figure 1.

#### 4. Biological properties of *Ocimum* oils

Currently, the interests in natural products are focused on sources of alternative, more natural and environmentally friendly antimicrobials, pesticides and therapeutic activities. The possibility of utilizing volatile oils is now being investigated. Generally the action of volatile oils is the result of the combined effect of both their active and inactive compounds. The inactive compounds might influence resorption, rate of reactions and bioavailability of the active compounds. Several active components might have a synergistic effect. Biological activity of an essential

oil is related to its chemical composition. The relation between composition and bioactivity of the essence from the aromatic plants may be attributable both to their major components (alcoholic, phenolic, terpenic or ketonic compounds) and the minor ones present in the oil. It may act together synergistically or antagonistically to contribute to some activity of the tested oil. Experiments on biological activities of essential oils from different *Ocimum* species have been established by numerous researchers. In the present review, we have described the some important bio-properties of different essential oils of *Ocimum* species from different parts of the world.


**Table 1**

Investigation on chemical composition of essential oils of different *Ocimum* species.

Plant species	Major constituents (%)	Investigator	Place
<i>O. basilicum</i> L.	Methyl chavicol (70.00), Linalool (25.00), Eugenol (5.00)	Mondawi <i>et al.</i> [118]	Um Ruaba
<i>O. gratissimum</i> L. 387E	Thymol (35.40), p-Cymene (18.30), Eugenol (10.70)	Ntezuriubanza <i>et al.</i> [119]	Southern Western Rwanda
<i>O. gratissimum</i> 345C	Thymol (46.60), $\gamma$ -Terpinene (22.40)		
<i>O. basilicum</i>	Linalool (45.70), Eugenol (13.40), Methyl eugenol (9.57), Fenchyl alcohol (3.64)	Akgul[120]	Turkey
<i>O. canum</i> Sims. 337C	Linalool (82.30)		
<i>O. gratissimum</i> 345C	Thymol (46.00–7.00)		
<i>O. gratissimum</i> 387E	Thymol (35.40), Eugenol (10.70)		
<i>Ocimum trichodon</i> 213C	Eugenol (74.70)		
<i>Ocimum utricifolium</i> 243C	Trans-Methylisoeugenol (49.70), Trans- $\beta$ -Ocimene (13.50), Cis- $\beta$ -ocimene (12.60)		
<i>Ocimum utricifolium</i> 298E	Methyleugenol (73.60)	Janseen <i>et al.</i> [26]	Rwanda
<i>Ocimum utricifolium</i> 364A	Trans-Methylisoeugenol (35.60), Cis- $\beta$ -Ocimene (29.20)		
<i>Ocimum utricifolium</i> 375I	Eugenol (39.40), Cis- $\beta$ -ocimene (41.00)		
<i>Ocimum utricifolium</i> 378I	Eugenol (37.60), Cis- $\beta$ -ocimene (39.00)		
<i>Ocimum utricifolium</i> 392G	Methyleugenol (86.60)		
<i>Ocimum utricifolium</i> 401C	Eugenol (71.00)		
<i>Ocimum utricifolium</i> 402C	Eugenol (71.00)		
<i>O. basilicum</i>	Methyl chavicol (87.30), Linalool (5.40), Methyl eugenol (1.50), $\beta$ -Caryophyllene (2.40), $\alpha$ -Pinene (1.00), $\beta$ -Pinene (0.80), Limonene (0.50), Camphene (0.20)	Khatri <i>et al.</i> [121]	Pakistan
<i>O. basilicum</i>	Linalool, Methyl chavicol, Eugenol	Marotti <i>et al.</i> [122]	Italy
<i>O. basilicum</i>	Limonene (10.40), Linalool (50.80)	Zallo <i>et al.</i> [28]	Western provinces of Cameroon
<i>O. canum</i>	1,8-Cineole (78.30), $\beta$ -Pinene (5.80)		
<i>O. gratissimum</i>	p-Cymene (7.00), $\gamma$ -Terpinene (20.00), Thymol (46.20)		
<i>O. basilicum</i>	Linalool, Methyl eugenol	Chalchat <i>et al.</i> [123]	Mali
<i>O. basilicum</i>	Linalool (69.00), Eugenol (10.00), (E)- $\alpha$ -Bergamotene (3.00), Thymol (2.00)	Keita <i>et al.</i> [124]	Guinea
<i>O. americanum</i> L. citral type	Geranial (37.70), Neral (27.90)	Mondello <i>et al.</i> [125]	Bangladesh
<i>O. americanum</i> L. camphor type	Camphor (38.60), Limonene (10.60)		
<i>O. basilicum</i> var. <i>purpurascens</i>	Linalool (29.70), Geranial (27.40), Geranyl acetate (13.80)		
<i>O. sanctum</i> L. green	Eugenol (41.70), Limonene (3.80), E. Caryophyllene (24.40)		
<i>O. sanctum</i> purple	Eugenol (77.50), E-Caryophyllene (10.70)		
<i>O. basilicum</i>	1,8-Cineole (11.00), Linalool (42.50), Estragole (52.20)		
<i>O. basilicum</i> var. <i>minimum</i>	Linalool (16.00), Estragole (52.20), 1,8-Cineole (7.40)	Silva <i>et al.</i> [126]	North Eastern Brazil
<i>O. basilicum</i> var. <i>purpurascens</i>	Linalool (39.30), $\alpha$ -Muurolol (11.00)		
<i>O. basilicum</i>	Methyl eugenol (78.02), $\alpha$ -Cubebene (6.17), Nerol (0.83), $\epsilon$ -Muurolene (0.74)	Ozcan and Chalchat[127]	Turkey
<i>Ocimum minimum</i>	Geranyl acetate (69.48), Terpinen-4-ol (2.35), Octan-3-yl-acetate (0.72)		

**Table 1, continued**Investigation on chemical composition of essential oils of different *Ocimum* species.

Plant species	Major constituents (%)	Investigator	Place
<i>O. gratissimum</i> L.	Eugenol (54.00), 1,8–Cineole (21.60)	Silva <i>et al.</i> [10]	Brazil
<i>O. micranthum</i> Willd.	Eugenol (64.80), (E)– $\beta$ –caryophyllene (14.30), Bicyclogermacrene (8.10)		
<i>Ocimum selloi</i> Benth.	Anethole (64.60), Linalool (20.60)		
		Tchoumboungang	
<i>O. gratissimum</i>	$\beta$ –Phellandrene (21.10), Limonene (11.40), $\gamma$ –Terpinene (21.90), Thymol (11.20)	<i>et al.</i> [80]	Yaounde
<i>O. basilicum</i>	Linalool (10.00), Methyl chavicol (60.30), Methyl cinnamate (6.30)	Kasali <i>et al.</i> [128]	Nigeria
<i>O. basilicum</i>	Linalool (44.18), 1,8–Cineole (13.65), Eugenol (8.59)	Ismail[107]	Egypt
<i>O. basilicum</i> cv. purple	Methyl chavicol (52.40), Linalool (20.10), Epi– $\alpha$ –cadinol (5.90), Trans– $\alpha$ –bergamotene (5.20)	Sajjadi[13]	Isfahan, Iran
<i>O. basilicum</i> cv. green	Methyl chavicol (40.50), Geranial (27.60), Neral (18.50), Caryophyllene oxide (5.40)		
<i>O. sanctum</i> Shyama	Methyl eugenol (67.80), E–Caryophyllene (17.10)	Awasthi and Dixit [129]	Northern India
<i>O. sanctum</i> Rama	Eugenol (46.20), E–Caryophyllene (27.60), $\beta$ –Elemene (16.30)		
<i>O. basilicum</i>	$\beta$ –Ocimene (2.27), Methyl chavicol (92.48)	Bunrathep <i>et al.</i> [55]	Pathumthani (Thailand)
<i>O. canum</i>	Linalool (5.65), Neral (34.01), Geranial (37.89), $\beta$ –Caryophyllene (6.79)		
<i>O. gratissimum</i>	Z– $\beta$ –ocimene (48.28), $\beta$ –Caryophyllene (3.06), $\gamma$ –Muurolene (9.32), $\alpha$ –Farnasene (4.33), Eugenol (25.02)		
<i>O. sanctum</i>	$\beta$ –Caryophyllene (35.20), Methyl eugenol (53.67)		
<i>O. basilicum</i>	1,8–Cineole (4.00), Linalool (28.60), Estragole (21.70), (E)–Methyl Cinnamate (14.30)	Politeo <i>et al.</i> [17]	Austria
<i>Ocimum campechianum</i> Mill.	Eugenol (32.20–60.60), Methyl eugenol (60.60–69.500), 1,8–Cineole (0.90–19.70), Elemicin (0.20–65.90)	Zoghbi <i>et al.</i> [130]	North Brazil
<i>O. gratissimum</i> (5 samples)	Thymol (13.10–36.20), $\gamma$ –Terpinene (0.20–28.10), 1,8–Cineole (0.00–25.20), p–Cymene (4.40–19.90)		
<i>O. basilicum</i>	Linalool (53.00), Methyl chavicol (29.00)	Zheljazkov <i>et al.</i> [11]	European basil
<i>O. basilicum</i>	Estragole (85.50), Linalool (1.71)	Koba <i>et al.</i> [43]	Lome (Togo)
<i>O. basilicum</i>	Methyl chevicol (70.04), Linalyl acetate (22.54) (13.56), Camphene (7.32)	Anand <i>et al.</i> [47]	Uttarakhand, India
<i>O. kilimandscharicum</i>	Camphor (56.07), DL–Limonene		
<i>O. gratissimum</i>	Eugenol (53.89), Cis–ocimene (23.97), Germacrene–D (10.36)		
<i>O. gratissimum</i>	Eugenol (63.70), (Z)– $\beta$ –ocimene (19.60), Germacrene–D (7.30)	Verma <i>et al.</i> [48]	Sub–tropical region of northern India
<i>O. kilimandscharicum</i>	Camphor (63.40), Limonene (7.90), Camphene (5.80), $\gamma$ –Terpinene (4.70)		
<i>O. basilicum</i> var. purpurascens	Methyl cinnamate (59.95), Linalool (16.40), tau–Cadinol (4.37)	Mohhiuddin <i>et al.</i> [131]	Bangladesh
<i>O. basilicum</i>	Methyl chavicol (62.00), Linalool (24.00)	Sastry <i>et al.</i> [132]	Deccan Region, India
<i>O. gratissimum</i>	Eugenol (74.80), Limonene 1,8–Cineole (10.80)		
<i>O. tenuiflorum</i>	Eugenol (84.00), $\beta$ –Caryophyllene (6.90)		

**Figure 1.** Chemical structure of some foremost compounds isolated from essential oils of *Ocimum* species.

#### 4.1. Antimicrobial activity

The indiscriminate use of antimicrobial agents has resulted in the emergence of a number of drug-resistant bacteria and fungi. To overcome the increasing resistance of pathogenic microbes, more effective antimicrobial agents with novel mode of action must be developed. Essential oils derived from several *Ocimum* species have been reported to be active against several Gram-positive and Gram-negative bacteria as well as against yeasts and fungi due to their terpenic constituents. Recently, essential oils and extracts of certain plants have been shown to have antimicrobial effects[25] as well as imparting flavour to foods[1]. Janssen *et al.* investigated the antimicrobial potential of four *Ocimum* species viz., *O. canum*, *O. gratissimum*, *Ocimum trichodon* and *O. urticifolium* grown in Rwanda against *Escherichia coli* (*E. coli*), *Bacillus subtilis* (*B. subtilis*), *Staphylococcus*



*aureus* (*S. aureus*) and *Trichophyton mentagrophytes* var. *interdigitale*[26]. They found that *O. canum* was the most effective amongst the studied oils. Essential oil extracted from the leaves of *O. sanctum* has been found to inhibit *in vitro* growth of *E. coli*, *B. anthracis* and *Pseudomonas aeruginosa* (*P. aeruginosa*) which showed its antibacterial activity[27].

As an example of antimycotic activity, the oil of *O. gratissimum* leaves was fungicidal at 78 mg/L for *Microsporium gypseum* and *Trichophyton rubrum*, but a concentration of 312 mg/L was required to inhibit growth of *Candida albicans* (*C. albicans*) and *Cryptococcus neoformans*[28]. Further, Basilico and Basilico reported the antifungal activity of *O. basilicum* oil against *Aspergillus ochraceus* and found that oil was effective at 500 mg/L[29]. At 500 µg/L, *O. canum* oil exhibited 100% mycelial inhibition against *Penicillium italicum*, post-harvest rotting fungus of *Citrus* fruits[30]. Moreira *et al.* worked on antimicrobial activity of *O. basilicum* on survival and growth of different strains of *E. coli* O157:H7[31]. The strains of *E. coli* exhibited similar susceptibilities to the action of the essential oil assayed with the minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) values were 1.9 mL/100mL and 2.0 mL/100 mL, respectively. Mbata and Saikia investigated the antibacterial effect of *O. gratissimum* oil against *Listeria monocytogenes* serotype and reported that at 20–250 µg/mL, the essential oil progressively inhibited the bacterial growth[32]. Similar antimicrobial activity is reported in essential oils isolated from Amazonian basil, *O. gratissimum* and *Ocimum micranthum* (*O. micranthum*)[33,34]. Weseler *et al.* found that MIC value of *O. basilicum* against *Helicobacter pylori* was of 286.7 µg/mL while MBC value increased upto 573.4 µg/mL[35]. Wannissorn *et al.* by using disc diffusion assay, evaluated the antimicrobial activity of *O. basilicum*, *O. basilicum* var. *citratum*, *O. gratissimum*, *O. tenuiflorum* essential oils from Thailand against zoonotic enteropathogens including *Salmonella* spp., *E. coli* O157, *Campylobacter jejunii* and *Clostridium perferingens* which are important for broiler export[36]. The zone of inhibition ranges of 9–80 mm with *O. basilicum* var. *citratum* was the most potent. The antifungal activity of eugenol isolated from *O. gratissimum* reported against *Alternaria* and *Penicillium chrysogenum*[37].

Bozin *et al.* reported the antibacterial activity of *O. basilicum* against 13 human pathogenic bacteria and find that the MIC ranges of 8–30 µL/mL[15]. Atanda *et al.* investigated the ability of sweet basil (*O. basilicum*) in the growth control of aflatoxigenic fungus *Aspergillus parasiticus*[38]. Lee *et al.* did the work on inhibitory effect of *O. basilicum* oil and its main constituent eugenol and linalool on *Microsporium gypseum* (guinea pig infection) at MIC of 0.01%–0.03%[39]. Similarly, *O. basilicum* inhibited the growth of *S. enteritidis* at MIC=20.0–80.0 µg/mL[40], *O. forskolei* Benth. oil of *B. subtilis* and *S. aureus* by diffusion test were at MIC 32 µg/mL and 128 µg/mL, respectively, and *O. sanctum*

of *Alternaria alternata*, *Aspergillus* species, *Penicillium* species, *Fusarium nivale* were at MIC 0.3 µL/mL[41,42]. The *in vitro* microbiological experiments of Koba *et al.* revealed that only the methyl eugenol and methyl eugenol/t-anethole chemotypes were active against tested fungi and bacteria[43]. Their MIC against fungi ranged from 80–150 µL/L and from 200–500 µL/L respectively. Likewise, on tested bacteria, the MIC varied from 200–400 µL/L and from 250–500 µL/L respectively. In this work, there is no doubt that the antifungal activity of *O. basilicum* chemotype methyleugenol against tested fungi is a predictable consequence of its high content in methyleugenol known as one of the phenolic volatile molecules endowed with antimicrobial properties[44]. Pandey *et al.* did work on the antimicrobial activity of *O. sanctum* against seven bacteria (*Bacillus pumilus*, *B. subtilis*, *S. aureus*, *E. coli*, *Klebsiella pneumoniae*, *P. aeruginosa* and *Salmonella typhi*), two filamentous fungi (*Aspergillus niger* and *Aspergillus flavus*) and a yeast (*C. albicans*)[45]. The oil inhibited 100% microbial growth at 700 mg/L for bacteria and 1300 mg/L for fungi. Recently, Singh *et al.* investigated that essential oil of *O. canum* strongly inhibited the mycobial growth of seven post-harvest pathogenic fungal species (*Aspergillus flavus*, *Aspergillus nidulans*, *Aspergillus niger*, *Aspergillus ochraceus*, *Mucor* sp., *Penicillium italicum*, *Penicillium oxalicum* and *Rhizopus arrhizus*) and found that oil was toxic at 500 mg/L against all the fungi[46]. Anand *et al.* look for the antimicrobial activity of essential oils of three *Ocimum* species *viz.*, *O. basilicum*, *O. kilimandscharicum* and *O. gratissimum* against Gram-positive (*S. aureus*, *Enterococcus faecalis*) and Gram-negative (*E. coli*, *P. aeruginosa*) bacteria as well as yeast *C. albicans*[47]. Report revealed that *O. gratissimum* oil was the most effective with MIC values ranging from 1.5–12.5 mg/mL. Verma *et al.* evaluated the antibacterial activity of *O. gratissimum* and *O. kilimandscharicum* against *S. aureus*, *Streptococcus mutans*, *Enterococcus faecalis* and found that *O. gratissimum* oil was more toxic against all tested bacterial strains and showed a zone of inhibition ranging from 8–16 mm[48]. Pandey *et al.* investigated that *O. sanctum* oil was found to be toxic against two plant pathogenic bacteria *Erwinia herbicola* and *Pseudomonas putida* by exhibiting complete inhibition of their microbial growth[49]. Oil exhibited the MIC 2.0 µL/mL for *Erwinia herbicola* and 1.0 µL/mL for *Pseudomonas putida*, but was bactericidal at higher doses of 8 and 4 µL/mL respectively.

#### 4.2. Antioxidant activity

Antioxidants are substances that inhibit the oxidation of our cells from toxins such as free radicals. The toxins can be from the natural digestion and metabolism of foods, alcohol, nicotine from cigarette smoke, environmental factors, OTC drugs such as acetaminophen, (Tylenol®) prescription drugs, preservatives, *etc.* Reactive oxygen species (ROS) including singlet oxygen (<sup>1</sup>O<sub>2</sub>), superoxide ion (O<sup>2-</sup>), hydroxyl ion (OH<sup>-</sup>),

and hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) are highly reactive and toxic molecules generated in cells during normal metabolism. However, in response to a variety of factors including tobacco smoke, pollutants, ionizing radiations, alcohol, synthetic pesticides, and solvent, their production increases. ROS can cause oxidative damage to proteins, lipids, enzymes, and DNA, and they have also been linked to pathogenesis of oxidative diseases. Living cells possess an excellent scavenging mechanism to avoid excess ROS induced cellular injury, however, with ageing and under influence of external stresses, these mechanisms become inefficient, and dietary supplementation of synthetic antioxidants is required[50]. In this context, aromatic plants, particularly their essential oils, are being evaluated for antioxidant activity. *Ocimum* plants contain large amounts of antioxidants other than vitamin C, vitamin E, flavonoids and carotenoids. The presence of many pharmacologically active compounds in *Ocimum* species provides them protection against free radical induced oxidative damage of cellular components. The antioxidant capacity of basil essential oils has been studied several times[51–54]. Bunarathep *et al.* reported the comparative antioxidant activity of essential oils of four *Ocimum* species (*O. basilicum*, *O. canum*, *O. gratissimum* and *O. sanctum*) by DPPH bioassay and found that *O. gratissimum* was the most antioxidative followed by *O. sanctum*, *O. canum* and *O. basilicum* with EC<sub>50</sub> values 30.20, 767.82, 8343.19 and 47057.45 µg/mL respectively[55]. Politeo *et al.* pointed out the antioxidant capacity of free volatile aglycons from basil (*O. basilicum*) by two different methods which were 2,2-diphenyl-1-picrylhydrazyl radical scavenging method (DPPH) and ferric reducing/antioxidant power assay (FRAP)[17]. DPPH method shows that free volatile aglycones possess good antioxidant properties comparable with that of the essential oil and well-known antioxidant butylated hydroxytoluene, but less than pure eugenol. The results obtained by FRAP method showed that these compounds are somewhat less effective antioxidants than essential oil and butylated hydroxytoluene. From time to time, potential uses of *O. sanctum*, *O. gratissimum* and *O. basilicum* essential oils as antioxidant agent have been explored[15,16,18]. Trevisan *et al.* evaluated the antioxidant capacity of essential oils obtained by steam hydrodistillation from five species of the genus *Ocimum*, namely *O. basilicum* var. *purpurascens*, *O. basilicum*, *O. gratissimum*, *O. micranthum* and *O. tenuiflorum* (syn. *O. sanctum*), using a high-performance liquid chromatography-based hypoxanthine/xanthine oxidase and the DPPH assays[56]. In the hypoxanthine/xanthine oxidase assay, strong antioxidant capacity was evident in all the oils, but the greater was shown by that obtained from *O. tenuiflorum* (syn. *O. sanctum*) (IC<sub>50</sub>=0.46 µL/mL) compared to *O. basilicum* var. *purpurascens* (IC<sub>50</sub>=1.84 µL/mL). In the recent research, *O. sanctum* and Omani basil oils were also found to possess significant antioxidant action[57,58].

### 4.3. Repellent, insecticidal, larvicidal and nematocidal activities

Plant based fumigants have long been touted as attractive alternatives to synthetic fumigants for arthropods management because botanicals reputedly pose little threat to the environment or to human health. Genus *Ocimum* is known for pesticidal properties due to diverse group of compounds in its essential oil. Basil oil contains bioactive constituents that are insecticidal and repellent[59–62]. Esther *et al.* observed that oil extracted from the leaves of tropical shrub, *O. suave* cause mortality in all stages of the tick *Rhipicephalus appendiculatus*[63]. A 10% solution was found to kill all immature and more than 70% of adults feeding on rabbits. The toxicity of the result in this study may be due to the toxic effects of the compounds eugenol, mono and sesquiterpenoids found in the plant extract[64]. Sosan *et al.* studied on larvicidal activity of *O. gratissimum* on larvae of *Aedes aegypti* L. (*Ae. aegypti*) and found that oil exhibited 100% mortality at 300 mg/L concentration of 24 h exposure[65]. However, a related species *O. canum* growing in Zimbabwe was reported to exhibit repellency against adults of *Ae. aegypti*[66]. Aslan *et al.* tested essential oil from *O. basilicum* for their toxicity against the nymphs and adults of *Tetranychus urticae* (Acari: Tetranychidae) and adults of *Bemisia tabaci* (Aleyrodidae)[67]. Cavalcanti *et al.* determined the larvicidal activity of essential oils from Brazilian plants *O. americanum* and *O. gratissimum* against *Ae. aegypti*[68]. The results showed that *O. americanum* and *O. gratissimum* have LC<sub>50</sub> of 67 and 60 mg/L respectively. Ilondu *et al.* observed that *O. suave* (wild basil) caused mortality of adult *Rhyzopertha dominica* in grain cereals at the 2–8 g plant materials per 50 g cereals (rice, maize and sorghum)[69]. This protects the cereals after 90 d post treatment. Olonisakin *et al.* reported the insecticidal activity of *O. shave* against *Callosobruchus maculatus* with LC<sub>50</sub> values of 0.66 mg/10 g seed[70]. Lopez *et al.* found that estragole is example of toxic fumigant compound in the basil essential oil (*O. basilicum*) that is active against insect pests[71]. Bakr *et al.* evaluated the insecticidal effect of volatile oil from basil (*O. basilicum*) against two museum insect pests black carpet beetle (*Attagenus fasciatus*, a pest in the museums where it attacks) and cigarette beetle (*Lasioderma serricorne*, a pest of stored tobacco), LC<sub>50</sub> values of oil against larvae and adults of the black carpet beetle were 1.78786 and 2.05647 mL/80 cm<sup>3</sup> as well as 1.6609 mL/80 cm<sup>3</sup> and 2.01459 mL/80 cm<sup>3</sup> for the larvae and adults of cigarette beetle, respectively[72]. Manzoor *et al.* reported the *O. sanctum* oil as toxicant and repellent agent against termite, *Heterotermes indicola*[73]. Similarly, Basil oil (*O. basilicum*) and its major constituent's trans-anethole, estragole, linalool were insecticidal to adult fruit flies of *Ceratitis capitata* (*C. capitata*), *Bactrocera dorsalis* (*B. dorsalis*), and *Bactrocera cucurbitae* (*B. cucurbitae*)[74]. The toxic action of basil oil in *C. capitata* occurred significantly

faster than that of in *B. cucurbitae*, but slightly faster than that of in *B. dorsalis*. Estragole acted faster in *B. dorsalis* than that of in *C. capitata* and *B. cucurbitae*. Linalool action was faster in *B. dorsalis* and *C. capitata* than that of in *B. cucurbitae*. Trans-anethole action was similar to all the three species. Methyl eugenol acted faster in *C. capitata* and *B. cucurbitae* than that of in *B. dorsalis*. When linalool was mixed with cuelure (attractant to *B. cucurbitae* male), its potency to the three species decreased as the concentration of cuelure increased. Oil of *O. suave* (wild basil) was used under ambient conditions with the aim of evaluating the toxicity of the plant oil on adult housefly—*Musca domestica*[75]. The oil applied at 0.05, 0.10, 0.15 and 0.20 mL/L caused significant mortality of the housefly. The LC<sub>50</sub> was 0.09 mL/50 mL of water, and the LT<sub>50</sub> was 4.40 h. This result shows that *O. suave* leaf oil could be used as an alternative to synthetic insecticides for the control of *M. domestica*. More recently, Pandey and Tripathi reported that the dose of 5 and 10 µL *O. sanctum* oil was responsible for absolute repellent and insecticidal activities against pulse beetles *Cryptocarya chinensis* and *Callosobruchus maculatus* respectively[76]. Rajamma *et al.* examined the comparative larvicidal activity of different species of *Ocimum* (*O. sanctum*, *O. basilicum* and *O. gratissimum*) against larvae of *Culex quinquefasciatus* in terms of LD<sub>50</sub> value on late third or early fourth instar larvae for a period of 24 h[77]. A comparison of LD<sub>50</sub> value has shown that *O. basilicum* oil is more active than the other two species. The LD<sub>50</sub> value of *O. basilicum* and *O. sanctum* oils were 39.31 and 40.02 µL/L on the laboratory reared larvae and 129.53 and 139.49 µL/L on field collected larvae. Laboratory reared larvae were more sensitive than field collected larvae.

Bioassay tests were conducted to find out the nematicidal activity of *O. basilicum* oil against *Meloidogyne inconita*[78]. The oil was highly toxic at 250 mg/L. Holetz *et al.* studied on effect of essential oil of *O. gratissimum* on the trypanosomatid *Herpetomonas samuelpeaoi*[79]. They observed that at from 20–250 µg/mL, the essential oil progressively inhibited the protozoan growth. The IC<sub>50</sub> in defined and complex media, at 28 °C were 100 and 91 µg/mL, respectively. Tchoumboungang *et al.* investigated the antimalarial activity of *O. gratissimum* essential oil on mice infected with *Plasmodium berghei*[80]. The oil showed maximum 77.5% activity at 500 mg/kg of mouse per day. Oil also showed significant suppression of parasitaemia (55%–77.8%) at oral dose ranges of 200–500 mg/kg per day. Santoro *et al.* worked out on *O. basilicum* oil and its active constituent eugenol, linalool and observed that they were strong inhibitor of *Trypanosoma cruzi* which causes epimastigotes, trypomastigotes with LC<sub>50</sub> values ranges of 99.5–57.5 lg/mL[81].

#### 4.4. Therapeutic properties of *Ocimum* oils

A variety of curative properties have been recognized to *Ocimum* species. The medicinal plants are rich in secondary

metabolites (which are potential sources of drugs) and essential oils of therapeutic importance. The important advantages claimed for therapeutic uses of medicinal plants in various ailments are their safety besides being economical, effective and easy available. Several studies have shown that steam distilled essential oils extracted from fresh leaves of *Ocimum* species have therapeutic importance.

##### 4.4.1. Anti-inflammatory, antinociceptive, antipyretic, antiulcer, analgesic and anthelmintic properties

Essential oil (200 mg/kg body weight) of fresh leaves and fixed oil (0.1 mL/kg, body weight) of seeds (fixed oil) of *Ocimum* species have proved as anti-inflammatory agent on experimental animal's hind paw edema induced by carrageenan, serotonin, histamine and prostaglandin-E-2, significantly reduce the edema when compared with the saline treated control[82]. The oil of *O. sanctum* was found to possess significant anti-inflammatory activity against carrageenan and different other mediator-induced paw edema in rats[83]. This result supports the dual inhibition of arachidonate metabolism and indicated by its activity in inflammation models that are intensive to selective cyclooxygenase inhibitors. The mechanism of action of the anti-inflammatory effects of Tulsi could be the cyclo-oxygenase and lipooxygenase pathways[83,84]. In earlier work, oil of *O. sanctum* was found to inhibit the inflammation induced by carrageenan[85]. The oil also possesses significant antipyretic, analgesic and antiarthritis activity without any noticeable toxicity[86]. Sahouo *et al.* reported inhibitory effect produced by chemical constituents of *O. gratissimum* essential oil as anti-inflammatory and analgesic drugs, *in vitro*, on soybean lipoxygenase L-1 and cyclooxygenase function of prostaglandin H synthase (PGHS), the two enzymes involved in the production of mediators of inflammation[87]. *O. gratissimum* inhibited the two enzymes, cyclooxygenase function of PGHS and lipoxygenase L-1, with an IC<sub>50</sub> values of 125 and 144 µg/mL respectively. The proteins implicated in production of mediators of inflammation, proposed in literature, were PGHS and lipoxygenase-5LO. Their natural substrate is arachidonic acid. Indeed, it is well established that PGHS catalyzed biotransformation of arachidonic acid into prostaglandins, thromboxanes and prostacyclins which are mediators of inflammation, hormonal modulation and platelet aggregation. Lipoxygenase 5LO catalyzed biotransformation of arachidonic acid into fatty acid hydroperoxides, lipoxines, and leukotrienes, compounds implicated in allergy and suffering. Lipoxygenase L-1 was a model of lipoxygenase 5LO and its natural substrate was linoleic acid[88]. Singh *et al.* showed that *O. sanctum* oil produced hypotensive effect in anaesthetized dog, which seems to be due to its peripheral vasodilatory action. The oil increased blood-clotting time and percentage increases were comparable to aspirin and could be due to inhibition of platelet aggregation[89]. The

oil also showed inhibition towards cytochromatic enzyme responsible for hepatic metabolism of pentobarbitone. Both eugenol and *O. gratissimum* oil presented anthelmintic activity against *Haemonchus contortus*, the main nematode of ovines and caprines in Northeastern Brazil[90].

The essential oil of *O. micranthum* was studied for a possible analgesic effect on the acetic acid induced writhing and formalin test in mice and antioedema activities on the carrageenan and dextran induced paw edema in rats. The essential oil demonstrated antinociceptive effects, and pretreatment with naloxone did not reverse the antinociception, indicating that the opioid system is not involved. On the other hand, pretreatment with L-arginine reversed the antinociception, suggesting involvement of the nitric oxide system. The essential oil did not present an antioedematogenic effect in the carrageenan and dextran models[91]. Rabelo *et al.* found that *O. gratissimum* oil produced a dose-dependent inhibition of acetic acid-induced writhing, causing up to a ~60% inhibition at the highest dose (300 mg/kg) used during their experiment antinociceptive effects of *O. gratissimum* essential oil in mice[92]. The therapeutic use of *O. sanctum* in treatment of gastric ulcer has been attributed to antiulcerogenic action of eugenol and essential oil extracted from Tulsi leaves[93,94].

#### 4.4.2. Anti-cancerous property

Estragole, a constituent of *O. basilicum* essential oil, has shown carcinogenic properties in rat and mouse[95,96]. Prakash and Gupta investigated the chemoprotective activity of *O. sanctum* oil against subcutaneously injected 20-methylcholanthrene induced fibrosarcoma tumors in the thigh region of Swiss albino mice and found that supplementation of maximal tolerated dose (100 µL/kg body weight) of the oil significantly reduced 20-methylcholanthrene induced tumor incidence and tumor volume[97]. Earlier they reported the antiproliferative activity of the *O. sanctum* oil against HeLa cells in culture[98]. Manosroi *et al.* have shown an inhibition of the proliferation of murine leukemia and human mouth epidermal carcinoma cell lines by essential oils of *O. sanctum*, *O. basilicum* and *O. americanum*[62].

#### 4.4.3. As skin permeation enhancers

The delivery of drugs via skin routes has been extensively studied. Dermal and transdermal drug delivery is often limited by poor permeability of the stratum corneum (SC) to drugs, which precludes their crossing the skin at therapeutic rates. The barrier properties of the SC can be reduced by the use of natural products. Essential oils are receiving considerable interest as enhancers to improve drug permeation. Essential oil of sweet basil (*O. basilicum*) was evaluated as skin permeation enhancer to promote the percutaneous absorption of drugs and reported it as noble enhancers[99].

#### 4.4.4. Cardiovascular and anti-lipidemic actions of *Ocimum* oils

Both *O. gratissimum* oil and eugenol decrease the blood pressure in conscious DOCA-salt hypertensive rats exhibiting its cardiovascular effects[100]. In earlier research, Lahlou *et al.* found that intravenous treatment of normotensive rats with *O. gratissimum* oil and its main constituent eugenol dose dependently decreased mean aortic pressure[101]. Eugenol was also reported to induce vasorelaxant effects on rat and rabbit thoracic aorta as well as on rat mesenteric vascular bed[102–104].

Essential oil from *O. sanctum* leaves has been shown to have a potential for lipid-lowering action. The anti-hyperlipidaemic ability of the essential oil extracted from *O. sanctum* leaves in rats fed with high cholesterol diet was investigated[105]. In conclusion, the researchers found that there was no significant differences in body weight gain, food intake, and heart weight in all groups of rats. Authors observed that phenylpropanoid compounds, the main composition of *O. sanctum* essential oil is responsible for the lipid lowering effect in high cholesterol rats[105]. In another finding by treating with *O. sanctum* oil to rats fed with a high cholesterol diet, it was found that essential oil suppressed the high serum lipid profile and atherogenic index as well as serum lactate dehydrogenase and creatine kinase MB subunit without significant effect on high serum levels of aspartate aminotransferase, alanine aminotransferase and alkaline phosphatase in rats fed with high cholesterol diet[57]. Oil was also found to decrease the high level of thiobarbituric acid reactive substances, glutathione peroxidase and superoxide dismutase without impacting catalase in the cardiac tissue while in the liver, it decreased high level of thiobarbituric acid reactive substances without significantly effecting glutathione peroxidase, superoxide dismutase and impacting catalase. Authors concluded that essential oil of *O. sanctum* has lipid-lowering and antioxidative effects that protect the heart against hypercholesterolemia.

#### 4.4.5. Immunomodulatory and CNS activity of *Ocimum* oils

In current competitive life, stress is a very common problem for human beings. Assembly of more free-radicals due to stress leads to activity adverse effects on various vital organs and tissues of the human body. *Ocimum* essential oil has gained special attention due to its biological properties, however, little is known about its immunomodulatory effects. The immunomodulatory potential of *O. sanctum* oil on some immunological parameters in stressed and non-stressed animal was carried out[106]. They concluded that by penetrating the oil it exert immunomodulatory effect by modulating GABAergic activity and influence both humoral and cell mediated immunological parameters in native non-stressed as well as stressed animals[106]. In this context, some CNS activities *viz.*, sedative, hypnotic, anticonvulsant, local anesthetic of *O. basilicum* essential oil were also investigated by Ismail[107]. When tested in mice, *O. basilicum* oil had no effect on motor activity up to a dose of 1.2 mL/kg at 90 mm post-administration. Pentobarbitone sleeping time tested in



mice was significantly increased in mice by all doses of essential oil higher than 0.2 mL/kg. Intraperitoneal of oil significantly increased in a dose dependent manner, the latency of convulsion and percent of animal exhibiting clonic seizures. He found ED<sub>50</sub> values of oil were 0.61, 0.43 and 1.27 mL/kg against convulsion induced by pentylenetetrazole, picrotoxin and strychnine, respectively. He noted that the observed anticonvulsant and hypnotic activities of oil could be related to the presence of variety of terpenic constituents in it. It has been reported that terpenes have a protective effect against pentylenetetrazole and picrotoxin-induced convulsions<sup>[108,109]</sup>. Modulation of glutamergic and GABAergic transmission are mechanisms indicated for anticonvulsant action on monoterpenes<sup>[110]</sup>. Linalool, the most abundant constituent of *Ocimum* oil, has anticonvulsant activity against pentylenetetrazole-induced convulsion and also through inhibition of glutamergic transmission as well as through suppression of voltage-gated current<sup>[111–113]</sup>. On the other hand, other main constituent of 1,8-cineole exerts anticonvulsant activity, potentiates phenobarbitone sleeping time, and has an inhibitory effect on locomotory activity<sup>[114]</sup>. Eugenol, the other representative constituent, also exerts anticonvulsant property through potentiating binding of GABA to its receptor and by increasing the affinity of these receptors to bind GABA<sup>[115]</sup>. Additionally, eugenol has anesthetic, sedative, and muscle relaxant effects<sup>[116]</sup>.

In research of Oliveira *et al.*, CNS depressant and anticonvulsant effects of *O. basilicum* (access “Maria Bonita”) leaf essential oil in different experimental models were investigated<sup>[117]</sup>. *O. basilicum* essential oil, at all doses, showed depressant CNS activity as revealed in the general pharmacological screening; decrease of spontaneous activity, ataxia, and sedation. Additionally, all doses of *O. basilicum* essential oil induced a significant increase of sleeping time ( $P < 0.05$ ) and decrease in the latency to sleep ( $P < 0.01$ ). *O. basilicum* essential oil also increased the latency for development of convulsions in pentylenetetrazol and picrotoxin tests ( $P < 0.05$ ). For pentylenetetrazol, the effect of *O. basilicum* essential oil was reversed by flumazenil. *O. basilicum* essential oil did not interfere with the convulsions induced by strychnine ( $P > 0.05$ ). Their data suggests that *O. basilicum* essential oil possesses CNS depressant and anticonvulsant properties which could be mediated by an interaction with central GABAergic receptors.

## 5. Conclusions

There are many herbal plants in the world among which *Ocimum* (Tulsi) is considered to be the ruler of herbs due to its great medicinal ethics. Various medicinal properties of Tulsi are well documented in the Hindu mythology. Considering the health beneficial effects of Tulsi, our ancestors in India insisted to plant a Tulsi sapling in everyone’s house. The composed literature revealed that

essential oils of collected *Ocimum* species from different geographical origins existed variety of chemical constituents, and the researchers found wide and varied application in traditional healthcare system. The review of past studies which has been discussed in this chapter showed that essential oils obtained from *Ocimum* plants have gained much appreciation among food scientists and researchers because of their multifold biological activities. The reviewed results aimed at attracting the attention of scientists and researchers looking for new drugs from natural products as well as those investigating the pharmaceutical diversity of essential oils. The data reported in the present study provide a basis for reviving the old art of “essential oil therapy” based on our modern scientific knowledge of their mode of action, supported by safety issues. Probably, such natural components might prove to be potentially beneficial but comparatively less toxic. Eventually, plants belonging to *Ocimum* genus could contribute a lot to solve certain economy and health problems. Thus, essential oils and their constituents of *Ocimum* species can hopefully be considered in the future for more clinical evaluations and applications, and as possible adjuvants to current medications.

## Conflict of interest statement

We declare that we have no conflict of interest.

## Acknowledgements

The authors thank the Head Department of Botany, DDU Gorakhpur University, Gorakhpur for providing necessary facilities and the Library officers, National Science Library, National Medical Library, New Delhi, FRI, Dehradun and CIMAP, Lucknow for providing library facilities. Dr. A. K. Pandey is grateful to CST UP Lucknow for financial assistance (Grant No. CST/AAS, D–09, April 3, 2007).

## References

- [1] Burt S. Essential oils: their antibacterial properties and potential applications in food—A review. *Int J Food Microbiol* 2004; **94**: 223–253.
- [2] Hedge C. A global survey of the biogeography of the Labiatae. In: Harley RM, Reynolds T, editors. *Advances in labiatae science*. UK: Royal Botanic Gardens, Kew; 1992, p. 7–17.
- [3] Baser KH. The Turkish *Origanum* species. In: Kintzios SE, editor. *Oregano: The genera Origanum and Lippia*. London: Taylor and Francis; 2002, p. 109.
- [4] Lawrence BM. Labiatae oils—mother nature’s chemical factory. Essential oils allured publishing carol stream IL; UK: Mendeley Ltd; 1993, p. 188–206. [Online] Available from: <http://legacy.library.ucsf.edu/documentStore/i/u/d/iud13c00/Siud13c00.pdf>. [Access on May 12, 2012]

- [5] Martins AP, Salgueiro LR, Vila R, Tomi F, Canigueral S, Casanova J, et al. Composition of essential oils of *Ocimum canum*, *Ocimum gratissimum* and *Ocimum minimum*. *Planta Med* 1999; **65**: 187–189.
- [6] Manosroia J, Dhumtanoma P, Manosroia A. Anti-proliferative activity of essential oil extracted from Thai medicinal plants on KB and P388 cell lines. *Cancer Lett* 2006; **235**: 114–120.
- [7] Baris O, Golloce M, Sahin R, Ozer H, Killic H, Ozkan H, et al. Biological activities of essential oil and methanol extract of *Achillea biebersteinii* Afan. (Asteraceae). *Turk J Biol* 2006; **30**: 65–73.
- [8] Jirovetz L, Buchbauer G. Analysis, chemotype and quality control of the essential oil of a new cultivated basil (*Ocimum basilicum* L.) plant from Bulgaria. *Sci Pharm* 2001; **69**: 85–89.
- [9] Mondal S, Mahapatra SC, Mirdha BR, Naik SN. Antimicrobial activity of essential oils obtained from fresh and dried leaves of *Ocimum sanctum* L. against enteric bacteria and yeast. *Acta Horticulturae* 2007; **756**: 267–270.
- [10] Silva MG, Matos FJ, Lopes PR, Silva FO, Holanda MT. Composition of essential oils from three *Ocimum* species obtained by steam and microwave distillation and supercritical CO<sub>2</sub> extraction. *ARKIVOC* 2004; **6**: 66–71.
- [11] Zheljajzkov VD, Cantrell CL, Tekwani B, Khan SI. Content, composition, and bioactivity of the essential oils of three basil genotypes as a function of harvest. *J Agric Food Chem* 2008; **56**: 380–385.
- [12] Telci I, Bayram E, Yilmaz G, Avci B. Variability in essential oil composition of Turkish basils (*Ocimum basilicum* L.). *Biochem Syst Ecol* 2006; **34**: 489–497.
- [13] Sajjadi SE. Analysis of the essential oils of two cultivated basil (*Ocimum basilicum* L.) from Iran. *Daru* 2006; **14**: 128–130.
- [14] Khalid KA, Hendawy SF, Gezawy EE. *Ocimum basilicum* L. production under organic farming. *Res J Agric Biol Sci* 2006; **2**: 25–32.
- [15] Bozin B, Mimica-Dukić N, Simin M, Anackov G. Characterization of the volatile composition of essential oils of some Lamiaceae spices and the antimicrobial and antioxidant activities of the entire oils. *J Agric Food Chem* 2006; **54**: 1822–1828.
- [16] Matasyoh LG, Matasyoh JC, Wachira FN, Kinyua MG, Muigai AW, Mukiyama TK. Chemical composition and antimicrobial activity of the essential oil of *Ocimum gratissimum* L. growing in Eastern Kenya. *Afr J Biotechnol* 2007; **6**: 760–765.
- [17] Politeo O, Jukic M, Milos M. Chemical composition and antioxidant capacity of free volatile aglycones from basil (*Ocimum basilicum* L.) compared with its essential oil. *Food Chem* 2007; **101**: 379–385.
- [18] Hussain AI, Anwar F, Sherazi ST, Przybylski R. Chemical composition, antioxidant and antimicrobial activities of basil (*Ocimum basilicum*) essential oils depends on seasonal variations. *Food Chem* 2008; **108**: 986–995.
- [19] Mondal S, Mirdha BR, Mahapatra SC. The science behind sacredness of Tulsi (*Ocimum sanctum* Linn.). *Indian J Physiol Pharmacol* 2009; **53**: 291–306.
- [20] Koul O, Suresh W, Dhaliwal GS. Essential oils as green pesticides: potential and constraints. *Biopestic Int* 2008; **4**: 63–84.
- [21] Lawrence BM. A review of the world production of essential oil. *Perfumer & Flavorist* 1985; **10**: 2–16.
- [22] Lawrence BM. A world perspective. Proceedings of the 10th international congress of essential oils, fragrances and flavors, Washington, DC, USA. Amsterdam: Elsevier Science Publisher BV; 1998, p. 161.
- [23] Yusuf M, Begum J, Mondello L, Stagno D, Alcontres I. Studies on the essential oil bearing plants of Bangladesh. Part VI. Composition of the oil of *Ocimum gratissimum*. *Flavour Frag J* 1998; **13**: 163–166.
- [24] Sanda K, Koba K, Nambo P, Gaset A. Chemical investigation of *Ocimum* species growing in Togo. *Flavour Frag J* 1998; **13**: 226–232.
- [25] Lang L, Buchbauer G. A review on recent research results (2008–2010) on essential oils as antimicrobials and antifungals. A review. *Flavour Frag J* 2012; **27**: 13–39.
- [26] Janssen AM, Scheffer JJ, Ntezurubanza L, Svendesen AB. Antimicrobial activities of some *Ocimum* species grown in Rwanda. *J Ethnopharmacol* 1998; **26**: 57–63.
- [27] Rajeshwari S. *Ocimum sanctum*. The Indian home remedy. In: *Current medical scene*. Bombay Central, Bombay: Cipla Ltd; 1992.
- [28] Zollo PH, Biyiti L, Tchoumboungang F, Menut C, Lamaty C, Bouchet PH. Aromatic plants of tropical Africa. Part XXXII. Chemical composition and antifungal activity of thirteen essential oils from aromatic plants of Cameroon. *Flavour Frag J* 1998; **13**: 107–114.
- [29] Basilico MZ, Basilico JC. Inhibitory effects of some spice essential oils on *Aspergillus ochraceus* NRRL 3174 growth and ochratoxin A production. *Lett Appl Microbiol* 1999; **29**: 238–241.
- [30] Tripathi P, Dubey NK, Banerji R, Chansouria JP. Evaluation of some essential oils as botanical fungitoxicants management of post-harvest rotting of *Citrus* fruits. *Word J Microbiol* 2004; **20**: 317–321.
- [31] Moreira JR, Ponceb AG, del Vallea CE, Rourab SI. Inhibitory parameters of essential oils to reduce a foodborne pathogen. *LWT-Food Sci Technol* 2005; **38**: 565–570.
- [32] Mbata TI, Saikia A. Antibacterial activity of essential oil from *Ocimum gratissimum* on *Listeria monocytogenes*. *Internet J Food Safety* 2005; **5**: 15–19.
- [33] Nakamura CV, Ueda-Nakamura T, Bando E, Melo AF, Cortez DA, Dias BP, et al. Antibacterial activity of *Ocimum gratissimum* L. essential oil. *Mem Inst Oswaldo Cruz* 1999; **94**: 675–678.
- [34] Sacchetti G, Medici A, Maietti S, Redice M, Muzzoli M, Manfredini S, et al. Composition and functional properties of essential oil Amazonian basil, *Ocimum micranthum* Willd. Labiatae in comparison with essential oil. *J Agric Food Chem* 2004; **52**: 3486–3491.
- [35] Weseler A, Geiss HK, Saller R, Reichling J. A novel colorimetric broth microdilution method to determine the minimum inhibitor concentration (MIC) of antibiotics and essential oils against *Helicobacter pylori*. *Pharmazie* 2005; **60**: 498–502.
- [36] Wannissorn B, Jarikasem S, Siriwangchai T, Thubthimthed S. Antibacterial properties of essential oils from Thai medicinal plants. *Fitoterapia* 2005; **76**: 233–236.
- [37] Faria TJ, Yassumoto F, Roberto RS, Souza L, Ishikawa J. Antifungal activity of essential oil isolated from *Ocimum*

- gratissimum* L. (eugenol chemotype) against phytopathogenic fungi. *Braz Arch Biol Technol* 2006; **49**: 867–871.
- [38] Atanda OO, Akpan I, Oluwafemi F. The potential of some spice essential oils in the control of *A. parasiticus* CFR 223 and aflatoxin production. *Food Control* 2007; **18**: 601–607.
- [39] Lee SJ, Han JL, Lee GS, Park MJ, Choi IG, Na KJ, et al. Antifungal effect of eugenol and nerolidol against *Microsporium gypseum* in a guinea pig model. *Biol Pharm Bull* 2007; **30**: 184–188.
- [40] Rattanachaikunsopon P, Phumkhachorn P. Antimicrobial activity of basil *Ocimum basilicum* oil against *Salmonella enteritidis* in vitro and in food. *Biosci Biotechnol Biochem* 2010; **74**: 1200.
- [41] Fatope MO, Marwah RG, Hadhrami NM, Onifade AK, Williams JR. Identification of the chemotypes of *Ocimum ferskolei* and *Ocimum basilicum* by NMR spectroscopy. *Chem Biodivers* 2008; **5**: 2457.
- [42] Kumar A, Shukla R, Singh P, Dubey NK. Chemical composition, antifungal and antiaflatoxigenic activities of *Ocimum sanctum* essential oil and its safety assessment as plant based antimicrobial. *Food Chem Toxicol* 2010; **48**: 539–543.
- [43] Koba K, Poutouli PW, Raynaud C, Chaumont J, Sanda K. Chemical composition and antimicrobial properties of different basil essential oils chemotypes from Togo. *Bangladesh J Pharmacol* 2009; **4**: 1–8.
- [44] Viillon C, Chaumont JP. Antifungal properties of essential oils and their main components upon *Cryptococcus neoformans*. *Mycopathologia* 1994; **128**: 151–153.
- [45] Pandey RR, Dubey RC, Saini S. Phytochemical and antimicrobial studies on essential oils of some aromatic plants. *Afr J Biotechnol* 2010; **9**: 4364–4368.
- [46] Singh P, Pandey AK, Sonker N, Tripathi NN. Preservation of *Buchnanania lanzan* Spreng. seeds by *Ocimum canum* Sims. essential oil. *Ann Plant Protect Sci* 2011; **19**: 407–410.
- [47] Anand AK, Manindra M, Haider SZ, Sharma A. Essential oil composition and antimicrobial activity of three *Ocimum* species from Uttarakhand (India). *Int J Pharm Sci* 2011; **3**: 223–225.
- [48] Verma RS, Bisht PS, Padalia RC, Saikia D, Chauhan A. Chemical composition and antibacterial activity of essential oil from two *Ocimum* spp. grown in sub-tropical India during spring-summer cropping season. *Asian J Tradit Med* 2011; **6**: 211–217.
- [49] Pandey AK, Singh P, Palni UT, Tripathi NN. *In vitro* antibacterial activities of the essential oils of aromatic plants against *Erwinia herbicola* (Lohnis) and *Pseudomonas putida* (Kris Hamilton). *J Serb Chem Soc* 2012; **77**: 313–323.
- [50] Halliwell. Antioxidants and human disease: a general introduction. *Nutr Rev* 1997; **55**: 44–52.
- [51] Baratta MT, Dorman HJ, Deans SG, Figueiredo AC, Barroso JG, Ruberto G. Antimicrobial and antioxidant properties of some commercial essential oils. *Flavour Frag J* 1998; **13**: 235–244.
- [52] Julisni HR, Simon JE. Antioxidant activity of Basil. In: Janick J, Whipkey A, editors. *Trends in new crops and new uses*. Alexandria: ASHS Press; 2002, p. 575–579.
- [53] Lee SJ, Umamo K, Shibamoto T, Lee KG. Identification of volatile components in basil (*Ocimum basilicum* L.) and thyme leaves (*Thymus vulgaris* L.) and their antioxidant properties. *Food Chem* 2005; **91**: 131–137.
- [54] Tomaino A, Cimino F, Zimbalatti V, Venuti V, Sulfaro V, De Pasquale A. Influence of heating on antioxidant activity and the chemical composition of some spice essential oils. *Food Chem* 2005; **89**: 549–554.
- [55] Bunrathep S, Palanuvez C, Ruangrungrasi N. Chemical composition and antioxidative activities of essential oils from four *Ocimum* species endemic to Thailand. *J Health Res* 2007; **21**: 201–206.
- [56] Trevisan MT, Silva MG, Pfundstein B, Spiegelhalter B, Owen RW. Characterization of the volatile pattern and antioxidant capacity of essential oils from different species of the genus *Ocimum*. *J Agric Food Chem* 2006; **54**: 4378–4382.
- [57] Suanarunsawat T, Ayutthaya WD, Songsak T, Thirawarapan S, Pongshompoo S. Antioxidant activity and lipid-lowering effect of essential oils extracted from *Ocimum sanctum* L. leaves in rats fed with a high cholesterol diet. *J Clin Biochem Nutr* 2010; **46**: 52–59.
- [58] Hanif MA, Al-Maskari MY, Al-Maskari A, Al-Shukaili A, Al-Maskari AY, Al-Sabahi JN. Essential oil composition, antimicrobial and antioxidant activities of unexplored Omani basil. *J Med Plant Res* 2011; **5**: 751–757.
- [59] Keita SM, Vincent C, Schmidt JP, Arnason JT, Belanger A. Efficacy of essential oil of *Ocimum basilicum* L. and *Ocimum gratissimum* L. applied as an insecticidal fumigant and powder to control *Callosobruchus maculatus* (F.) Coleoptera: Bruchidae. *J Stored Prod Res* 2001; **37**: 339–349.
- [60] Salvatore A, Borkosky S, Willink E, Bardon A. Toxic effects of lemon peel constituents on *Ceratitis capitata*. *J Chem Ecol* 2004; **30**: 323–333.
- [61] Paula JP, Farago PV, Checchia LE, Hirose KM, Ribas JL. Repellent activity of *Ocimum selloi* Benth. essential oil (variety eugenol) against *Anopheles braziliensis* Chagas. *Acta Farm Bonaerense* 2004; **23**: 376–378.
- [62] Popovic Z, Kostic M, Popovic S, Skoric S. Bioactivities of essential oils from basil and sage to *Sitophilus oryzae* L. *Biotechnol Biotec Equip* 2006; **20**: 36–40.
- [63] Esther NM, Ahmed H, Suliman E, Edward ML, Mark K. Repellent and acaricidal properties of *Ocimum suave* against *Rhipicephalus appendiculatus* tick. *Exp Appl Acarol* 1995; **19**: 11–18.
- [64] Hassanli A, Iwande W, Ole-Sitayo N, Moreka L, Nokoe S. Weevil repellent constituents of *Ocimum suave* leaves and *Eugenia caryophyllata* cloves used as grain protectant in parts of Eastern Africa. *Discov Innovat* 1990; **2**: 91–95.
- [65] Sosan MB, Adewoyin FB, Adewunmi CO. Laticidal properties of 3 indigenous plant oils on the mosquito *Aedes aegypti*. *Nig J Nat Prod Med* 2001; **5**: 30–33.
- [66] Luckwa N, Masidza C, Nyazema NZ, Curtis CF, Mwaiko GL. Efficacy and duration of activity of *Lippia javanica* Spreng, *Ocimum canum* Sims. and a commercial repellent against *Aedes aegypti*. In: Hostettman K, Chinyanganya F, Maillard M, Wolfende JL, editors. International organization for chemistry in science and development (IOCD) Symposium; Zimbabwe: Victoria Falls; 1996, p. 321–325.
- [67] Aslan I, Ozbek H, Çalmasur O, Sahin F. Toxicity of essential oil vapours to two greenhouse pests, *Tetranychus urticae* Koch and

- Bemisia tabaci* Genn. *Ind Crops Prod* 2004; **19**: 167–173.
- [68] Cavalcanti ES, Morais SM, Lima MA, Santana EW. Larvicidal activity of essential oils from Brazilian plants against *Aedes aegypti* L. *Mem Inst Oswaldo Cruz* 2004; **99**: 541–544.
- [69] Ilondu EM, Egwunyenga OA, Iloh AC. The effects of wild Basil (*Ocimum suave*) Wild Labiateae: (Lamiaceae) leaf dust as protectant of cereals, against *Rhyzopertha dominica* (F) Coleoptera: (Bostryidae). *J Natl Acad Adv Sci* 2004; **3**: 129–134.
- [70] Olonisakin A, Oladimeji MO, Lajide L. Bioactivity of steam distilled oils against the cowpea bruchids, *Callosobruchus maculatus* (F) infesting stored cowpea seeds. *Pak J Boil Sci* 2006; **9**: 1271–1275.
- [71] Lopez MD, Jordan MJ, Pascual-villalobos MJ. Toxic compounds in essential oils of coriander, caraway and basil active against stored rice pests. *J Stored Prod Res* 2008; **44**: 273–278.
- [72] Bakr RF, Fattah HM, Salim NM, Atiya NH. Insecticidal activity of four volatile oils on two museum insects pests of Egypt. *Acad J Biolog Sci* 2010; **2**: 57–66.
- [73] Manzoor F, Beena W, Malik S, Naz N, Naz S, Syed WH. Preliminary evaluation of *Ocimum sanctum* as toxicant and repellent against termite, *Heterotermes indicola* (wasmann) (Isoptera: Rhinotermitidae). *Pak J Sci* 2011; **63**: 59–62.
- [74] Chang CL, Cho IK, Li QX. Insecticidal activity of Basil oil, trans-Anethole, Estragole, and Linalool to adult fruit flies of *Ceratitis capitata*, *Bactrocera dorsalis*, and *Bactrocera cucurbitae*. *J Econ Entomol* 2009; **102**: 203–209.
- [75] Ojjanwuna CC, Edafemakor AG, Iloh AC. Toxicity of *Ocimum suave* (wild basil) leaf oil on adult housefly (*Musca domestica*). *Int Res J Agric Sci Soil Sci* 2011; **10**: 417–420.
- [76] Pandey AK, Tripathi NN. Post-harvest fungal and insect deterioration of pigeon pea seeds and their management by plant volatiles. *J Indian Bot Soc* 2011; **90**: 326–331.
- [77] Rajamma AJ, Dubey S, Sateesha SB, Tiwari SN, Ghosh SK. Comparative larvicidal activity of different species of *Ocimum* against *Culex quinquefasciatus*. *Nat Prod Res* 2011; **25**(20): 1916–1922.
- [78] Pandey R, Kalra A, Tandon S, Mehrotra N, Singh HN, Kumar S. Essential oils as potent sources of nematicidal compounds. *J Phytopathol* 2000; **148**: 501–502.
- [79] Holetz FB, Ueda-Nakamura T, Filho PD, Cortez DA, Morgado-diaz JA, Nakamura CV. Effect of essential oil of *Ocimum gratissimum* on the trypanosomatid *Herpetomonas samuelpessoai*. *Acta Protozool* 2003; **42**: 269–276.
- [80] Tchoumboungang F, Zollo PH, Dagne E, Mekonnen Y. *In vivo* antimalarial activity of essential oils from *Cymbopogon citratus* and *Ocimum gratissimum* on mice infected with *Plasmodium berghei*. *Planta Med* 2005; **71**: 20–23.
- [81] Santoro GF, Cardoso MG, Guimaraes LG, Mendonca LZ, Soares MJ. *Trypanosoma cruzi*: activity of essential oils from *Achillea millefolium* L., *Syzygium aromaticum* L. and *Ocimum basilicum* L. on epimastigotes and trypomastigotes. *Exp Parasitol* 2007; **116**: 283–290.
- [82] Singh S, Agrawal SS. Anti-asthmatic and anti-inflammatory activity of *Ocimum sanctum* Linn. *J Res Edu Indian Med* 1991; **7–9**: 23–28.
- [83] Singh S, Majumdar DK, Rehan HM. Evaluation of anti-inflammatory potential of fixed oil of *Ocimum sanctum* (Holybasil) and its possible mechanism of action. *J Ethnopharmacol* 1996; **54**: 19–26.
- [84] Singh S, Majumdar DK. Evaluation of anti-inflammatory activity of fatty acids of *Ocimum sanctum* fixed oil. *Indian J Exp Biol* 1997; **35**: 380–383.
- [85] Singh S, Majumdar DK. Anti-inflammatory and antipyretic activities of *Ocimum sanctum* fixed oil. *Pharmaceut Biol* 1995; **33**: 288–292.
- [86] Singh S, Majumdar DK. Analgesic activity of *Ocimum sanctum* and its possible mechanism of action. *Pharmaceut Biol* 1995; **33**: 188–192.
- [87] Sahouo GB, Tonzibo ZF, Boti B, Chopard C, Mahy JP, Yao TN. Anti-inflammatory and analgesic activities: Chemical constituents of essential oils of *Ocimum gratissimum*, *Eucalyptus citriodora* and *Cymbopogon giganteus* inhibited lipooxygenase L-1 and cyclooxygenase of PGHS. *B Chem Soc Ethiopia* 2003; **17**: 191–197.
- [88] Van der Oudera J, Buytenhek M, Slikkerveer FJ, Van Dorp DA. Purification and characterization of prostaglandin endoperoxide synthetase from sheep vascular glands. *Biochem Biophys Acta* 1977; **487**: 315–331.
- [89] Singh S, Rehan HM, Majumdar DK. Effect of *Ocimum sanctum* fixed oil on blood pressure, blood clotting time and pentobarbitone-induced sleeping time. *J Ethnopharmacol* 2001; **78**: 139–143.
- [90] Pessoa LM, Morais SM, Bevilaqua CM, Luciano JH. Anthelmintic activity of essential oil of *Ocimum gratissimum* Linn. and eugenol against *Haemonchus contortus*. *Vet Parasitol* 2002; **109**: 59–63.
- [91] Lino CS, Gomes PB, Lucetti DL, Diógenes JPL, Sousa FC, Silva MG, et al. Evaluation of antinociceptive and anti-inflammatory activities of the essential oil (EO) of *Ocimum micranthum* Willd. from Northeastern Brazil. *Phytother Res* 2005; **19**: 708–712.
- [92] Rabelo M, Souza EP, Soares PM, Miranda AV, Matos FJ, Criddle DN. Antinociceptive properties of the essential oil of *Ocimum gratissimum* L. (Labiatae) in mice. *Braz J Med Biol Res* 2003; **36**: 521–524.
- [93] Sen P. Therapeutic potentials of Tulsi: from experience to facts. *Drugs News Views* 1993; **1**: 15–21.
- [94] Mnadal S, Das DN, Dey K. *Ocimum sanctum* Linn—A study on gastric ulceration and gastric secretion in rats. *Indian J Physiol Pharmacol* 1993; **37**: 91–92.
- [95] Miller EC, Swanson AB, Phillips DH, Fletcher TL, Liem A, Miller JA. Structure-activity studies of the carcinogenicities in the mouse and rat of some naturally occurring and synthetic alkenylbenzene derivatives related to safrole and estragole. *Cancer Res* 1983; **43**: 1124–1134.
- [96] Anthony A, Caldwell G, Hutt AG, Smith RL. Metabolism of estragole in rat and mouse and influence of dose size on excretion of the proximate carcinogen 10-hydroxyestragole. *Food Chem Toxicol* 1987; **25**: 799–806.
- [97] Prakash J, Gupta SK. Chemo-preventive activity of *Ocimum sanctum* seed oil. *J Ethnopharmacol* 2000; **72**: 29–34.
- [98] Prakash J, Gupta SK, Joshi S. Antiproliferative studies on *Ocimum sanctum*. *Indian J Pharmacol* 1999; **31**: 79.
- [99] Fang JY, Leu Y, Hwang T, Cheng H. Essential oils from sweet basil (*Ocimum basilicum*) as novel enhancers to accelerate transdermal drug delivery. *Biol Pharm Bull* 2004; **27**: 1819–1825.



- [100] Interaminense LF, Leal-Cardoso JH, Magalhaes PJ, Duarte GP, Lahlou S. Enhanced hypotensive effects of the essential oil of *Ocimum gratissimum* leaves and its main constituent, eugenol, on DOCA-Salt hypertensive conscious rats. *Planta Med* 2005; **71**: 376–378.
- [101] Lahlou S, Interaminense LF, Magalhaes PJ, Leal-Cardoso JH, Duarte GP. Cardiovascular effects of eugenol, a phenolic compound present in many plant essential oils, in normotensive rats. *J Cardiovasc Pharmacol* 2004; **43**: 250–257.
- [102] Damiani CE, Rossoni LV, Vassallo DV. Vasorelaxant effects of eugenol on rat thoracic aorta. *Vasc Pharmacol* 2003; **40**: 59–66.
- [103] Nishijima H, Uchida R, Kameyama K, Kawakami N, Ohkubo T, Kitamura K. Mechanisms mediating the vasorelaxing action of eugenol, a pungent oil, on rabbit arterial tissue. *Jpn J Pharmacol* 1999; **79**: 327–334.
- [104] Criddle DN, Madeira SV, de Moura RS. Endothelium-dependent and independent vasodilator effects of eugenol in the rat mesenteric vascular bed. *J Pharm Pharmacol* 2003; **55**: 359–365.
- [105] Suanarunsawat T, Ayuthaya WD, Songsak T, Rattanamahaphoom J. Anti-lipidemic actions of essential oil extracted from *Ocimum sanctum* L. leaves in rats fed with high cholesterol diet. *J Appl Biomed* 2009; **7**: 45–53.
- [106] Mediratta PK, Sharma KK, Singh S. Evaluation of immunomodulatory potential of *Ocimum sanctum* seed oil and its possible mechanism of action. *J Ethnopharmacol* 2002; **80**: 15–20.
- [107] Ismail M. Central properties and chemical composition of *Ocimum basilicum* essential oil. *Pharm Biol* 2006; **44**: 619–626.
- [108] Librowski T, Czarnecki R, Mendyk A, Jastrzebska M. Influence of new monoterpene homologues of GABA on the central nervous system activity in mice. *Pol J Pharmacol* 2000; **52**: 317–321.
- [109] Brum LF, Elisabetsky E, Souza D. Effects of linalool on [(3)H] MK801 and [(3)H] muscimol binding in mouse cortical membranes. *Phytother Res* 2001; **15**: 422–425.
- [110] Szabadiss J, Erdelyi L. Pre- and postsynaptic effects of eugenol and related compounds on *Helix pomatia* L. neurons. *Acta Biol Hung* 2000; **51**: 265–273.
- [111] Elisabetsky E, Brum L, Souza D. Anticonvulsant properties of linalool in glutamate-related seizure models. *Phytomedicine* 1995; **6**: 107–113.
- [112] Silva L, Emanuelli T, Souza D, Elisabetsky E. Effects of linalool on glutamate release and uptake in mouse cortical synaptosomes. *Neuro Chem Res* 2001; **26**: 191–194.
- [113] Narusuye K, Kawai F, Matsuzaki K, Miyachi E. Linalool suppresses voltage-gated currents in sensory neurons and cerebellar Purkinje cells. *J Neural Transm* 2005; **112**: 193–203.
- [114] Santos FA, Rao VS. Anti-inflammatory and antinociceptive effects of 1, 8-cineole, a terpenoid oxide present in many plant essential oils. *Phytother Res* 2000; **4**: 240–244.
- [115] Aoshima H, Hamamoto K. Potentiation of GABA receptors expressed in *Xenopus* oocytes by perfume and phytoncid. *Biosci Biotechnol Biochem* 1999; **63**: 743–748.
- [116] Dallmerier K, Carlini EA. Anesthetic, hypothermic, myorelaxant and anticonvulsant effects of synthetic eugenol derivatives and natural analogues. *Pharmacology* 1981; **22**: 113–127.
- [117] Oliveira JS, Porto LA, Estevam CS, Siqueira RS, Alves PB, Niculau ES, et al. Phytochemical screening and anticonvulsant property of *Ocimum basilicum* leaf essential oil. *Boletim Latinoamericano y del Caribe de Plantas Medicinales y Aromaticas* 2009; **8**: 195–202.
- [118] Mondawi BM, Duprey RJ, Magboul AZ, Satti AM. Constituents of essential oil of *Ocimum basilicum* var. *thyrsiflorum*. *Fitoterapia* 1984; **55**: 60–61.
- [119] Ntezurbanza L, Scheffer JJ, Svendsen B. Composition of the essential oil of *Ocimum gratissimum* grown in Rwanda. *Planta Med* 1987; **53**: 421–423.
- [120] Akgul A. Volatile oil composition of sweet basil (*Ocimum basilicum* L.) cultivating in Turkey. *Nahrung* 1989; **33**: 87–88.
- [121] Khatri LM, Nasir MK, Saleem R, Noor F. Evaluation of Pakistani sweet basil oil for commercial exploitation. *Pak J Sci Ind Res* 1995; **38**: 281–282.
- [122] Marotti M, Piccaglia R, Giovanelli E. Differences in essential oil composition of basil (*Ocimum basilicum* L.) Italian cultivars related to morphological characteristics. *J Agric Food Chem* 1996; **14**: 3926–3929.
- [123] Chalchat JC, Garry RP, Sidibe L, Marama M. Aromatic plants of Mali (I): Chemical composition of essential oils of *Ocimum basilicum* L. *J Essent Oil Res* 1999; **11**: 375–380.
- [124] Keita SM, Vincent C, Schmit JP, Belanger A. Essential oil composition of *Ocimum basilicum* L., *Ocimum gratissimum* L. and *Ocimum suave* L. in the Republic of Guinea. *Flavour Frag J* 2000; **15**: 339–341.
- [125] Mondell L, Zappia G, Cotroneo A, Bonaccorsi I, Chowdhury JU, Yusuf M, et al. Studies on the essential oil-bearing plants of Bangladesh Part VIII. Composition of some *Ocimum* oils *Ocimum basilicum* L. var. *purpurascens*, *Ocimum sanctum* L. green, *Ocimum sanctum* L. purple, *Ocimum americanum* L., citral type, *Ocimum americanum* L., camphor type. *Flavour Frag J* 2002; **17**: 335–340.
- [126] Silva MG, Matos FJ, Machado MI, Craveiro AA. Essential oils of *Ocimum basilicum* L., *Ocimum basilicum* var. *minimum* L. and *Ocimum basilicum* var. *purpurascens* Benth. grown in north-eastern Brazil. *Flavour Frag J* 2003; **18**: 13–14.
- [127] Ozcan M, Chalchat J. Essential oil composition of *Ocimum basilicum* L. and *Ocimum minimum* L. in Turkey. *Czech J Food Sci* 2002; **20**: 223–228.
- [128] Kasali AA, Eshilokun AO, Adeola S, Winterhalter P, Knapp H, Bonlander B, et al. Volatile oil composition of new chemotype of *Ocimum basilicum* L. from Nigeria. *Flavour Frag J* 2005; **20**: 45–47.
- [129] Awasthi PK, Dixit SC. Chemical compositions of *Ocimum sanctum* Shyama and *Ocimum sanctum* Rama oils from the plains of Northern India. *J Essent Oil Bear Plants* 2007; **10**: 292–296.
- [130] Zoghbi MG, Oliveira J, Andrade EH, Trigo JR, Fonseca RG, Rocha AE. Variation in volatiles of *Ocimum campechianum* Mill. and *Ocimum gratissimum* L. cultivated in north of Brazil. *J Essent Oil Bear Plants* 2007; **10**: 229–240.
- [131] Mohhiuddin M, Chowdhury MJ, Alam MK, Hossain MK. Chemical composition of essential oil of four flavouring plants used by the tribal people of Bandarban hill district in Bangladesh. *Int J Med Arom Plants* 2012; **2**: 106–113.
- [132] Sastry KP, Kumar RR, Kumar AN, Sneha G, Elizabeth M. Morpho-chemical description and antimicrobial activity of different *Ocimum* species. *J Plant Dev* 2012; **19**: 53–64.