



Contents lists available at ScienceDirect

Asian Pacific Journal of Tropical Biomedicine

journal homepage: [www.elsevier.com/locate/apjtb](http://www.elsevier.com/locate/apjtb)

Document heading

# Aldose reductase inhibitory principles from the whole plant of *Hybanthus enneaspermus* (Linn) F. Muell

Patel DK, Prasad SK, Sairam K, Hemalatha S\*

Pharmacognosy Research Laboratory, Department of Pharmaceutics, Institute of Technology, Banaras Hindu University, Varanasi–221005, India

## ARTICLE INFO

## Article history:

Received 8 January 2012

Received in revised form 12 January 2012

Accepted March 3 2012

Available online 28 April 2012

## Keywords:

*Hybanthus enneaspermus*

Aldose reductase

Rat lens

Beta sitosterol

diabet

flavonoids

## ABSTRACT

**Objective:** To evaluate active aldose reductase (AR) inhibitory principles from ethanolic extract and one isolated compound beta sitostirrol from *Hybanthus enneaspermus* (*H. enneaspermus*) Linn F. Muell. (Violaceae), used in the treatment of various complications including diabetes. **Methods:** AR inhibitory activities of *H. enneaspermus* and isolated compound beta sitosterol against rat lens AR enzyme were investigated *in vitro* using quercetin as a standard drug. **Results:** There was a significant difference in the AR inhibitory activity, and it was found to be more in *H. enneaspermus* [ $IC_{50}$  ( $49.46 \pm 2.26 \mu\text{g/mL}$ )] followed by beta sitosterol [ $IC_{50}$  ( $82.71 \pm 0.42 \mu\text{g/mL}$ )] but was less compared to standard quercetin [ $IC_{50}$  ( $3.326 \pm 0.11 \mu\text{g/mL}$ )]. From the value of  $V_{max}$ ,  $K_m$  and  $K_i$  it was found that *H. enneaspermus* inhibited enzyme in a non competitive manner, whereas beta sitosterol inhibited enzyme in a competitive manner. **Conclusions:** *H. enneaspermus* showed significant AR inhibitory activity compared to the isolated compound beta sitosterol, which may be due to its high phenolic and flavonoid content. So this plant can be used to treat diabetic complications in the future, but further evaluation is necessary to know the exact mode of action involved in this process.

## 1. Introduction

Chronic hyperglycemia can cause various complications including blindness, renal failure, neuropathy, limb amputation, myocardial infarction and stroke, which are mainly due to elevated levels of glucose in tissue such as the nerve, kidney, retina and lens. Many biochemical pathways are involved in the process associated with this complication such as polyol pathway[1,2]. Aldose reductase (AR) which belongs to aldo–keto reductase is the first and rate–limiting enzyme in the polyol pathway which converts glucose to sorbitol utilizing Nicotinamide Adenine Dinucleotide Phosphate (NADPH) as a cofactor. Accumulation of sorbitol leads to osmotic swelling, changes in membrane permeability, and also oxidative stress culminating in tissue injury. It also changes the reduction potential of the cell and induces oxidative stress. Oxidative stress is mainly due to depleted intracellular levels of reduced glutathione, increased non–enzymatic glycation

and activation of protein kinase C[2]. From earlier study it was proven that inhibition of AR enzyme could be effective in the treatment of cataract, retinopathy, nephropathy and neuropathy. Eventhough both synthetic and natural AR inhibitors have been reported to prevent some diabetic complications in animal models and clinical trials, still very limited number of AR inhibitor is there due to deleterious side effects and poor penetration of target tissues such as nerve and retina[3].

Many AR inhibitors obtained from natural sources such as quercitrin, guaijaverin, and desmanthin–1 showed significant inhibitory activity against AR enzyme as mentioned in some other study[4]. In the earlier studies it has been shown that cataract progression can be slowed down or prevented by the use of natural therapies. Plants which have high flavonoid content were found to have significant inhibitory potential against AR enzyme[5]. Quercitrin–3–rhamnosid a type of flavonoid was the first synthetic AR inhibitor tested *in vivo* against AR enzyme. In another study it was found that oral consumption of quercetin and naringin decreases sorbitol accumulation in lenses of diabetic rats[6]. From the data obtained in earlier study it was found that, 4–oxo–4h–chromen ring in flavonoid structure is necessary to inhibit the AR enzyme. High ascorbic acid concentration

\*Corresponding author: S Hemalatha, Pharmacognosy Research Laboratory, Department of Pharmaceutics, Institute of Technology, Banaras Hindu University, Varanasi–221005, India.

Tel: +91 9415256481

E–mail: [shemalatha.phe@itbhu.ac.in](mailto:shemalatha.phe@itbhu.ac.in)

Foundation Project: Supported by University Grants Commission, New Delhi.

can also inhibit erythrocyte AR where aminoguanidine prevented the retinopathy in diabetic rats[7]. Two other isoflavone compounds, tectorigenin and irigenin, were also shown strong inhibitory activity against AR enzyme[8].

*Hybanthus enneaspermus* (*H. enneaspermus*) Linn F. Muell. (Violaceae) is an herb distributed in the tropical and subtropical regions of world. The plant is used to treat urinary infections, diarrhea, cholera, leucorrhoea, gonorrhoea, dysuria, inflammation, sterility and diabetes in the different systems of medicine[9,10]. Pharmacologically it has been tested for anti-inflammatory, antitussive, antiplasmodial, antimicrobial, anticonvulsant and antidiabetic activities[11,12]. Due to presence of significant amount of phenol and flavonoid contents with relevant antioxidant and antidiabetic potential, the aim of the present study was to evaluate a potential new source of AR inhibitors from the ethanolic extract of *H. enneaspermus* which will prove beneficial for treatment of diabetes related secondary complication such as cataract.

## 2. Materials and methods

### 2.1. Chemicals and reagents

All the chemicals and reagents used in this study were of analytical grade and procured from Sigma Chemical Co., New Delhi and Merck India Ltd. For absorbance measurements ultraviolet–visible spectrophotometer (Shimadzu, Pharmaspec–1700) was used.

### 2.2. Procurement of plant material

Plant material (*H. enneaspermus*) was procured from herbal vendors in Chennai, and authenticated by the chief botanist TAMPOL Anna Hospital Chennai, voucher specimen (Cog/HE/01/08) was kept for further reference at Laboratory herbarium, Department of Pharmaceutics, Institute of Technology, Banaras Hindu University, Varanasi, India.

### 2.3. Preparation of extract

The air-dried whole plant parts of *H. enneaspermus* (1 kg) were extracted with ethanol in a Soxhlet apparatus for 12 h. After filtration of the solvent, the organic phases were concentrated under a vacuum to get dry extract.

### 2.4. Phytochemical evaluation

#### 2.4.1. Fingerprint analysis

HPTLC fingerprint analysis of *H. enneaspermus* was performed in chloroform : methanol (8:2) solvent system, mode of application was automatic, volume of sample was 5  $\mu$ L, development mode was ascending and scanning was done at 254 nm using win cats software.

#### 2.4.2. Isolation process

Powdered material of *H. enneaspermus* was refluxed with n-Hexane for 2 h, after filtration organic solvent was concentrated under vacuum to get dry extract. n-Hexane soluble fraction (15 g) was packed on column with silica gel used as a stationary phase and successively eluted with a stepwise gradient of n-Hexane–ethyl acetate (0, 1, 2, 5, 10, 20 and so on up to 100%) and then were washed with methanol. The active guided fractions (6:4) were collected, mix all the same fractions which was checked by TLC, concentrated and finally kept for 5 day. Solid mass was dissolved in the methanol and were separated. This process was then repeated until the colour of the solution became colourless. Finally the colourless solid mass was dissolved in the chloroform and 2 mL of methanol was further added to it and was kept at room temperature for 5 days from which white soft crystal were obtained. For the determination of the structure and nature of compound, various type of data analysis such as melting point, infrared spectrum, NMR spectral analysis have been done. Finally the isolated compound was confirmed by TLC fingerprint analysis using standard beta sitosterol as a reference compound.

### 2.5. Measurement of AR inhibitory activity in vitro

#### 2.5.1. Experimental animals

Healthy albino rats of Charles foster strain (140–180 g) were used in this study, they were housed under standard laboratory conditions [12 h light/12 h darkness, (21  $\pm$  2)  $^{\circ}$ C]. Animals were given standard pellets diet (Mona laboratory animal feed) and water *ad libitum* throughout the experimental period. Experimental study was approved by the Institutional Animal Ethical Committee of Institute of Medical Sciences, Banaras Hindu University, Varanasi, India.

#### 2.5.2. Purification of lens AR

Eyes of normal Wistar rats were removed immediately after sacrificing them, washed with saline and their fresh weights were measured. 10% homogenate was prepared from cleared lenses in 0.1 M phosphate buffer saline (PBS, pH 7.4). After centrifugation at 5000  $\times$  g for 10 min in a refrigerated centrifuge, the supernatant was collected and kept in ice[5].

#### 2.5.3. Determination of protein content

Protein content of supernatant was determined by the method of Lowry *et al*[5].

#### 2.5.4. Determination of AR activity

A sample cuvette containing 0.7 mL of phosphate buffer (0.067 M), 0.1 mL of NADPH (25  $\times$  10<sup>-5</sup> M), 0.1 mL of lens supernatant, 0.1 mL of DL-glyceraldehyde (5 $\times$ 10<sup>-4</sup> M) as a substrate to a final volume of 1 mL was read against a reference cuvette containing all components but the substrate, DL-glyceraldehyde. The final pH of the reaction mixture was 6.2. The enzymatic reaction was started by

the addition of the substrate and the absorbance (OD) was recorded at 340 nm for 3 min at 30 second interval. AR activity was expressed as OD/min/mg protein[5].

### 2.5.5. Determination of AR activity and compound

Dried extracts and beta sitosterol were reconstituted in PBS to prepare stock solutions. To determine their AR inhibiting activity, 0.1 mL of each concentration (25, 50, 75, 100, 200 and 300  $\mu$ g/mL) from various stock solutions prepared in the PBS was added to both the reference and standard cuvettes. The reaction was initiated by the addition of 0.1 mL DL-glyceraldehyde and the rate of reaction was measured as described above[5].

### 2.5.6. Determination of enzyme kinetic

Enzyme kinetics  $K_m$  and  $V_{max}$  of AR were determined with varying concentrations of DL-glyceraldehyde as substrate in the absence and presence of different concentrations of *H. enneaspermus* and beta sitosterol by Lineweaver-Burk double reciprocal plots. Inhibitory constant ( $K_i$ ) was derived by plotting slopes obtained from Lineweaver-Burk plots versus *H. enneaspermus* and beta sitosterol concentration[5].

## 2.6. Statistical analysis


Results are expressed as Mean  $\pm$  SEM of three independent determinations. Linear regression analysis was performed, quoting the correlation coefficient  $r^2$ . Two-ways ANOVA followed by Bonferroni post test was performed for evaluation of all data. GraphPad Prism (version 5) software was used for all statistical analysis, and  $P < 0.05$  was considered as significance.

## 3. Results

### 3.1. Phytochemical evaluation

From the column chromatography study it was found that *H. enneaspermus* contain significant amount of beta sitosterol, % yield and structure of the isolated compound

were presented in Table 1 and Figure 1. HPTLC fingerprint analysis showed that *H. enneaspermus* contain nine spots, where two spots were found to be prominent, which was presented in the Figure 2.



**Figure 1.** Structure of the isolated compound (Beta sitosterol).

### 3.2. AR inhibitory activity

The present study deals with AR inhibitory potential of *H. enneaspermus* and isolated compound beta sitosterol on rat lens AR enzyme. *H. enneaspermus* have been chosen in the present study based on the ethnopharmacological uses. From the data obtained in the present study it was found that the inhibitory potential of *H. enneaspermus* and beta sitosterol in the form of  $IC_{50}$  value was found to be significant against AR enzyme. The AR concentration in normal rat lens was found to be  $0.0148 \pm 0.0006$  which was considered 100%. At the concentration of 100  $\mu$ g/mL *H. enneaspermus* extract showed more percentage of inhibition (86.61%) compare to the isolated beta sitosterol (78.78%). The  $IC_{50}$  value of the *H. enneaspermus* and beta sitosterol were presented in Table 1 and Figure 3.

### 3.3. Study of enzyme kinetic

Study of the enzyme kinetic was performed in Lineweaver-Burk plot using DL-glyceraldehyde as a substrate. The Lineweaver-Burk plot was made in between 1/velocity Vs

**Table 1**  
AR inhibitory potential of *H. enneaspermus* and beta sitosterol.

S. No	Extract/compounds	% yield	AR inhibitory activity* [ $IC_{50}$ ( $\mu$ g/mL)]
1	<i>H. enneaspermus</i>	13.45	$49.460 \pm 2.260$
2	Beta sitosterol	5.98 <sup>#</sup>	$82.710 \pm 0.420$
3	Quercetin	NA	$3.326 \pm 0.110$

\*- Values represent Mean  $\pm$  SEM of three independent measurements; NA- Not analysed; #- compared to the n-hexane extract .

**Table 2**  
Kinetic parameter of rat lens AR.

S. No	AR activity	$V_{max}$	$K_m$	$K_i$
1	DL-glyceraldehyde	$0.1686 \pm 0.0006$	$0.6562 \pm 0.0045$	$0.0000 \pm 0.0000$
2	DL-glyceraldehyde + Beta sitosterol	$0.1393 \pm 0.0003^a$	$0.6903 \pm 0.0035^a$	$0.1945 \pm 0.0004$
3	DL-glyceraldehyde + <i>H. enneaspermus</i>	$0.1429 \pm 0.0030^a$	$0.9627 \pm 0.0004^{ab}$	$0.1622 \pm 0.0003^b$

Values are Mean  $\pm$  SEM of three independent data in each group; a: compare to the DL-glyceraldehyde, b: compare to the DL-glyceraldehyde + Beta sitosterol (Two-way ANOVA followed by Bonferroni post test,  $P < 0.05$  was considered as significance).

1/ [DL-glyceraldehyde]. From the value of  $V_{max}$ ,  $K_m$  and  $K_i$  it was found that *H. enneaspermus* inhibited enzyme in non competitive manner, whereas beta sitosterol inhibited enzyme in competitive manner. Values of  $V_{max}$ ,  $K_m$  and  $K_i$  were presented in the Table 2, where the Lineweaver–Burk plot was presented in Figure 3.




Figure 2. HPTLC fingerprint analysis of *H. enneaspermus*.




Figure 3. (A) Effect of *H. enneaspermus*, Beta sitosterol and quercetin on AR inhibitory activity. (B) Effects of *H. enneaspermus* and Beta sitosterol on the Lineweaver–Burk plot of AR activity with DL-glyceraldehyde as substrate.

#### 4. Discussion

Through the literature review it was found that *H. enneaspermus* has impressive ethnopharmacological profile which is used in the different systems of the medicine for the treatment of the various type of complication including diabetic[10]. On the basis of these ethnopharmacological data present work dealt with the evaluation of AR inhibitory potential of the *H. enneaspermus* and isolated compound beta sitosterol on rat lens AR enzyme. For this purpose extraction was performed, further isolation was done through column chromatography. Further HPTLC fingerprint analysis of ethanolic extract of *H. enneaspermus* was carried out followed by performing the AR inhibitory activity of *H.*

*enneaspermus* and beta sitosterol on rat lens AR enzyme.

The phytoconstituents reported in the plant revealed the presence of majorly flavonoids, terpenes, phenols, anthraquinones, glycosides, polyoses, alkaloids, and saponins, tannins. Among these compound phenolic compounds (71.95 mg/g) and flavonoids (20 mg/g) were found to be in abundant quantity[12]. The above phytoconstituents could be used as an analytical tool for the standardization of *H. enneaspermus*, if any adulterants or contaminants were present in the plant material it can be identified by the use of this parameter. HPTLC fingerprint analysis data can also be used for the comparative analysis of the sample, collected in different places and time interval. Ethanol was used for the extraction process due to its high extractability and less toxicity. For the isolation purpose n-Hexane was used as a solvent due the non polar nature of the isolated compound beta sitosterol. In the ethanol extract total phenol and flavonoid was found to be at significant level which could be explained by the possible formation of complexes between phenolic compounds with other components, which are more extractable in ethanol[13,14].

Since mice are devoid of lens AR enzyme, which do not develop sugar cataract in hyperglycemic conditions therefore, Wister rat lens were used as AR enzyme source for initial screening of *H. enneaspermus* and beta sitosterol[15]. Further compared to other tissues, AR enzyme was found to be in higher percentage in the eye lens, particularly rat lens[16]. The inhibitory effects of plant phytochemicals including polyphenols, known as natural antioxidants, and their antioxidant activities are important for human health, which can lower the blood glucose level in diabetes and is capable of reducing oxidative stress by scavenging reactive oxygen species, and prevent cell damage[17–19]. Some flavonoids and polyphenols were found to be effective against  $\alpha$ -glucosidase and AR. For the development of effective, safe and potentially useful products such as commercial  $\alpha$ -glucosidase and AR inhibitors from natural sources, various researches have been done[20]. From our earlier study it was found that *H. enneaspermus* up to 5 g/kg bw has no toxic effect in animal experimental study suggesting that this plant is safe for human being.

From earlier work it was found that root of *Salacia oblonga*, *Salviae multiorrhizae*, *Glycerrhiza uralensis*, *Radix astragali*, puerarin and curcumin have significant inhibitory potential against AR enzyme. Further, theses plants have also been reported to contain significant amount of bioflavonoids, which have also been reported to have AR inhibitory potential against AR enzyme. So in the present study AR inhibitory activity of the *H. enneaspermus* might be due to the presence of these compounds[3,5]. In another study it was found that flavonoids are among the most potent AR inhibitors known, whereas *Ocimum sanctum* was found to be most effective against AR enzyme compared to other extract such as *Aralia* extract. Other compound such as quercetin, flavanone and flavonol glucosides has also been reported to have significant AR inhibitory activity[21,22]. Some of the

AR inhibitors (zopolrestat, ponalrestat, tolerest, sorbinil, eparetat, and ranirestat) have been developed in the past years but have some side effects and limited efficacy<sup>[23]</sup>. The depletion of plasma antioxidants is a major cause of diabetes related complications such as atherosclerosis and coronary heart disease. From our earlier study it was found that *H. enneaspermus* has significant antioxidant activity, so it can be used to treat such type of complication<sup>[12]</sup>.

In conclusion, the present study reveals that *H. enneaspermus* has significant AR inhibitory potential compared to the isolated compound beta sitosterol. Further, the relationship between phenolic, flavonoid and AR inhibitory activity was also investigated. The ethanolic extracts have been chosen due to its expected flavonoid contents that were reported to have AR inhibitory activity. The drug has shown significant AR inhibitory potential but, further *in vivo* study is necessary to know the exact mechanism of action and safety profile of the *H. enneaspermus* and beta sitosterol.

### Conflict of interest statement

We declare that we have no conflict of interest.

### Acknowledgements

The financial assistance from University Grants Commission, New Delhi, for Dinesh Kumar Patel (Senior Research Fellowship) is greatly acknowledged.

### References

- [1] Brownlee M. Biochemistry and molecular cell biology of diabetic complications. *Nature* 2001; **414**(6865): 813–820.
- [2] Van Zandt MC, Doan B, Sawicki DR, Sredy J, Podjarny AD. Discovery of [3-(4,5,7-trifluoro-benzothiazol-2-ylmethyl)-pyrrolo[2,3-b]pyridin-1-yl] acetic acids as highly potent and selective inhibitors of aldose reductase for treatment of chronic diabetic complications. *Bioorg Med Chem Lett* 2009; **19**(7): 2006–2008.
- [3] Muthenna P, Suryanarayana P, Gunda SK, Petrash JM, Reddy GB. Inhibition of aldose reductase by dietary antioxidant curcumin: Mechanism of inhibition, specificity and significance. *FEBS Lett* 2009; **583**(22): 3637–3642.
- [4] Kato A, Yasuko H, Goto H, Hollinshead J, Nash RJ, Adachi I. Inhibitory effect of rhesinine isolated from *Evodia rutaecarpa* on aldose reductase activity. *Phytomedicine* 2009; **16**(2–3): 258–261.
- [5] Halder N, Joshi S, Gupta SK. Lens aldose reductase inhibiting potential of some indigenous plants. *J Ethnopharmacol* 2003; **86**(1): 113–116.
- [6] Vincent TE, Mendiratta S, May JM. Inhibition of aldose reductase in human erythrocytes by vitamin C. *Diabetes Res Clin Pract* 1999; **43**(1): 1–8.
- [7] Goodarzi MT, Zal F, Malakooti M, Safari MR, Sadeghian S. Inhibitory activity of flavonoids on the lens aldose reductase of healthy and diabetic rats. *Acta Med Iranica* 2006; **44**(1): 41–45.
- [8] Jung SH, Lee YS, Lee S, Lim SS, Kim YS, Shin KH. Isoflavonoids from the rhizomes of *Belamcanda chinensis* and their effects on aldose reductase and sorbitol accumulation in streptozotocin induced diabetic rat tissues. *Arch Pharm Res* 2002; **25**(3): 306–312.
- [9] Hemalatha S, Wahi AK, Singh PN, Chansouria JPN. Anticonvulsant and free radical scavenging activity of *Hybanthus enneaspermus*: A preliminary screening. *Indian J Tradit Knowledge* 2003; **2**(4): 383–388.
- [10] Das S, Dash SK, Padhy SN. Ethno botanical information from Orissa state, India. A review. *J Hum Ecol* 2004; **14**(3): 165–227.
- [11] Tripathy S, Sahoo SP, Pradhan D, Sahoo S, Satapathy DK. Evaluation of anti arthritic potential of *Hybanthus enneaspermus*. *Afr J Pharm Pharmacol* 2009; **3**(12): 611–614.
- [12] Patel DK, Kumar R, Prasad SK, Sairam K, Hemalatha S. Antidiabetic and *in vitro* antioxidant potential of *Hybanthus enneaspermus* (Linn) F. Muell in streptozotocin-induced diabetic rats. *Asian Pac J Trop Biomed* 2011; **1**(4): 316–322.
- [13] Zhu KX, Lian CX, Guo XN, Peng W, Zhou HM. Antioxidant activities and total phenolic contents of various extracts from defatted wheat germ. *Food Chem* 2011; **126**(3): 1122–1126.
- [14] Zhao B, Hall CA. Composition and antioxidant activity of raisin extracts obtained from various solvents. *Food Chem* 2008; **108**(2): 511–518.
- [15] Cheung AK, Fung MK, Lo AC, Lam TT, So KF, Chung SS, et al. Aldose reductase deficiency prevents diabetes-induced blood-retinal barrier breakdown, apoptosis, and glial reactivation in the retina of db/db mice. *Diabetes* 2005; **54**(11): 3119–3125.
- [16] Saraswat M, Muthenna P, Suryanarayana P, Petrash JM, Reddy GB. Dietary sources of aldose reductase inhibitors: Prospects for alleviating diabetic complications. *Asian Pac J Clin Nutr* 2008; **17**(4): 558–565.
- [17] Viswanatha GLS, Vaidya SK, Ramesh C, Krishnadas N, Rangappa S. Antioxidant and antimutagenic activities of bark extract of *Terminalia Arjuna*. *Asian Pac J Trop Med* 2010; **3**(12): 965–970.
- [18] Patel DK, Kumar R, Prasad SK, Hemalatha S. *Pedalium murex* Linn (Pedaliaceae) fruits: A comparative antioxidant activity of its different fractions. *Asian Pac J Trop Biomed* 2011; **1**(5): 395–400.
- [19] Girija K, Lakshman K, Chandrika U, Ghosh SS, Divya T. Anti-diabetic and anti-cholesterolemic activity of methanol extracts of three species of *Amaranthus*. *Asian Pac J Trop Biomed* 2011; **1**(2): 133–138.
- [20] Hsieh P, Huang G, Ho Y, Lin Y, Huang S, Chiang Y, et al. Activities of antioxidants,  $\alpha$ -Glucosidase inhibitors and aldose reductase inhibitors of the aqueous extracts of four *Flemingia* species in Taiwan. *Botanical Stud* 2010; **51**: 293–302.
- [21] Chung YS, Choi YH, Lee SJ, Choi SA, Lee JH, Kim H, et al. Water extract of *Aralia elata* prevents cataractogenesis *in vitro* and *in vivo*. *J Ethnopharmacol* 2005; **101**(1–3): 49–54.
- [22] Tiwari AK, Rao JM. Diabetes mellitus and multiple therapeutic approaches of phytochemicals: Present status and future prospects. *Curr Sci* 2002; **83**(1): 30–38.
- [23] Kim HM, Mok S, Lee JM, Choi CK, Ku J, Park K, et al. Inhibition of aldose reductase from rat lenses by methanol extracts from Korean folk plants. *Nat Prod Sci* 2010; **16**(4): 285–290.