

HOSTED BY



ELSEVIER

Contents lists available at ScienceDirect

Asian Pacific Journal of Tropical Biomedicine

journal homepage: www.elsevier.com/locate/apjtbOriginal article <http://dx.doi.org/10.1016/j.apjtb.2016.04.007>

Knowledge, attitude and practice of healthcare workers concerning Crimean-Congo hemorrhagic fever in Western Iran

Mojtaba Salimi^{1,2}, Abbas Aghaei Afshar^{3*}, Mojtaba Limoei⁴, Soraya Babakhani², Omid Chatrabgoun⁵, Ahmad Ali Hanafi-Bojd¹, Gidiglo Godwin Nutifafa⁶¹Department of Medical Entomology & Vector Control, School of Public Health, Tehran University of Medical Sciences (TUMS), Tehran, Iran²Department of Health Promotion and Network Expansion, Health Deputy, Kermanshah University of Medical Sciences, Kermanshah, Iran³Leishmaniasis Research Center, Kerman University of Medical Sciences, Kerman, Iran⁴Departments of Public Health and Health Research Center, Kermanshah University of Medical Sciences, Kermanshah, Iran⁵Department of Mathematical, Statistics and Probability, Malayer University, Iran⁶Department of Medical Entomology & Vector Control, School of Public Health, International Campus, Tehran University of Medical Sciences, Tehran, Iran

ARTICLE INFO

Article history:

Received 15 Jan 2016

Received in revised form 12 Feb, 2nd revised form 19 Feb 2016

Accepted 23 Mar 2016

Available online 22 Apr 2016

Keywords:

Crimean-Congo hemorrhagic fever

Healthcare worker

Knowledge

Iran

ABSTRACT

Objective: To determine the knowledge, attitude and practice of healthcare workers in Kermanshah Province about Crimean-Congo hemorrhagic fever (CCHF).**Methods:** This study was conducted in 2014 on healthcare personnel in different job categories including physicians, nurses, midwives, laboratory staff and network health staff of Kermanshah Province by direct interview.**Results:** A total of 367 respondents who had more than 5 years of experience in their jobs were interviewed. Among them 91% of physicians and nurses, 97% of midwives and health workers and 96% of laboratory staff stated that they had not been confronted with CCHF patients so far. Regarding knowledge, 76% of physicians, 78% of nurses, 77% of midwives and 58% of laboratory staff believed that the disease is remediable. Most of the interviewed participants stated that the disease pertains to people who are in close contact with domestic animals, but they did not consider their own occupations as one of the risk factors. More than 70% of the respondents believed that the disease may exist in the province or their work field. Generally, the knowledge about CCHF was inadequate, with nurses having the lowest level of knowledge.**Conclusions:** Knowledge of Kermanshah healthcare staff about CCHF was poor, especially nurses in a high risk job category. Therefore, it is necessary to conduct specific training programs for the disease identification, transmission, prevention, and treatment as well as the use of personal protection and safety devices.

1. Introduction

Crimean-Congo hemorrhagic fever (CCHF) is an acute hemorrhagic fever caused by segmented, negative-stranded RNA virus belonging to the family Bunyaviridae, genus *Nairovirus*, which is the second most widespread of all medically important arboviruses, after dengue virus. Basically, this disease is transmitted to humans by domestic animals and bite of an infected tick or via aerosol generated from infected animals' excreta. Human to human transmission occurs following contact with an infected person's blood, tissue or fluid discharge. The vectors of this arthropod-borne disease are generally hard ticks of Ixodidae family,

*Corresponding author: Abbas Aghaei Afshar, Leishmaniasis Research Center, Kerman University of Medical Sciences, Kerman, Iran.

Tel: +98 341 3224616

E-mail: afshara2@yahoo.com

The study protocol was approved by the Ethic Committee of Kermanshah University of Medical Sciences which approved the investigations. Informed written consent was obtained from all participants in this study. Confidentiality on the content of the records was kept by the investigators and information was only utilized for the research purpose.

Foundation Project: Supported by Kerman University of Medical Sciences, Kerman, Iran (Grant No. 94251).

Peer review under responsibility of Hainan Medical University. The journal implements double-blind peer review practiced by specially invited international editorial board members.

including some species of *Rhipicephalus*, *Boophilus*, *Dermacentor* and *Hyalomma* (in particular *Hyalomma marginatum*). Some species of *Argas* and *Ornithodoros* in Argasidae family have been reported to be infected [1–5]. The virus of the disease can be transmitted via transovarial transmission through the eggs of tick vectors to their next generation and can pass through different stages of their life cycle. Therefore, if a tick bites an infected vertebrate, it will be infected in all its life and can transmit the disease to humans by bites.

Climate has also been found as important predictor of CCHF risk; areas regularly experiencing long period of low rainfall and humidity were associated with increased incidence of CCHF in Iran and Senegal, and higher temperatures were indicators of CCHF occurrence in Turkey, Bulgaria, and Iran [6,7]. Outbreak of the disease occurs in warm seasons when ticks are more active and questing hosts for blood feeding. Most cases were reported between April and September. Increasing mean temperature, normalized difference vegetation index, savannah-type land coverage or habitat fragmentation increased significantly the incidence of CCHF in the CCHF-affected areas [7]. Climate, environmental factors and human behavior that may influence CCHF epidemiology and spread should be further studied. Facilitated tick reproduction and global warming which are mainly attributed to climatic changes should be considered in the potential changing epidemiology of CCHF [8].

Hosts of the disease include a range of domestic and wild animals such as cattle, sheep and goat. The birds are resistant to infection but ostrich is sensitive. In endemic areas where human infection is the source, the disease is of high prevalence. There are no clinical symptoms in animals, and this increases the risk of infection in humans when slaughter the infected animal or contact animal's skin or carrion shortly after slaughtering. CCHF has been ranked as one of the most important nosocomial infections and hospitalized patients could infect others, for which reason they must be isolated. CCHF virus is on the US list of select agent considered to have bioterrorism potential because of its association with rapid-onset of hemorrhagic fever and 30% case fatality rate [5,8–10]. There is considerable difference among mortality rates of the disease in different parts of the world. The difference is not only related to that of diagnostic services-medical care, but also to virulence and pathogenicity of different virus strains across the world. For example, the disease is milder in South Russia with a mortality rate of 5%–10%, but in South Africa and Middle East the mortality rates were reported as 35% and 35%–50%, respectively. It should be noted that the mortality was higher in hospital epidemics [11]. The World Health Organization has listed CCHF among the emerging diseases for which control and prevention measures should be renewed and intensified. According to our consensus measure, the five countries currently having the strongest evidence for presence of CCHF are Turkey, Iran, Afghanistan, Tajikistan, and Pakistan [12,13].

The disease has been reported in several countries from Africa (Senegal, Nigeria, South Africa, Kenya, Tanzania, Ethiopia, Mauritania, Congo, and Uganda), Europe (Bulgaria, Turkey, Albania, Kosovo, Greece, Yugoslavia, Hungary, Russia and Ukraine) and Asia (Iraq, Pakistan, India, Afghanistan, Oman, Iran, China and United Arab Emirates) [13,14], and imported CCHF has never been reported in Northern Europe or in America [15]. CCHF in Iran was first reported in 1970, when virus antibodies were identified in the sera of 45 sheep sent from Tehran to Moscow [16]. After that antibodies of the virus were detected in 13% of people who were living in Caspian Sea littoral from East

Azerbaijan Province [17]. Mortality rate approaching 20% in 2 000 remarkably dropped to 6% in 2007. CCHF, however, is an emerging disease in the country. The disease exists in almost all parts of Iran according to the epidemiological evidences and confirmed cases [18]. In recent decades, confirmed human cases that resulted in death were reported in provinces of Sistan-va-Baluchestan, Khuzestan, Chaharmahal-va-Bakhtiari, Azerbaijan-e-Gharbi, Bushehr, Yazd, Kerman, Tehran, Esfahan, Golestan, Fars, Qom and Khorasan in Iran [6,14,18].

CCHF is an occupational disease for butchers, slaughter house workers, livestock workers, animal husbandry workers, veterinarians and healthcare workers who belong to the group of people at high risk of the disease in the endemic areas of CCHF [12]. In a survey conducted in south of Iran, livestock was identified as the source of infection and CCHF outbreak occurred via two routes: contact with tissue and blood of infected livestock, as well as nosocomial transmission [16]. Hence, the knowledge of people who are in contact with the disease should be evaluated periodically.

Kermanshah Province is important from CCHF point of view because of cattle trafficking with infected areas, illegal slaughtering in the area and confirmed cases of disease and death in the neighboring provinces [11,12,19]. Therefore, it is necessary to evaluate the knowledge of at-risk occupational groups. The aim of this study was to evaluate the knowledge, attitude and practice of the healthcare staff as one of the high risk occupational groups of people towards CCHF.

2. Materials and methods

Kermanshah Province, located at the extreme west of Iran shares border with Iraq, and is located at the coordinates of 33.36–35.15° N and 45.24–48.30° E (Figure 1). The weather is moderate and both cold and warm regions can be found in the province. Annual mean temperature is reported as 15.4 °C and the mean annual rainfall is recorded between 300 and 800 mm. The capital city is Kermanshah and 13 other cities are located in this province. There are many livestock farms in this province and CCHF virus has been detected from ticks and patients in recent years.



Figure 1. Map of Iran and location of Kermanshah Province in Western Iran.

2.1. Study population and sample size

This descriptive cross-sectional study was conducted on healthcare staff of Kermanshah Province in 2014 to determine their knowledge, attitude and practice concerning CCHF. A total of 367 healthcare workers were selected using cluster random sampling from different cities of the province for interview, based on the number of people employed in each occupational group.

2.2. Data collection and analysis

Data were collected by the use of a pre-designed questionnaire and interviews were conducted on healthcare staff in different occupational categories (physicians, nurses, midwives, laboratory staff and health network personnel). Questions covered demographic information, as well as knowledge (7 questions), attitudes (5 questions) and practice (4 questions) of the respondents about CCHF. Reliability of the questionnaire was examined based on a few staff members in occupational groups of health service providers. The collected data were analyzed using SPSS 16.0, Chi-square test, One-way ANOVA and least significant difference tests to find the correlation of knowledge, attitude and practice of the respondents with their job categories.

2.3. Ethical consideration

The study protocol was approved by Ethic Committee of Kermanshah University of Medical Sciences which approved the investigations. Informed written consent was obtained from all participants in this study. Confidentiality on the content of the records was kept by the investigators and information was only utilized for the research purpose.

3. Results

A total of 367 individuals were interviewed including 85 physicians, 116 nurses, 65 midwives, 72 laboratory staff and 29 health network personnel. Among the interviewed population, 49% of physicians, 94% of nurses, 17% of midwives and 91% of laboratory staff were employed in hospitals and other health centers. Academic education is the main source of knowledge acquisition about CCHF for physicians (62%), midwives (53%) and health staff (33%); job training and press and media are the main sources of knowledge acquisition about CCHF for nurses (45%) and laboratory workers (39%), respectively (Figure 2).




Figure 2. Main source of knowledge acquisition about CCHF in healthcare workers.

Regarding knowledge of the studied community, 76% of physicians, 78% of nurses, 77% of midwives and 58% of laboratory staff believed that CCHF is treatable. The level of knowledge about transmission, reservoir host, treatment, dealing with patients and personal protection in physicians, nurses, midwives and laboratory staff in hospitals was lower than that of staff in health centers. Many occupational groups particularly in the private sector have not had enough information on their occupational hazards yet. By using Chi-square test, there was no significant relationship between job and level of knowledge ($P > 0.05$), but with respect to One-way ANOVA there was significant relationship between the levels of knowledge in different occupational groups. This difference was significant between physician and nurse, nurse and laboratory worker, as well as nurse and health workers ($P < 0.05$). The knowledge and awareness of occupational groups were compared as binary by least significant difference test. It was found that there is a significant difference between occupational groups (nurses and physicians, and laboratory staff and nurses). Level of knowledge in laboratory staff and physicians was higher than that in nurses (Table 1). In this study, there was no significant relationship between the place of work (urban or rural) and the level of their knowledge ($P = 0.08$). Concerning attitude, overall 88% of physicians, 93% of nurses, 97% of midwives and 92% of laboratory staff believed that CCHF patients must be isolated in the hospital (Table 1). About the prevention, 95% of physicians, 93% of nurses, 92% of midwives and 96% of laboratory staff believed that prevention is essential for CCHF and control can avoid or minimize the exposure to the virus.

Table 1

Results of the interview with various occupational groups in Kermanshah Province, Iran. %.

Interview questions	Positive response in different occupational groups				
	Physician	Nurse	Midwife	Laboratory staff	Health staff
Have you ever encountered a CCHF patient?	91	91	97	96	3
General information about the disease	81	80	85	90	89
Knowledge about the transmission routes of the disease	89	85	94	99	100
Knowledge about the etiology of CCHF	86	66	71	83	83
Do you believe that the disease is treatable?	76	78	77	58	86
CCHF can be considered as an occupational disease	89	89	92	94	97
Knowledge about the host reservoir(s)	51	53	53	61	48
Patients must be isolated in hospital	88	93	97	92	79
Treatment of the patient's family is necessary	46	82	47	88	48
Death occurs due to lack of treatment	94	88	85	96	97
Identifying risk factors	95	93	92	96	93

(continued on next page)

Table 1 (continued)

Interview questions	Positive response in different occupational groups				
	Physician	Nurse	Midwife	Laboratory staff	Health staff
Personal protection	80	90	74	81	69
The need to use preventive medications by staff	64	71	70	71	72
Inter-sector collaborations are required	58	52	56	69	72
Prevention measures are necessary	95	93	92	96	93
Presence of the disease in their working areas	71	76	79	69	72

4. Discussion

Transmission of CCHF through infected secretions and blood has put it as one of the most important nosocomial infections, and reports of outbreak of the disease in the hospital have ranked it as one of the main sources of infection among service providers and staff in health centers. Thus helping the healthcare providers to prevent the spread of the disease in a community and hospital setting has significant roles. The first priority and necessity in this regard is to assess the knowledge, attitude and practice of related staff as one of the high risk groups in order to find their weaknesses, so that essential steps can be taken to overcome the disease by designing a proper educational plan with assistance from related authorities at different educational and executive levels. This study refers to the role of education in promoting the knowledge of healthcare staff in the face of CCHF. According to the results, there was no significant relationship between job and level of knowledge. The results of this study confirmed those of previous studies conducted in other areas [13,16,20–24].

Despite results of Gozel *et al.* showing that CCHF virus is not easily transmitted from person to person through respiratory or physical contact, use of equipments to prevent transmission of this disease, including gloves, face masks, face visors and box coats, should be introduced to emergency room health care workers, and encouragement should be provided for using these equipments [15]. In this study we identified the use of personal protective equipment in some occupational groups, particularly those have direct contact with the disease. Some occupational groups, especially nurses, have direct contact with hospitalized patients. Therefore, their low level of knowledge about the disease is regarded as one of the important risk factors to the spread of this disease in the community. Another important point in these results is that most occupational groups did not consider their occupations as risk factors of the disease, and treatment with ribavirin in suspected cases and post-exposure prophylaxis for healthcare workers who potentially exposed to CCHF virus should be considered.

We found that level of knowledge in laboratory staff and physicians was higher than that in nurses. This finding is contrary to other studies which have assessed the knowledge of nurses at a high level [20,25,26]. Some other studies on healthcare workers concluded that educational programs, especially for nurses, plays a major role in prevention of disease

transmission [12,27]. In Kermanshah Province, due to insufficient training of staff especially in the health sector where most of the selected nurses were from, their knowledge about CCHF was inadequate, and educational programs must be codified for them. Here, the role of insufficient job training, especially for high-risk groups is visible and should be considered by the relevant authorities. In conclusion, it is recommended that specific training courses be conducted for healthcare staff to recognize early symptoms of CCHF and include the disease in their initial differential diagnosis, thereby ensuring early detection of the disease. Strengthening national and regional surveillance is necessary to provide a clearer feature of CCHF in its geographically distributed area. Isolation facilities should be available in hospitals and strict adherence to safety measures must be learned and practiced by healthcare workers.

Conflict of interest statement

The authors declare that there is no conflict of interest.

Acknowledgments

The authors are grateful to president of the health center of Kermanshah for his support as well as personnel of Kermanshah University of Medical Science for cooperation in this study. This study has received financial grant from Kerman University of Medical Sciences (Grant No. 94251).

References

- [1] Duh D, Saksida A, Petrovec M, Ahmeti S, Dedushaj I, Panning M, et al. Viral load as predictor of Crimean-Congo hemorrhagic fever outcome. *Emerg Infect Dis* 2007; **13**(11): 1769-72.
- [2] Messina JP, Pigott DM, Golding N, Duda KA, Brownstein JS, Weiss DJ, et al. The global distribution of Crimean-Congo hemorrhagic fever. *Trans R Soc Trop Med Hyg* 2015; **109**(8): 503-13.
- [3] Tahmasebi F, Ghiasi SM, Mostafavi E, Moradi M, Piazak N, Mozafari A, et al. Molecular epidemiology of Crimean-Congo hemorrhagic fever virus genome isolated from ticks of Hamadan Province of Iran. *J Vector Borne Dis* 2010; **47**(4): 211-6.
- [4] Telmadarraiy Z, Ghiasi SM, Moradi M, Vatandoost H, Eshraghian MR, Faghihi F, et al. A survey of Crimean-Congo haemorrhagic fever in livestock and ticks in Ardabil Province, Iran during 2004–2005. *Scand J Infect Dis* 2010; **42**(2): 137-41.
- [5] Wölfel R, Paweska JT, Petersen N, Grobelaar AA, Leman PA, Hewson R, et al. Virus detection and monitoring of viral load in Crimean-Congo hemorrhagic fever virus patients. *Emerg Infect Dis* 2007; **13**(7): 1097-100.
- [6] Chinikar S. An overview of Crimean-Congo hemorrhagic fever in Iran. *Iran J Microbiol* 2009; **1**(1): 7-12.
- [7] Vescio FM, Busani L, Mughini-Gras L, Khoury C, Avellis L, Taseva E, et al. Environmental correlates of Crimean-Congo haemorrhagic fever incidence in Bulgaria. *BMC Public Health* 2012; **12**: 1116.
- [8] Maltezou HC, Andonova L, Andraghetti R, Bouloy M, Ergonul O, Jongejan F, et al. Crimean-Congo hemorrhagic fever in Europe: current situation calls for preparedness. *Euro Surveill* 2010; **15**(10): 19504.
- [9] Gozel MG, Bakir M, Oztop AY, Engin A, Dokmetas I, Elaldi N. Investigation of Crimean-Congo hemorrhagic fever virus transmission from patients to relatives: a prospective contact tracing study. *Am J Trop Med Hyg* 2014; **90**(1): 160-2.
- [10] Vanhomwegen J, Alves MJ, Županc TA, Bino S, Chinikar S, Karlberg H, et al. Diagnostic assays for Crimean-Congo hemorrhagic fever. *Emerg Infect Dis* 2012; **18**(12): 1958-65.

- [11] Sisman A. Epidemiologic features and risk factors of Crimean-Congo hemorrhagic fever in Samsun Province, Turkey. *J Epidemiol* 2013; **23**(2): 95-102.
- [12] Yolcu S, Kader C, Kayipmaz AE, Ozbay S, Erbay A. Knowledge levels regarding Crimean-Congo hemorrhagic fever among emergency healthcare workers in an endemic region. *J Clin Med Res* 2014; **6**(3): 197-204.
- [13] Ali Z, Kumar R, Ahmed J, Ghaffar A, Mureed S. Knowledge, attitude and practice of Crimean-Congo hemorrhagic fever among rural population of Baluchistan, Pakistan. A public health nutritional assessment of elderly in Islamabad: a mixed method study. *J Publ Health* 2013; **3**(4): 11-3.
- [14] Ansari H, Shahbaz B, Izadi S, Zeinali M, Tabatabaee SM, Mahmoodi M, et al. Crimean-Congo hemorrhagic fever and its relationship with climate factors in southeast Iran: a 13-year experience. *J Infect Dev Ctries* 2014; **8**(6): 749-57.
- [15] Gozel MG, Dokmetas I, Oztop AY, Engin A, Elaldi N, Bakir M. Recommended precaution procedures protect healthcare workers from Crimean-Congo hemorrhagic fever virus. *Int J Infect Dis* 2013; **17**(11): e1046-50.
- [16] Izadi S, Salehi M, Holakouie-Naieni K, Chinikar S. The risk of transmission of Crimean-Congo hemorrhagic fever virus from human cases to first-degree relatives. *Jpn J Infect Dis* 2008; **61**(6): 494-6.
- [17] Jabbari A, Tabasi S, Abbasi A, Alijanpour E. Crimean-Congo hemorrhagic fever: treatment and control strategy in admitted patients. *Casp J Intern Med* 2012; **3**(2): 443-4.
- [18] Chinikar S, Ghiasi SM, Moradi M, Goya MM, Shirzadi MR, Zeinali M, et al. Geographical distribution and surveillance of Crimean-Congo hemorrhagic fever in Iran. *Vector Borne Zoonotic Dis* 2010; **10**(7): 705-8.
- [19] Majeed B, Dicker R, Nawar A, Badri S, Noah A, Muslem H. Morbidity and mortality of Crimean-Congo hemorrhagic fever in Iraq: cases reported to the national surveillance system, 1990-2010. *Trans R Soc Trop Med Hyg* 2012; **106**(8): 480-3.
- [20] Ozer A, Miraloglu M, Ekerbicer HC, Cevik F, Aloglu N. Knowledge levels about Crimean-Congo hemorrhagic fever among midwifery and nursing students in Kahramanmaraş, Turkey. *Southeast Asian J Trop Med Public Health* 2010; **41**(1): 77-84.
- [21] Sharifi-Mood B, Mardani M, Keshkar-Jahromi M, Rahnavardi M, Hatami H, Metanat M. Clinical and epidemiologic features of Crimean-Congo hemorrhagic fever among children and adolescents from southeastern Iran. *Pediatr Infect Dis J* 2008; **27**(6): 561-3.
- [22] Sheikh NS, Sheikh AS, Sheikh AA. Knowledge, attitude and practices regarding Crimean-Congo haemorrhagic fever among healthcare workers in Balochistan. *J Ayub Med Coll Abbottabad* 2004; **16**(3): 39-42.
- [23] Sharifinia N, Rafinejad J, Hanafi-Bojd AA, Biglarian A, Chinikar S, Baniardalani M, et al. Knowledge and attitudes of the rural population and veterinary and health personnel concerning Crimean-Congo hemorrhagic fever in Western Iran in 2012. *Fla Entomol* 2013; **96**(3): 922-8.
- [24] Yilmaz R, Ozcetin M, Erkorkmaz U, Ozer S, Ekici F. Public knowledge and attitude toward Crimean Congo hemorrhagic fever in Tokat Turkey. *Iran J Arthropod Borne Dis* 2009; **3**(2): 12-7.
- [25] Çilingiroğlu N, Temel F, Altıntaş H. Public's knowledge, opinions and behaviors about Crimean-Congo hemorrhagic fever: an example from Turkey. *Kafkas Univ Vet Fak Derg* 2010; **16**: S17-22.
- [26] Mofleh JA, Ashgar RJ, Kakar RS, Zaidi SZ, Kazi BM. Nosocomial outbreak of Crimean-Congo hemorrhagic fever in Holy Family Hospital, Rawalpindi, Pakistan, 2010. *J Public Health Epidemiol* 2013; **5**(4): 173-7.
- [27] Taghdisi M, Sasooli H, Ramezani A. A study of level the nurse's knowledge of the CCHF disease in the Sistan. *Zahedan J Res Med Sci* 2012; **13**(1): 65.