



Contents lists available at ScienceDirect

Journal of Acute Disease

journal homepage: www.jadweb.org



Document heading doi: 10.1016/S2221-6189(13)60126-2

A review: Anti diabetic medicinal plants used for diabetes mellitus

G Arumugam¹, P Manjula², N Paari^{2*}

¹Department of Biochemistry, Adhiparasakthi College of Science, Kalavai, Vellore

²Dr. N. Paari's Diabetic Centre, MKB Nagar, Chennai-600039

ARTICLE INFO

Article history:

Received 14 May 2013

Received in revised form 22 May 2013

Accepted 29 May 2013

Available online 20 September 2013

Keywords:

Diabetes mellitus

Ayurveda

Medicinal plants

Hypoglycemic

ABSTRACT

Aim of the present study is evaluated various medicinal plants used for antidiabetic activity. Diabetes mellitus is one of the most common non-communicable diseases globally. It is the fourth leading causes of death in the most developed countries and there in substantial evidenced that it in epidemic in many developing and newly industrialized nations. This posing a serious threat to be met within 21st century. Since ancient time plants have been exemplary source of medicine. Ayurveda and other Indian literature mentioned the used of plants in treatment of various ailments. Out of an estimated 250 000 higher plants, less than 1% have been screened pharmacologically and very few in regard to diabetes mellitus. Systematic studies on the folklore medicinal plants that combat diabetes mellitus are scanty.

1. Introduction

Diabetes mellitus is a common and very prevalent disease affecting the citizens of both developed and developing countries. It is estimated that 25% of the world population is affected by this disease. Diabetes mellitus is caused by the abnormality of carbohydrate metabolism which is linked to low blood insulin level or insensitivity of target organs to insulin^[1]. Despite considerable progress in the treatment of diabetes by oral hypoglycemic agents, search for newer drugs continues because the existing synthetic drugs have several limitations. The herbal drugs with antidiabetic activity are yet to be commercially formulated as modern medicines, even though they have been acclaimed for their therapeutic properties in the traditional systems of medicine^[2]. Type 2 diabetes usually occurs in obese individuals and is associated with hypertension and dyslipidemia. Thus the treatment aims to reduce insulin resistance and to stimulate insulin secretion. Diabetes is a metabolic disorder where in human body does not produce or properly use insulin, a hormone that is

required to convert sugar, starches, and other food into energy. Diabetes mellitus is characterized by constant high levels of blood glucose (sugar). Human body has to maintain the blood glucose levels at a very narrow range which is done with insulin and glucagon. The function of glucagon is causing the liver to release glucose from its cells into the blood for the production of energy. Type 1 Diabetes leads to inability to release insulin results in low rates of glucose uptake into muscles and adipose tissue^[3]. Traditional medicine (herbal) is used for treatment of diabetes in developing countries where the cost of conventional medicines is a burden to the population^[4]. Despite the introduction of hypoglycemic agents from natural and synthetic sources, diabetes and its secondary complications continue to be a major medical problem. Many indigenous Indian medicinal plants have been found to be useful to successfully manage diabetes. One of the great advantages of medicinal plants is that these are readily available and have very low side effects. Plants have always been an exemplary source of drugs and many of the currently available drugs have been derived directly or indirectly from them.

The ethnobotanical information reports about 800 plants that may possess antidiabetic potential^[5]. Several herbs have shown antidiabetic activity when assessed

*Corresponding author: Dr. N. Paari, Diabetologist, Dr. N. Paari's Diabetic Centre, No; 28; 1st Main Road, MKB. Nagar, Chennai-600039.

Tel: 07845345239

E-mail: bioaru.g@gmail.com

using presently available experimental techniques[6]. This review article enumerates some medicinal plants possessing antidiabetic activity and elucidating their mechanisms of action such as *Brassica juncea* (*B. juncea*), *Combretum micranthum* (*C. micranthum*), *Elephantopus scaber* (*E. scaber*), *Gymnema sylvestre* (*G. sylvestre*), *Liriope spicata* (*L. spicata*), *Parinari excelsa* (*P. excelsa*), *Ricinus communis* (*R. communis*), *Sarcopoterium spinosum* (*S. spinosum*), *Smallanthus sonchifolius* (*S. sonchifolius*), *Swertia punicea* (*S. punicea*), *Vernonia anthelmintica* (*V. anthelmintica*) etc. and method of experiment on animals and therapeutic efficiency of plant extracts were exploited. Some of the important anti-diabetic potential herbal plants sources are given in the Table 1.

2. Antidiabetic effect of folklore medicinal plants

2.1. *Brassica juncea*

It is commonly used spice in various food items in Tamilnadu. *B. juncea* is a traditional medicinal plant which belongs to family Cruciferae. *B. juncea* aqueous seed extract has a potent hypoglycemic activity which was investigated in STZ induced diabetic male albino rat. Doses which have hypoglycemic activity was reported as 250, 350, 450 mg/kg[7].

2.2. *Eugenia jambolana*

Eugenia jambolana (*E. jambolana*) popularly known as Jamun or Indian blackberry has been indicated in

Table 1

Medicinal plants having antidiabetic activity.

S.No	Plant name	Family	Parts used	Type of extract	Activity	References
1	<i>Alangium lamarkii</i>	Alangiaceae	Leaves	Alcoholic	Antidiabetic	[15]
2	<i>Albizia odoratissima</i>	Mimosaceae	Bark	Methanol	Antidiabetic	[19]
3	<i>Axonopus compressus</i>	Poaceae	Leaves	Methanol	Antidiabetic	[13]
4	<i>Berberis vulgaris</i>	Berberidaceae	Root	Aqueous	Hypoglycaemic	[14]
5	<i>Brassica juncea</i>	Cruciferae	Seed	Aqueous	Hypoglycemic	[7]
6	<i>Caesalpinia digyna</i>	Fabaceae	Root	Methanol	Antidiabetic	[10]
7	<i>Catharanthus roseus</i>	Apocynaceae	Leaf	Methanol	Hypoglycemic	[16]
8	<i>Centaurium erythraea</i>	Gentianaceae	Leaf	Aqueous	Antidiabetic	[17]
9	<i>Chaenomeles sinensis</i>	Rosaceae	Fruits	ethyl acetate	Antidiabetic	[18]
10	<i>Cocos nucifera</i>	Arecaceae	Leaf	hydro-methanol	Antihyperglycemic	[20]
11	<i>Costus speciosus</i>	Costaceae	rhizome	hexane	Antidiabetic	[21]
12	<i>Cyclocarya paliurus</i>	Cyclocaryaceae	Bark	Aqueous, PE, chloroform, ethyl acetate & n-butanol	Hypoglycemic	[22]
13	<i>Dillenia indica</i>	Dilleniaceae	Leaves	Methanolic	Antidiabetic	[23]
14	<i>Embelia ribes</i>	Myrsinaceae	Berries	Hexane	Antidiabetic	[24]
15	<i>Hybanthus enneaspermus</i>	Violaceae	Whole plant	Alcoholic	Antidiabetic	[25]
16	<i>Lippa nodiflora</i>	Verbenaceae	Whole plant	Methanol	Antidiabetic and Hypolipidemic	[26]
17	<i>Lithocarpus polystachyus</i>	Fagaceae	Leaves	Ethanol & Aqueous	Hypoglycemic	[27]
18	<i>Marrubium vulgare</i>	Lamiaceae	Aerial part	Methanol	Hyperglycemia and dyslipidemia	[28]
19	<i>Ocimum sanctum</i>	Lamiaceae	Aerial part	Hydroalcoholic	Antidiabetic	[29]
20	<i>Opuntia streptacantha</i>	Cactaceae	Leaves	Ethanol	Antihyperglycemia	[30]
21	<i>Psidium guajava</i>	Myrtaceae	Fruits	Ethanol	Antihyperglycemic	[31]
22	<i>Semecarpus anacardium</i>	Anacardiaceae	nut	Milk	Antidiabetic	[32]
23	<i>Prosopis glandulosa</i>	Fabaceae	Whole plant	Gelatine/Jelly	Antidiabetic	[33]
24	<i>Ophiopogon japonicus</i>	Asparagaceae	Root	Ethanol	Hypoglycemic	[34]
26	<i>Setaria italica</i>	Poaceae	Seed	Aqueous	Antihyperglycemic	[35]
25	<i>Solanum torvum</i>	Solanaceae	Fruit	Methanol	Antihyperglycemic	[36]
26	<i>Cassia auriculata</i>	Caesalpiniaceae	Leaves	Aqueous	Antihyperglycemic	[37]
27	<i>Zygophyllum album</i>	Zygophyllaceae	Whole plant	Ethanol	Antidiabetic	[38]
28	<i>Vitex negundo</i>	Lamiaceae	Leaves	Methanol	Antihyperglycemic	[39]
29	<i>Viscum schimperi</i>	Viscaceae	aerial parts	Methanolic	Antihyperglycemic & Hypolipidaemic	[40]
30	<i>Symplocos cochinchinensis</i>	Symplocaceae	Leaves	Hexane	Antidiabetic	[41]
31	<i>Enicostemma littorale</i>	Gentianaceae	Whole plant	aqueous	Antidiabetic	[42]
32	<i>Vaccinium arctostaphylos</i>	Ericaceae	Fruit	Ethanol	antidiabetic	[43]
33	<i>Solanum xanthocarpum</i>	Solanaceae	Leaves	Aqueous and Methanol	Antihyperglycemic	[44]

Ayurveda, an ancient system of Indian medicine, for use in DM. In accordance to its claimed anti-diabetic effect in traditional medicine, *E. jambolana* has been reported to have hypoglycemic effects both in experimental models and clinical studies^[8].

2.3. *Coccinia grandis*

Hypoglycemic activity was evaluated in alcoholic extracts of *Coccinia grandis* (*C. grandis*) leaves. Alcoholic extract 600 mg/kg bw was injected orally to mice. Oral administration of alcoholic extract of leaves of *C. grandis* showed significant hypoglycemic effect on blood glucose level in normal fasted rats^[9].

2.4. *Alangium lamarckii*

Antidiabetic effect of alcoholic extract of *Alangium lamarckii* (*A. lamarckii*). Alcoholic leaves extract 250 and 500 mg/kg bw was used for these studies. *A. lamarckii* have significant antidiabetic activity in STZ-nicotinamide induced diabetic rat^[10].

2.5. *Albizia odoratissima*

Antidiabetic effect of methanolic bark extract of *Albizia odoratissima* (*A. odoratissima*) in alloxan induced diabetic mice. The methanolic extracts were fed to the animals at a dosage of 250 and 500 mg/kg body weight. The significant reduced in the levels of serum cholesterol, triglycerides, SGOT, SGPT, alkaline phosphatase and decrement of total proteins in alloxan induced albino mice^[11].

2.6. *Artemis sphaerocephala* Krasch

Antioxidant effect of *Artemis sphaerocephala* (*A. sphaerocephala*) gum on STZ induced diabetic rat. Levels of serum and liver tissue thiobarbituric acid reactive substances (TBARS) and +OH were increased in STZ induced rat. The activity levels of liver and serum superoxide dismutase were decreased. After administration of extract of *A. sphaerocephala*, levels of TBARS and +OH were decreased in serum and liver tissue. The significant increments in the levels of liver and serum SOD. *A. sphaerocephala* is very good antioxidant activity^[12].

2.7. *Axonopus compressus*

The anti-diabetic effect of the methanolic leaf extract of the plant. Diabetes was induced in the rats by injection of alloxan. Methanolic leaves extract 250, 500 and 1 000 mg/kg bw was used for these studies. Methanolic leaf extract of *Axonopus compressus* (*A. compressus*) at all

the doses (250, 500 and 1 000 mg/kg) were significant reduction (by 31.5%, 19.8% and 24.5%) of the blood glucose levels in the diabetic rats when compared to the control group. *A. compressus* may possess very good antidiabetic property^[13].

2.8. *Berberis vulgaris*

Hypoglycaemic effect of *Berberis vulgaris* (*B. vulgaris*) L. in streptozotocin-induced diabetic rats *B. vulgaris* a traditional medicinal plant which belongs to family Berberidaceae. The results indicated that water extract and saponins shows significant hypoglycemic effect. The serum cholesterol and serum triglycerides levels were significantly increased^[14].

2.9. *Caesalpinia digyna*

Antidiabetic effect of bergenin from the roots of *Caesalpinia digyna* (*C. digyna*). The levels of plasma total cholesterol (TC), triglycerides (TG) and LDL-C were significantly increased, whereas levels of (HDL-C) were significantly decreased in diabetic rats when compared to control. After administration of bergenin (10 mg/kg; p.o.) the lipid profile were significantly increased when compared with that of glibenclamide (10 mg/kg; p.o.). The activity levels of antioxidant enzymes such as SOD and Cat were decreased. The level of TBARS was significantly increased in diabetic rat compared to control rat. The administration of bergenin (10 mg/kg; p.o.) significantly increased the SOD and CAT respectively and reduced TBARS level. Bergenin is very good antidiabetic properties^[15].

2.10. *Catharanthus roseus*

Hypoglycemic effect of the methanolic leaf extract of *Catharanthus roseus* (*C. roseus*) in alloxan induced diabetic rats. The levels of blood glucose were significantly decreased when compared with Control rat. The blood glucose lowering effect of *C. roseus* methanolic extract was more pronounced than Glibenclamide and Metformin^[16].

2.11. *Centaurium erythrea*

Diabetes was induced by a single dose of STZ (65 mg/kg) administered by intraperitoneal way. The oxidative stress was measured by tissue MDA. The estimation of pancreas antioxidant enzymes such as superoxide dismutase (SOD), catalase (CAT), and glutathione peroxidase (GPx). A significant decrement in the levels of pancreas tissue TBARS was recorded in diabetic treated rats when compared to that of normal animals. The activity levels of pancreas antioxidant defense enzymes viz. SOD, CAT,

GPx and GST were significantly increased in the diabetic treated animals. Antioxidant effect of the aqueous leaf extract of *Centaurium erythraea* (*C. erythraea*)^[17].

2.12. *Chaenomeles sinensis*

Ethyl acetate fraction of *Chaenomeles sinensis* (*C. sinensis*) (Thouin) Koehne fruits is very good Antidiabetic effect. *Chaenomeles sinensis* is belongs to family Rosaceae. Doses which have antidiabetic activity were reported as 50 and 100 mg/kg body weight^[18].

3. Conclusion

In this review we discussed about folklore medicinal plants for the treatment of Diabetes mellitus. Folklore medicinal plants are mostly used for rural areas; because the availability of lavish amount of medicinal plants those areas. Therefore, treating diabetes mellitus with plant derived compounds which are accessible and do not require laborious pharmaceutical synthesis seems highly attractive. In the present review an attempt has been made to investigate the antidiabetic medicinal plants and may be useful to the health professionals, scientists and scholars working in the field of pharmacology and therapeutics to develop antidiabetic drugs.

Conflict of interest statement

We declare that we have no conflict of interest.

References

- [1] Maiti R, Jana D, Das UK, Ghosh D. Antidiabetic effect of aqueous extract of seed of *Tamarindus indica* in streptozotocin induced diabetic rats. *J Ethnopharmacol* 2004; **92**: 85–91.
- [2] Wadkar KA, Magdum CS, Patil SS, Naikwade NS. Antidiabetic potential and Indian medicinal plants. *J Herbal Med Toxicol* 2008; **2**: 45–50.
- [3] AL Lehninger, DL Nelson, MM Cox. *Principle of Biochemistry*. New York: Worth Publishers; 2010.
- [4] Saravanan G, Pari L. Hypoglycaemic and antihyperglycaemic effect of *Syzygium cumini* bark in streptozotocin–induced diabetic rats. *J Pharmacol Toxicol* 2008; **3**: 1–10.
- [5] Alarcon–Aguilara FJ, Roman–Ramos R, Perez–Gutierrez S, Aguilar–Contreras A, Contreras–Weber CC, Flores–Saenz JL. Study of the anti–hyperglycemic effect of plants used as antidiabetics. *J Ethnopharmacol* 1998; **61**: 101–110.
- [6] Jafri MA, Aslam M, Javed K, Singh S. Effect of *Punica granatum* Linn (flowers) on blood glucose level in normal and alloxan–induced diabetic rats. *J Ethnopharmacol* 2000; **70**: 309–314.
- [7] Thirumalai T, Therasa VS, Elumalai EK, David E. Hypoglycemic effect of *Brassica juncea* (seeds) on streptozotocin induced diabetic male albino rat. *Asian Pac J Trop Biomed* 2011; **4**: 323–325.
- [8] Ravi K, Ramachandran B, Subramanian S. Effect of *Eugenia Jambolana* seed kernel on antioxidant defense system in streptozotocin induced diabetes in rats. *Life Sci* 2004; **75**(22): 2717–2731.
- [9] Ajay SS. Hypoglycemic activity of *Coccinia indica* (Cucurbitaceae) leaves. *Int J Pharm Tech Res* 2009; **1**(3): 892–893.
- [10] Rajesh Kumar, Dinesh Kumar Pate, Satyendra Kuldip Prasad, Kirshnamurthy Sairam, Siva Hemalatha. Antidiabetic activity of alcoholic leaves extract of *Alangium lamarckii* Thwaites on streptozotocin–nicotinamide induced type 2 diabetic rats. *Asian Pac J Tropical Med* 2011; 904–909.
- [11] Dinesh Kumar, Sunil Kumar, Sonia Kohli, Renu Arya, Jyoti Gupta. Antidiabetic activity of methanolic bark extract of *Albizia odoratissima* Benth. in alloxan induced diabetic albino mice. *Asian Pac J Trop Med* 2011; 900–903.
- [12] Xin–Zhong Hu, Xiao–Hui Xing, Zheng–Mao Zhang, Rui–Qin Wu, Qingbin Guo, Steve W. Cui, et al. Antioxidant effects of *Artemis sphaerocephala* Krasch. gum, on streptozotocin–induced type 2 diabetic rats. *Food Hydrocolloids* 2011; **25**: 207–213.
- [13] Ibeh BO, Ezeaja MI. Preliminary study of antidiabetic activity of the methanolic leaf extract of *Axonopus compressus* (P.Beauv) in alloxan induced diabetic rats. *J Ethnopharmacol* 2011; **138**: 713–716.
- [14] Meliani N, Amine Dib ME, Allali H, Tabti B. Hypoglycaemic effect of *Berberis vulgaris* L. in normal and streptozotocin induced diabetic rats. *Asian Pac J Trop Biomed* 2011; **6**: 468–471.
- [15] Kumar R, Patel DK, Prasad SK, Laloo D, Krishnamurthy S, Hemalatha S. Type 2 antidiabetic activity of bergenin from the roots of *Caesalpinia digyna* Rottler. *Fitoterapia* 2012; **83**(2): 395–401.
- [16] Ohadoma SC, Michael HU. Effects of co–administration of methanol leaf extract of *Catharanthus roseus* on the hypoglycemic activity of metformin and glibenclamide in rats. *Asian Pac J Trop Med* 2011; 475–477.
- [17] Sefi M, Fetoui H, Lachkar N, Tahraoui A, Lyoussi B, Boudawara T, et al. *Centaurium erythraea* (Gentianaceae) leaf extract alleviates streptozotocin–induced oxidative stress and β – cell damage in rat pancreas. *J Ethnopharmacol* 2011; **135**: 243–250.
- [18] Sancheti S, Sancheti S, Seo SY. Antidiabetic and antiacetylcholinesterase effects of ethyl acetate fraction of *Chaenomeles sinensis* (Thouin) Koehne fruits in streptozotocin–induced diabetic rats. *Exp Toxicol Pathol* 2011; **65**(1–2): 55–60.
- [19] Kumar D, Kumar S, kohli S, Arya R, Gupta J. Antidiabetic activity of methanolic bark extract of *Albizia odoratissima* Benth in alloxan induced diabetic albino mice. *Asian Pac J*

- Trop Med* 2011; **4**: 900–903.
- [20]Naskar S, Mazumder UK, Pramanik G, Gupta M, Sureshkumar RB, Bala A, et al. Evaluation of antihyperglycemic activity of *Cocos nucifera* Linn.on streptozotocin induced type 2 diabetic rats. *J Ethnopharmacol* 2011; **138**: 769–773.
- [21]Eliza J, Diasy P, Ignacimuthu S, Duraipandiyar V. Antidiabetic and antilipidemic effect of eremanthin from *Costus speciosus* (Koen.)Sm., in STZ–induced diabetic rats. *Chem Biol Interact* 2009; **182**: 67–72.
- [22]Li S, Li J, Guan XL, Li J, Deng SP, Li LQ, et al. Hypoglycemic effects and constituents of the barks of *Cyclocarya paliurus* and their inhibiting activities to glucosidase and glycogen phosphorylase. *Fitoterapia* 2011; **82**: 1081–1085.
- [23]Kumar S, Kumar V, Om Prakash. Antidiabetic, hypolipidemic and histopathological analysis of *Dillenia indica* (L.) leaves extract on alloxan induced diabetic rats. *Asian Pac J Trop Med* 2011; 347–352.
- [24]Mahendran S, Badami S, Maithili V. Evaluation of antidiabetic effect of embelin from *Embelia ribes* in alloxan induced diabetes in rats. *Biomed Preventive Nutr* 2011; **1**: 25–31.
- [25]Patel DK, Kumar R, Prasad SK, Sairam K, Hemalatha S. Antidiabetic and *in vitro* antioxidant potential of *Hybanthus enneaspermus* (Linn) F. Muell in streptozotocin–induced diabetic rats. *Asian Pac J Trop Med* 2011; **4**: 316–322.
- [26]Balamurugan R, Ignacimuthu S. Antidiabetic and hypolipidemic effect of methanol extract of *Lippia nodiflora* L. in STZ induced diabetic rats. *Asian Pac J Trop Biomed* 2011; **1**: S30–36.
- [27]Hou SZ, Chen SX, Huang S, Jiang DX, Zhou CJ, Chen CQ, et al. The hypoglycemic activity of *Lithocarpus polystachyus* Rehd. leaves in the experimental hyperglycemic rats. *J Ethnopharmacol* 2011; **138**: 142–149.
- [28]Elberry AA, Harraz FM, Ghareib SA, Gabr SA, Nagy AA, Sattar EA. Methanolic extract of *Marrubium vulgare* ameliorates hyperglycemia and dyslipidemia in streptozotocin–induced diabetic rats. *Int J Diabetes Mellitus* (2011). In press.
- [29]Patil R, Patil R, Ahirwar B, Ahirwar D. Isolation and characterization of anti–diabetic component (bioactivity guided fractionation) from *Ocimum sanctum* L. (Lamiaceae) aerial part. *Asian Pac J Trop Med* 2011; 278–282.
- [30]Cetto AA, Wiedenfeld H. Anti–hyperglycemic effect of *Opuntia streptacantha* Lem. *J Ethnopharmacol* 2011; **133**: 940–943.
- [31]Huang CS, Yin MC, Chiu LC. Antihyperglycemic and antioxidative potential of *Psidium guajava* fruit in streptozotocin–induced diabetic rats. *Food Chem Toxicol* 2011; **41**: 2189–2195.
- [32]Hedayathullah Khan HB, Vinayagam KS, Palanivelu S, Panchanatham S. Anti–diabetic effect of *Semecarpus anacardium* Linn nut milk extract in a high fat diet STZ–induced type 2 diabetic rat model. *Comp Clin Pathol* 2012; **21**(6): 1395–1400.
- [33]Georgea C, Lochnera A, Huisamen B. The efficacy of *Prosopis glandulosa* as antidiabetic treatment in rat models of diabetes and insulin resistance. *J Ethnopharmacol* 2011; **137**: 298–304.
- [34]Chen X, Jin J, Tang J, Wang Z, Wanga J, Jin L, et al. Extraction, purification, characterization and hypoglycemic activity of a polysaccharide isolated from the root of *Ophiopogon japonicas*. *Carbohydrate Polymers* 2011; **83**: 749–754.
- [35]Sireesh Y, Kasetti RB, Nabi SA, Swapna S, Apparao C. Antihyperglycemic and hypolipidemic activities of *Setaria italica* seeds in STZ diabetic rats. *Pathophysiology* 2011; **18**: 159–164.
- [36]Gandhi GR, Ignacimuthu S, Paulraj MG, Sasikumar P. Antihyperglycemic activity and antidiabetic effect of methyl caffeate isolated from *Solanum torvum* Swartz. fruit in streptozotocin induced diabetic rats. *Eur J Pharmacol* 2011; **670**: 623–631.
- [37]Gupta S, Sharma SB, Singh UR, Bansal SK. Salutary effect of *Cassia auriculata* L. leaves on hyperglycemia–induced atherosclerotic environment in streptozotocin rats. *Cardiovasc Toxicol* 2011; **11**: 308–315.
- [38]Ghoul JE, Boughanmi NG, Attia MB. Biochemical study on the protective effect of ethanolic extract of *Zygophyllum album* on streptozotocin induced oxidative stress and toxicity in mice. *Biomed Preventive Nutr* 2011; **1**(2): 79–83.
- [39]Sundaram R, Naresh R, Shanthi P, Sachdanandam P. Antihyperglycemic effect of iridoid glucoside, isolated from the leaves of *Vitex negundo* in streptozotocin–induced diabetic rats with special reference to glycoprotein components. *Phytomedicine* 2012; **19**(3–4): 211–216.
- [40]Sattar EA, Elberry AA, Harraz FM, Ghareib SA, Nagy AA, Gabr SA. Antihyperglycemic and hypolipidaemic effects of the methanolic extract of Saudi mistletoe (*Viscum schimperi* Engl.). *J Adv Res* 2011; **2**: 171–177.
- [41]Sunil C, Ignacimuthu S, Agastian P. Antidiabetic effect of *Symplocos cochinchinensis* (Lour.) S. Moore. in type 2 diabetic rats. *J Ethnopharmacol* 2011; **134**: 298–304.
- [42]Sonawane RD, Vishwakarma SL, Lakshmi S, Rajani M, Padh H, Goyal RK. Amelioration of STZ–induced type 1 diabetic nephropathy by aqueous extract of *Enicostemma littorale* Blume and swertiamarin in rats. *Mol Cell Biochem* 2010; **340**: 1–6.
- [43]Feshani AM, Kouhsari SM, Mohammadi S. Vaccinium arctostaphylos, a common herbal medicine in Iran: Molecular and biochemical study of its antidiabetic effects on alloxan–diabetic Wistar rats. *J Ethnopharmacol* 2011; **133**: 67–74.
- [44]Poongothai K, Ponmurugan P, Syed Zameer Ahmed K, Senthil Kumar B, Sheriff SA. Antihyperglycemic and antioxidant effects of *Solanum xanthocarpum* leaves (field grown & *in vitro* raised) extracts on alloxan induced diabetic rats. *Asian Pac J Trop Med* 2011; 778–785.