



Contents lists available at ScienceDirect

Journal of Acute Disease

journal homepage: www.jadweb.org



Document heading doi: 10.1016/S2221-6189(13)60145-6

# Development of bioanalytical parameters for standardization of *Terminalia arjuna*

Dinesh K. Patel<sup>1,2</sup>, Kanika Patel<sup>3</sup>, Dhanabal S.P.<sup>2\*</sup><sup>1</sup>Department of Pharmaceutics, Indian Institute of Technology, Banaras Hindu University, Varanasi-221005, India<sup>2</sup>J.S.S. College of Pharmacy, Ooty- 643 001, India<sup>3</sup>G.L.A Institute of Pharmaceutical Research, Mathura-281406, India

## ARTICLE INFO

## Article history:

Received 23 May 2013

Received in revised form 3 June 2013

Accepted 18 June 2013

Available online 20 December 2013

## Keywords:

HPTLC

Phytochemical analysis

Quercetin

Rutin

Standardization

*Terminalia arjuna*

## ABSTRACT

**Objective:** To develop a novel bioanalytical parameters for standardization of *Terminalia arjuna* (*T. arjuna*) extract. **Methods:** In the present investigation, parameters such as preliminary phytochemical analysis, total phenol, flavonoid, tannin content, solubility test, heavy metal analysis, antimicrobial study and quantitative analysis by HPTLC method were performed in order to standardization. **Results:** Preliminary phytochemical analysis showed the presence of alkaloid, carbohydrate, tannin, steroid, triterpenoid, glycoside, saponin, flavonoid, amino acid and protein. Loss on drying and solubility in water was found to be 5.04% and 81.10%. Total flavonoid and phenol content was found to be 10.4% and 0.44%. Total tannin content was found to be 24.8%. The content of quercetin and rutin in *T. arjuna* was found to be 1.08%w/w and 0.16%w/w respectively. Further the level of heavy metal and microorganism were found to be under the limit. **Conclusions:** These bioanalytical parameters can be used as an important tool for the food scientists, researchers and even the consumers for its standards.

## 1. Introduction

Phytochemicals are compounds mainly responsible for different color, flavor and smell of plants and they play an important role in the plant's natural defense mechanism against various diseases[1]. Natural products offer unbounded opportunities for new drug development due to the chemical diversity. According to WHO, 80% of the World's population relies on traditional medicine to meet their daily health need. A large number of drugs prescribed worldwide are derived directly or indirectly from natural sources such as aspirin from *Filipendula ulmar* (*F. ulmar*), morphine from *Papaver somniferum* (*P. somniferum*) and ephedrin from ephedra. A number of plant based drug is included in the WHO's essential medicine list[2,3].

*Terminalia arjuna* (*T. arjuna*) belonging to family Combretaceae, was used in the Ayurvedic medicine since

ancient times for the treatment of diseases. Plant parts, such as fruit and bark are mainly used to maintain good health[4]. *T. arjuna* bark powder and extract available as over-the-counter supplements for maintaining a healthy heart in the US. *T. arjuna* barks decrease blood pressure and heart rate, and counteract actions of norepinephrine and isoproterenol[5]. *T. arjuna* is appropriately known as "Hridya", as it possesses heart strengthening and cardiogenic properties. *T. arjuna* has been used in different forms such as Asava (alcoholic decoction), Ghrita (clarified butter), Kshirpka (boiled with milk) and dried bark powder for different medicinal purposes. It is also used to treat obesity, hypertension, ulcer and hyperglycemia. It has wound healing, antibacterial, antimutagenic/anticarcinogenic, antioxidant, hypocholesterolemic and antifertility activity[6–8]. *T. arjuna* is a famous Indian folk medicinal plant used as a cardiogenic in heart failure, ischaemic cardiomyopathy, atherosclerosis and myocardium necrosis. It is an essential ingredient of many Ayurvedic preparations which are sold as cardiogenics. It is used in the treatment of fractures, ulcers, blood diseases,

\*Corresponding author: Dr. S. P. Dhanabal, Professor & Head, Department of Phytochemistry & Phytomedicine, JSS college of Pharmacy, Ooty- 643 001, India.

Tel: 09443095697

E-mail: dhanskanaksahil@rediffmail.com

anaemia and asthma and has the ability to cure hepatic, congenital, venereal and viral diseases. Many oleanane type triterpenoids from *T. arjuna* plant showed antitumoral, antioxidant, antiallergic, antiasthmatic, antifeedant and cardioprotective activities<sup>[9,10]</sup>.

Phytochemically *T. arjuna* contain tannin, saponin, ester, sugar, steroids, acids, triterpenoid saponins (e.g., arjunic acid and derivatives), ellagic acid, gallic acid, oligomeric proanthocyanidins, phytosterols, flavonoids (arjunone, arjunolone, luteolin), polyphenols, calcium, magnesium, zinc and copper<sup>[4,5,7]</sup>. Ursane triterpene glucosyl ester, 2  $\alpha$ , 3  $\beta$  -dihydroxyurs-12, 18-dien-28-oic acid 28-O- $\beta$  -D-glucopyranosyl ester, 2  $\alpha$ , 3  $\beta$ , 23-trihydroxyurs-12,18-dien-28-oic acid 28-O- $\beta$  -D-glucopyranosyl ester, quadranoside VIII, kajiichigoside F1 and 2  $\alpha$ , 3  $\beta$ , 23-trihydroxyurs-12, 19-dien-28-oic acid 28-O- $\beta$  -D-glucopyranosyl ester, as well as phenolic compounds, 3-O-methylellagic acid 4  $\beta$ -O- $\alpha$ -L-rhamnopyranoside and (-)- epicatechin were isolated from *T. arjuna* plant<sup>[10,11]</sup>.

## 2. Material and methods

### 2.1. Plant extract and Chemicals

Crude plant extract of *T. arjuna* was procured from Garlico Herbal Concentrate (M.P.), India. High Performance Thin Layer Chromatography (HPTLC) precoated plates Silica Gel Merck 60F<sub>254</sub> was used as a stationary phase. Rutin and quercetin were used as a marker compound. All the chemicals and reagents used in the present analysis of analytical grade.

### 2.2. Development of bioanalytical parameters

A phytochemical screening was conducted on the *T. arjuna* extract using standard qualitative methods to confirm the presence of different phytoconstituents<sup>[12,13]</sup>. The presence of phytoconstituents in the extract was also analysed through TLC analysis<sup>[14]</sup>. Study of parameters such as solubility in water, tannin content, loss on drying, heavy metal analysis and microbiological assay were also performed as per method of IP, 1996 and WHO guidelines<sup>[15,16]</sup>. Total phenol and flavonoid content were also determined according to the standard methods<sup>[17,18]</sup>. Different combination of solvent system has been used for the optimization of solvent system for quantitative analysis through HPTLC methods.

#### 2.2.1. Development of HPTLC methods for standardization

The quantification of rutin and quercetin in *T. arjuna* were determined through HPTLC. Different concentration of standard solution and extracts were applied on HPTLC

plates. The HPTLC plates were developed in the optimized solvent system, dried in air and scanned at 254 nm using CAMAG TLC scanner 3. Further rutin and quercetin content was determined through advance high performance thin layer chromatography (HPTLC) method. For the preparation of the calibration curve in the quantitative analysis different concentration of the standard stock solution were prepared in the HPLC grade methanol. For the preparation of the samples solution, extract was dissolve in the methanol and then sonicated for 10 min and the final volume of the solutions was made up to 5 mL to get stock solutions. All the needed concentration of the samples was prepared from the stock solution by suitable dilution.

The chromatographic conditions for the HPTLC analysis used in the present investigation are as follows:

Analysis:	Estimation of rutin and quercetin in Terminalia arjuna extract.
Plate material:	HPTLC Precoated plates Silica Gel Merck 60F <sub>254</sub> .
Solvent system:	Ethyl acetate: formic acid: glacial acetic acid: water (100:11:11:26).
Syringe:	100 $\mu$ L Hamilton (Bonadzu, Switzerland).
Application mode:	CAMAG Automatic TLC Sampler III.
Development mode:	Ascending.
Scanning:	CAMAG TLC scanner 3 with Cats software.
Experimental conditions:	Temperature (25 $\pm$ 2) $^{\circ}$ C, relative humidity 40%.

## 3. Results

Preliminary Phytochemical analysis showed that alkaloid, carbohydrate, tannin, steroid, triterpenoid, glycoside, saponin, flavonoid, amino acid and protein were found to be present. TLC analysis showed three spots  $R_f$  (0.28, 0.19, 0.06) in chloroform: methanol: ethanol (90:05:05) solvent system. Loss on drying and solubility in water was found to be 5.04%, 81.10%. The total flavonoid, phenol and tannin content were found to be 10.4%, 0.44% and 24.8%. Further heavy metal analysis was also performed in the current task and the level of lead, arsenic, mercury and cadmium were found to be under the limit. Microbiological assay was also performed in the current task and result showed that *E. coli* and salmonella was found to be absent whereas total bacterial count and yeast & moulds contents were found to be 445 and 56 CFU/GM.


Fingerprinting analysis of sample was done through HPTLC method, and the data were presented in the Table 1. For quantitative analysis through HPTLC techniques, optimization of solvent system was done using combination of solvent system of varying polarity and the most suitable solvent system were taken for the quantitative analysis. Ethyl acetate: formic acid: glacial acetic acid: water (100:11:11:26)

was found to be suitable solvent system for quantitative analysis through HPTLC. The content of quercetin and rutin in *T. arjuna* was found to be 1.08%w/w and 0.16%w/w respectively. The respective HPTLC chromatogram of rutin, quercetin, and *T. arjuna* extract were presented in the Figure 1, Figure 2 and Figure 3. The interpretations of result were done using standard calibration curve of quercetin and rutin (Figure 4 and Figure 5). The results suggest that the sample contained considerable amount of rutin, and quercetin.


**Table 1**

Fingerprint analysis of *T. arjuna* extract.


Solvent system	R <sub>f</sub> value	Maximum peak height	Peak area (%)
<i>n</i> -butanol : acetic acid : H <sub>2</sub> O (4:1:5)	0.14	248.8	24.40
	0.16	247.8	14.91
	0.28	15.3	0.76
	0.38	20.2	1.24
	0.56	66.3	7.06
	0.74	271.3	43.76
	0.86	41.1	3.23
	0.95	66.6	4.65




**Figure 3.** Standard HPTLC chromatogram of *T. arjuna* extract.




**Figure 1.** Standard HPTLC chromatogram of quercetin.



**Figure 4.** Standard calibration curve of quercetin.



**Figure 2.** Standard HPTLC chromatogram of rutin.



**Figure 5.** Standard calibration curve of rutin.

#### 4. Discussion

Physicochemical and phytochemical analysis are used to check the genuine nature of the crude drug, thus it plays an important role in preventing the possible steps of adulteration<sup>[19]</sup>. Based on the reported traditional use in the literature, in the present investigation *T. arjuna* extract was taken.

The medicinal properties of plants material are mainly due to the presence of various phytoconstituents<sup>[20]</sup>. The presence of different phytoconstituents such as flavonoid, tannin, saponins, alkaloid and glycoside in the phytochemical tests justifies their therapeutic potential<sup>[21–23]</sup>.

These phytoconstituents have been reported to have multiple biological effects such as anti-inflammatory, anti allergic, antioxidant, antidiabetic, analgesic, antispasmodic, antibacterial, anti-viral, anti-cancer and aldose reductase inhibitory activities. It is also used for the treatment of diarrhea and dysentery<sup>[24–26]</sup>. Phytoconstituents obtained from natural sources have been gaining importance in the day by day due to the health promoting activity. So it is necessary to check the quality safety and efficacy of herbal drugs before its consumption<sup>[27,28]</sup>.

Phytochemical standardization plays an important role to ensure the quality safety and efficacy of the herbal drug. In the last few decades, an HPTLC technique has gained much popularity for standardization of the herbal drugs and formulations. Analysis of several samples simultaneously using a small quantity of marker compound and mobile phase with very less time is the major advantage of HPTLC<sup>[28]</sup>.

TLC and HPTLC techniques have been used as important analytical tools in pharmaceuticals, medicine, chemistry, food analysis, toxicology and environmental science<sup>[29]</sup>. Quality evaluation and standardization of the herbal preparation is a fundamental requirement of industry and other organization dealing with ayurvedic and herbal products. According to WHO guidelines, an herbal product needs to be standardized with respect to safety before releasing it into the market<sup>[30]</sup>.

These bioanalytical parameters can be utilized for the simultaneous analysis of different phytoconstituents present in the *T. arjuna* plant material.

In future, this information may be useful as a standard to identify and to differentiate from its adulterants and other related species.

#### Conflict of interest statement

The authors report no conflict of interest.

#### Acknowledgments

This work was financially supported by the TIFAC CORE in Herbal Drugs by DST in J.S.S. College of Pharmacy Ooty, T.N., India.

#### References

- [1] Eleazu CO, Eleazu KC, Awa E, Chukwuma SC. Comparative study of the phytochemical composition of the leaves of five Nigerian medicinal plants. *J Biotechnol Pharm Res* 2012; **3**: 42–46.
- [2] Dangoggo SM, Hassan LG, Sadiq IS, Manga SB. Phytochemical analysis and antibacterial screening of leaves of *Diospyros Mespiliformis* and *Ziziphus Spina-Christi*. *J Chem Eng* 2012; **1**: 31–37.
- [3] Patel DK, Laloo D, Kumar R, Hemalatha S. *Pedaliium murex* Linn.: an overview of its phytopharmacological aspects. *Asian Pac J Trop Med* 2011; **4**: 748–755.
- [4] Ghosh J, Das J, Manna P, Sil PC. Protective effect of the fruits of *Terminalia arjuna* against cadmium-induced oxidant stress and hepatic cell injury via MAPK activation and mitochondria dependent pathway. *Food Chem* 2010; **123**: 1062–1075.
- [5] Oberoi L, Akiyama T, Lee KH, Liu SJ. The aqueous extract, not organic extracts, of *Terminalia arjuna* bark exerts cardioprotective effect on adult ventricular myocytes. *Phytomedicine* 2011; **18**: 259–65.
- [6] Singh G, Singh AT, Abraham A, Bhat B, Mukherjee A, Verma R, et al. Protective effects of *Terminalia arjuna* against Doxorubicin-induced cardiotoxicity. *J Ethnopharmacol* 2008; **117**: 123–129.
- [7] Devi RS, Narayan S, Vani G, Shyamala Devi CS. Gastroprotective effect of *Terminalia arjuna* bark on diclofenac sodium induced gastric ulcer. *Chem Biol Interact* 2007; **167**: 71–83.
- [8] Singh DV, Verma RK, Singh SC, Gupta MM. RP–LC determination of olean derivatives in *Terminalia arjuna*. *J Pharm Biomed Anal* 2002; **28**: 447–452.
- [9] Kaur K, Arora S, Kumar S, Nagpal A. Antimutagenic activities of acetone and methanol fractions of *Terminalia arjuna*. *Food Chem Toxicol* 2002; **40**: 1475–1482.
- [10] Wang W, Ali Z, Shen Y, Li XC, Khan IA. Ursane triterpenoids from the bark of *Terminalia arjuna*. *Fitoterapia* 2010; **81**: 480–484.

- [11] Pawar RS, Bhutani KK. Effect of oleanane triterpenoids from *Terminalia arjuna*—a cardioprotective drug on the process of respiratory oxyburst. *Phytomedicine* 2005; **12**: 391–393.
- [12] Khandelwal KR. *Practical pharmacognosy techniques and experiments*. 17th Ed. India. 2007.
- [13] Harbourne JB. *Phytochemical methods: A guide to modern techniques of plant analysis*. London: Chapman and Hall; 1984.
- [14] Wagner H, Bladt S. *Plant drug analysis, A thin layer chromatography atlas*. Heidelberg: Springer-Verlog Berlin; 1996.
- [15] Indian Pharmacopoeia. Government of India, Ministry of Health and Family Welfare. Controller of publications publishers, Delhi. Vol-II (P-Z), p A-54, 1996.
- [16] World Health Organization. Traditional medicine.[Online]. Available from: <http://www.who.int/mediacentre/factsheets/fs134/en/>. [Retrieved April 12, 2009].
- [17] Singleton VL, Rossi JA. Colorimetry of total phenolics with phosphomolybdicphosphotungstic acid reagents. *Am J Enol Vitic* 1965; **16**: 144–158.
- [18] Chang CC, Yang MH, Wen HM, Chern JC. Estimation of total flavonoid content in propolis by two complementary colorimetric methods. *J Food Drug Anal* 2002; **10**: 178–182.
- [19] Mohan VR, Chenturpandy P, Kalidass C. Pharmacognostic and phytochemical investigation of *Elephantopus scaber* L. (Asteraceae). *J Pharm Sci Technol* 2010; **2**: 191–197.
- [20] Lalnundanga NL, Lalrinkima R. Phytochemical analysis of the methanol extract of root bark of *Hiptage benghalensis* (L.). *Kurz Sci Vis* 2012; **12**: 8–10.
- [21] Kavitha T, Nelson R, Thenmozhi R, Priya E. Antimicrobial activity and phytochemical analysis of *Anisomeles malabarica* (L) R.BR. *J Microbiol Biotech Res* 2012; **2**: 1–5.
- [22] Patel DK, Kumar R, Kumar M, Sairam K, Hemalatha S. Evaluation of aldose reductase inhibitory potential of different fraction of *Hybanthus enneaspermus* linn f. muell. *Asian Pac J Trop Biomed* 2012; **2**: 134–139.
- [23] Ayoola PB, Adeyeye A. Phytochemical and nutrient evaluation of *Carica papaya* (pawpaw) leaves. *IJRRAS* 2010; **5**: 325–328.
- [24] Patel DK, Kumar R, Laloo D, Hemalatha S. Evaluation of phytochemical and antioxidant activities of the different fractions of *Hybanthus enneaspermus* (Linn.) F. Muell. (Violaceae). *Asian Pac J Trop Med* 2011; **4**: 412–420.
- [25] Patel DK, Kumar R, Prasad SK, Sairam K, Hemalatha S. Antidiabetic and *in vitro* antioxidant potential of *Hybanthus enneaspermus* linn f. muell in streptozotocin-induced-diabetic rats. *Asian Pac J Trop Biomed* 2011; **1**: 316–322.
- [26] Patel DK, Kumar R, Prasad SK, Hemalatha S. *Petalium murex* Linn (Pedaliaceae) fruits: A comparative antioxidant activity of its different fractions. *Asian Pac J Trop Biomed* 2011; **1**: 395–400.
- [27] Patel DK, Kumar R, Laloo D, Hemalatha S. Diabetes mellitus: An overview on its pharmacological aspects and reported medicinal plants having anti-diabetic activity. *Asian Pac J Trop Biomed* 2012; **2**: 411–420.
- [28] Kshirsagar VB, Deokate UA, Bharkad VB, Khadabadi SS. HPTLC method development and validation for the simultaneous estimation of diosgenin and levodopa in marketed formulation. *Asian J Res Chem* 2008; **1**: 36–39.
- [29] Choudhary N, Sekhon BS. An overview of advances in the standardization of herbal drugs. *J Pharm Educ Res* 2011; **2**: 55–70.
- [30] Jain A, Lodhi S, Singhai AK. Simultaneous estimation of quercetin and rutin in *Tephrosia purpurea* Pers by high performance thinlayer chromatography. *Asian J Traditional Med* 2009; **4**: 104–109.