

Original article

Synergistic effects of plant extracts and antibiotics on *Staphylococcus aureus* strains isolated from clinical specimens

Ghaleb Adwan¹, Mohammad Mhanna²

¹Department of Biology and Biotechnology, An-Najah N. University, P. O. Box (7)-Nablus, Palestine

²Faculty of Graduate Studies, Department of Environmental Sciences, An-Najah N. University, P. O. Box (7)-Nablus, Palestine

Abstract

Objective: This study has been done to evaluate the interaction between water extracts of *Psidium guajava*, *Rosmarinus officinalis*, *Salvia fruticosa*, *Majorana syriaca*, *Ocimum basilicum*, *Syzygium aromaticum*, *Laurus nobilis*, and *Rosa damascena* alone and then synergy testing of these extracts with known antimicrobial agents including oxytetracycline HCl, gentamicin sulfate, penicillin G, cephalexin and enrofloxacin. This study was conducted against five *S. aureus* isolates; one is Methicillin – resistant *Staphylococcus aureus* (MRSA) and 4 Methicillin – sensitive *Staphylococcus aureus* (MSSA). **Methods:** Evaluation of the interaction between plant extracts and different antimicrobial agents has been done using well – diffusion and microdilution methods.

Results: The results of the conducted experiments using well – diffusion method demonstrate that these plants showed in vitro interactions between antimicrobial agents and plant extracts were additive, while using microdilution method showed synergistic effects with significant reduction in the MICs of the test antibiotics against these strains of *S. aureus*. This change in MIC was noticed in all plant extracts against test antibiotics including these plants showed weak antibacterial activity by well diffusion method. Synergism effect was occurred in both sensitive and resistant strains but the magnitude of minimum fold reduction of inhibitory concentration in resistant strains especially MRSA strain was higher than the sensitive strains. **Conclusion:** This study probably suggests the possibility of concurrent use of these antimicrobial drugs and plant extracts in combination in treating infections caused by *S. aureus* strains or at least the concomitant administration may not impair the antimicrobial activity of these antibiotics.

Keywords: Plant extracts; Synergistic effects; Antimicrobial agents; Microdilution method; Well diffusion method

INTRODUCTION

The wide use of antibiotics in the treatment of bacterial infections has led to the emergence and spread of resistant strains. Methicillin-resistant *Staphylococcus aureus* (MRSA) is a major cause of nosocomial in-

fections. MRSA infections are very difficult to cure because MRSA strains are resistance against almost all clinically available antibiotics. For most MRSA strains, glycopeptide-type drugs such as vancomycin are the only effective antimicrobial agents^[1]. However, vancomycin-resistant *S. aureus* (VRSA) has been reported^[2-4]. Thus, it is extremely important to find new antimicrobial agents or new ways that are effective for the treatment of infectious diseases caused by drug-resistant bacteria including MRSA^[5].

Few studies have found that the efficacy of anti-

Correspondence to: Ghaleb Adwan, Department of Biology and Biotechnology, An-Najah N. University, P. O. Box (7)-Nablus, Palestine.

Fax: +970-9-2347488

E-mail : adwang@najah.edu



icrobial agents can be improved by combining them with crude plant extracts against different pathogens including *S. aureus*, *P. aeruginosa*, *E. coli*, Extended Spectrum β -lactamases-producing multidrug-resistant *E. coli*, and vancomycin-resistant enterococci (*Enterococcus fecalis*)^[6-18]. Some Palestinian plants exhibit significant potency against human bacterial^[19-21]. But plant extracts as antimicrobials are rarely used as systemic antibiotics at present, this may be due to their low level of activity, especially against Gram-negative bacteria. Here we are trying to investigate an alternative approach to the treatment of bacterial infections by able to change the phenotype of a resistant pathogen to certain antibiotics to more susceptible pathogen to that antibiotics. This alternative approach has been evaluated using in vitro interaction between water plant extracts of the following plants: *Psidium guajava*, *Rosmarinus officinalis*, *Salvia fruticosa*, *Majorana syriaca*, *Ocimum basilicum*, *Syzygium aromaticum*, *Laurus nobilis*, *Rosa damascena* and certain known antimicrobial drugs such as oxytetracycline HCl, gentamicin sulphate, penicillin G, cephalixin, and enrofloxacin using well diffusion method and minimum inhibitory concentration (MIC) by microdilution method against 4 Methicillin-sensitive *Staphylococcus aureus* (MSSA) and one strain MRSA.

MATERIALS AND METHODS

Plant material and extract preparation

The plant materials used in this study consisted of *Psidium guajava* (leaf), *Rosmarinus officinalis* (leaf), *Salvia fruticosa* (leaf), *Majorana syriaca* (leaf), *Ocimum basilicum* (leaf), *Rosa damascena* (flower) which are growing in Palestine, and *Syzygium aromaticum* (dried flowerbud), *Laurus nobilis* (leaf) were collected from Palestinian markets. These plants were identified by Dr. Firas. D. Sawalha, Department of Plant Production, Faculty of Agriculture, An-Najah National University, Nablus, Palestine. Water extract was prepared as describe previously^[21]. Approximately of 30 g of dried plant materials were separately powdered, extracted by adding 150-200 mL boiled distilled water, and then kept for 1 h. The extracts were filtered through

Whatman No. 2 filter paper under vacuum. Extracts were concentrated to dryness at 37°C. Then, 100 mg of the dry residue was dissolved in 1 mL of sterile distilled water.

Antimicrobial agents and antibiotics

Five antimicrobial agents were evaluated for synergism assays with different plant extracts. These agents included oxytetracycline HCl (10%), enrofloxacin (10%), getamicin Sulphate (50%), cephalixin (0.15%) and penicillin G (penicillin G procaine 900 000 and penicillin G sodium 300 000 I. U). All these antimicrobial agents were produced by Jerusalem Pharmaceutical CO. Balsam branch except penicillin G was produced by Birzeit-Palestine Pharamaceutical CO, and were diluted to a final concentration 100 μ g/mL except penicillin G to 100 I. U/mL for well diffusion method and 200 μ g/mL, and 200 I. U/mL for MIC test.

Bacterial strains

Five *S. aureus* isolates (one MRSA and 4 MSSA) used in this study were recovered from urine sample, Thabet Thabet's Hospital, Toulkarm; semen sample, Al-Zaka Hospital, Toulkarm; diabetic foot wound, Al-Makased Hospital, Jerusalem; chest wound, Al-Makased Hospital, Jerusalem and leg wound, Al-Makased Hospital, Jerusalem. All isolates were identified in the microbiology laboratories of An-Najah National University, Palestine, by Gram stain, culture properties on nutrient agar and mannitol salt agar, catalase test, detection of hemolysis on 5% blood agar and by tube coagulase reaction. *S. aureus* strains were tested for methicillin resistance using the disk diffusion method^[22]. Oxacillin (1 μ g), and methicillin (5 μ g) disks (Oxoid) were used. Zones of inhibition were determined in accordance with procedures of the National Committee for Clinical Laboratory Standard^[23], *S. aureus* isolates considered susceptible to methicillin and oxacillin if inhibition zones are ≥ 14 and ≥ 13 mm, respectively. A reference strain (*Bacillus subtilis* ATCC6633) was also tested.

Antimicrobial activity tests

a. Determination of the combined activity using

Well-diffusion method; Antibacterial activity was measured using a well diffusion method according to the Clinical and Laboratory Standards Institute (formerly, the NCCLS) [24]. Briefly, Petri plates containing approximately 25-30 mL of Mueller Hinton agar medium were inoculated using a cotton swab with a 4-6 h old culture of the bacterial strains. Wells (6mm diameter) were punched in the agar and filled with 30 μ L of plant extracts or antibiotics and in case of synergism effect 30 μ L of each has been added into well. Replicate of each plate has been done. The plates were incubated at 37 °C for 18-24 h. The antibacterial activity was assessed by measuring the inhibition zone diameter (mm) around the well. The average of three replicates for each extract, antibiotic, and combination has been calculated. Synergism effect was considered when combinations exhibited with enlargement of combined inhibition zone size by 5 mm^[9].

b. Determination of MIC by microdilution method; MIC of antibiotic was determined by the microdilution method as described by Clinical and Laboratory Standards Institute [25]. The antibiotic was serially diluted in Mueller Hinton broth. Plant extracts solution were separately added into wells in a final concentration 1.5 mg/mL, then bacterial inoculum size of 10⁴ CFU/mL was added to each well. Controls without plant extracts, without bacterial inoculum or with plant extracts only were also included in the experiment. Each plant extract was run in duplicate. The test plates were incubated at 37°C for 18 h. The MIC was taken as the minimum concentration of the dilutions that inhibited the growth of the test microorganism.

RESULTS

The results of the conducted experiments using well-diffusion method demonstrate that these plants contain bioactive compounds some of them has a weak effect such as *Ocimum basilicum* and *L. nobilis*. *In vitro* interactions between antimicrobial agents and plant extracts using the previous method were additive against the five strains of *S. aureus* (data not shown), while using microdilution method showed synergistic effects between combination of antibiotics and plant extracts with significant reduction in the

MICs of the test antibiotics against 5 strains of *S. aureus*. This change in MIC was noticed in all plant extracts against test antibiotics including these plants showed weak antibacterial activity by well diffusion method. Also our results showed that synergism effect between antimicrobial agent and plant extract was occurred in both sensitive and resistant strains but the magnitude of minimum fold inhibition in resistant strains especially MRSA strain was higher than the sensitive strains. Minimum fold inhibition with drug-plant extract combinations against these strains is presented in Table 1.

DISCUSSION

Combined antibiotic therapy has been shown to delay the emergency of bacteria resistance and may also produce desirable synergistic effects in the treatment of bacteria infection. Drug synergism between known antibiotics and bioactive plant extracts is a novel concept, and could be beneficial (synergistic or additive interaction) or deleterious (antagonistic or toxic outcome). Despite the abundant literature about the antimicrobial properties of plant extracts, none of the plant derived chemicals have successfully been used for clinical use as antibiotics [26].

In our experiments, despite that some plant extracts showed weak antimicrobial effect using well diffusion method, the interactions between antibiotics and plant extracts were mainly additive against the five strains of MSSA and MRSA. This could be attributed to the inability of higher concentrations of plant extracts to diffuse through the nutrient agar medium. This impairment in drug diffusion is a major limitation in the evaluation of the antimicrobial effects of plant extracts using the agar diffusion method [11].

In this study, synergism effect resulting from the combination of antimicrobial agents with crude plant extracts was verified for all plants. Our results were consistent with previous *in vitro* studies which reported synergistic effects with significant reduction in the MICs of the antibiotics due to combination of different antimicrobial agents with different crude plant extracts against *S. aureus* strains [10-11, 13-15, 18] and stand out as veritable sources of potential resistance modifying agents [27-29].

Table 1 Minimum inhibitory concentration of antibiotics alone and in combination with aqueous plant extracts against 5 clinical isolates (4 MSSA and 1 MRSA) of *S. aureus* using microdilution method.

Antibiotic ^a /Plant extract	MIC (mg/L)		MFRIC ^b for MSSA strains	MFRIC ^b for MRSA strain
	Four strains of MSSA	One strain of MRSA		
CN	1.563-3.125	25-50		
<i>P. guajava</i> + CN	0.024 4-0.048 8	0.195	32	128
<i>R. officinalis</i> + CN	0.024 4	0.39	64	64
<i>S. fruticosa</i> + CN	0.012 2-0.024 4	0.195	64	128
<i>M. syriaca</i> + CN	0.024 4-0.048 8	0.78	32	32
<i>O. basilicum</i> + CN	0.012 2-0.024 4	0.39	32	64
<i>S. aromaticum</i> + CN	0.012 2-0.024 4	<0.048 8	64	>512
<i>L. nobilis</i> + CN	0.048 8-0.097 7	0.39	16	64
<i>R. damascena</i> + CN	0.012 2-0.024 4	<0.048 8	64	>512
ENR	0.195-0.39	0.39		
<i>P. guajava</i> + ENR	<6.1 × 10 ⁻³	<6.1 × 10 ⁻³	>32	>64
<i>R. officinalis</i> + ENR	<6.1 × 10 ⁻³	<6.1 × 10 ⁻³	>32	>64
<i>S. fruticosa</i> + ENR	<6.1 × 10 ⁻³	<6.1 × 10 ⁻³	>32	>64
<i>M. syriaca</i> + ENR	<6.1 × 10 ⁻³	<6.1 × 10 ⁻³	>32	>64
<i>O. basilicum</i> + ENR	<6.1 × 10 ⁻³	<6.1 × 10 ⁻³	>32	>64
<i>S. aromaticum</i> + ENR	<6.1 × 10 ⁻³	<6.1 × 10 ⁻³	>32	>64
<i>L. nobilis</i> + ENR	<6.1 × 10 ⁻³	<6.1 × 10 ⁻³	>32	>64
<i>R. damascena</i> + ENR	<6.1 × 10 ⁻³	<6.1 × 10 ⁻³	>32	>64
OT	0.78-6.25	25-50		
<i>P. guajava</i> + OT	<0.024 4	0.097 7-0.195	≥32	128
<i>R. officinalis</i> + OT	<0.024 4-0.048 8	0.39	≥16	64
<i>S. fruticosa</i> + OT	<0.024 4	0.195	>32	128
<i>M. syriaca</i> + OT	0.048 8	0.195-0.39	16	64
<i>O. basilicum</i> + OT	0.048 8	0.39	16	64
<i>S. aromaticum</i> + OT	<0.024 4	<0.048 8	>32	>512
<i>L. nobilis</i> + OT	0.048 8	0.78	16	32
<i>R. damascena</i> + OT	<0.024 4	<0.048 8	>32	>512
CL	0.195-0.39	50		
<i>P. guajava</i> + CL	<6.1 × 10 ⁻³	0.048 8	>32	1024
<i>R. officinalis</i> + CL	<6.1 × 10 ⁻³	0.097 7	>32	512
<i>S. fruticosa</i> + CL	<6.1 × 10 ⁻³	0.097 7	>32	512
<i>M. syriaca</i> + CL	<6.1 × 10 ⁻³	0.39	>32	128
<i>O. basilicum</i> + CL	<6.1 × 10 ⁻³	0.39	>32	128
<i>S. aromaticum</i> + CL	<6.1 × 10 ⁻³	<0.048 8	>32	>1 024
<i>L. nobilis</i> + CL	<6.1 × 10 ⁻³	0.39-0.78	>32	64
<i>R. damascena</i> + CL	<6.1 × 10 ⁻³	<0.048 8	>32	>1 024
P	>100	>100		
<i>P. guajava</i> + P	1.563-3.125	0.097 7	>32	1 024
<i>R. officinalis</i> + P	0.78-1.563	0.195	>64	512
<i>S. fruticosa</i> + P	0.39-1.563	0.048 8	>64	2 048
<i>M. syriaca</i> + P	0.78-1.563	0.39	>64	256
<i>O. basilicum</i> + P	1.563-3.125	0.39	>32	256
<i>S. aromaticum</i> + P	0.78	<0.048 8	>128	>2 048
<i>L. nobilis</i> + P	1.563-3.125	0.39	>32	256
<i>R. damascena</i> + P	0.78	<0.048 8	>128	>2 048

^a P, Penicillin G; CN, Gentamicin sulphate; CL, Cephalexin; ENR, Enrofloxacin; OT, Oxytetracycline HCl.

^b MFRIC; Minimum fold reduction of inhibitory concentration.

In these experiments, the change in MIC was noticed in all plant extracts against test antibiotics including these plants showed weak antibacterial activity. These results were in agreement with a previous report who mentioned a synergetic effect even these extracts did not show any activity by themselves^[17]. In addition, results showed that synergism effect was occurred in both sensitive and resistant strains but the magnitude of minimum fold inhibition in sensitive is less than resistant strains. These results were consistent with that which showed that synergistic interactions occurred in both resistant and sensitive *S. aureus*^[15,16].

All plant extracts showed a decrease in MIC to test antimicrobial agents, and this could be referred to that these crude extracts have many different phytochemicals^[30], which might inhibit bacteria by different mechanisms. This double attack of both agents on different target sites of the bacteria could theoretically lead to either an additive or a synergistic effect^[11]. Screening for such activities in crude extracts is the first step in identifying leads for isolation of such compounds, and some plants have provided good indications of these potentials for use in combination with antimicrobial therapy. Further separation and purification of the crude extracts might show an increase in bioactivity than the crude extracts. This may be due to numerous compounds within the crude extracts may have interfered with the actions of one another. Once they were separated by various purification methods however, the inhibiting effect of one on the other had reduced significantly^[6].

Here we recommended the evaluation of the exact drug-plant ratio at which the interaction in maximal between the plant extract and antimicrobial drug. A wider study with increase in the number of drugs, increase number of clinical isolates especially MRSA, and the identification of the effective compounds in the crude extract are also necessary in order to establish the mode of action against the *S. aureus* isolates and the mechanism of synergy, which is fundamental to development of pharmacological agents to treat diseases by *S. aureus* using medicinal plants.

In conclusion, the results of this study were encouraging, although clinical controlled studies are needed to define the real efficacy and possible toxic effects *in vivo*. This study probably suggests the possibility of concurrent use of these antimicrobial drugs

and extracts in combination in treating infections caused by *S. aureus* strains or at least the concomitant administration of these plants and antimicrobial drugs may not impair the antimicrobial activity of these antibiotics. However, it is hard to predict synergistic effects *in vivo* on the basis of the presented *in vitro* evidence alone because it is difficult to estimate the *in vivo* concentration of active ingredients, especially the bioavailable concentration of free (active) ingredients, after plants have been ingested. Therefore, our results revealed the importance of plant extracts when associated with antibiotics to control bacteria.

REFERENCES

- 1 **Patterson JE**. New gram-positive agents in nosocomial infection. *Curr Opin Dis.* 2000; 13: 593-598.
- 2 Centers for Disease Control and Prevention. National Nosocomial Infections Surveillance (NNIS) report, data summary from January 1992-June 2001. *Am J Infect Control.* 2001; 29: 404-421.
- 3 Centers for Disease Control and Prevention. *Staphylococcus aureus* resistant to vancomycin-United States, 2002. *Morb Mortal Wkly Rep.* 2002; 51: 565-567.
- 4 **Chang S**, Sievert DM, Hageman JC, Boulton ML, Tenover FCT, Downes FP, et al. Infection with vancomycin-resistant *Staphylococcus aureus* containing the *vanA* resistance gene. *N Engl J Med.* 2003; 348: 1342-1347.
- 5 **Taylor PW**, Stapleton PD, Luzio JP. New ways to treat bacterial infections. *Drug Discov Today.* 2002; 7: 1086-1091.
- 6 **Adwan MG**, Abu-Shanab AB, Adwan MK. In vitro activity of certain drugs in combination with plant extracts against *Staphylococcus aureus* infections. *Pak J Med.* 2008; 24: 541-544.
- 7 **Chang PC**, Li, HY, Tang HJ, Liu JW, Wang JJ, Chuang YC. In vitro synergy of baicalein and gentamicin against vancomycin-resistant *Enterococcus*. *J Microbiol Immunol Infect.* 2007; 40: 56-61.
- 8 **Horiuchi K**, Shiota S, Kuroda T, Hatano T, Yoshida T, Tsuchiya T. Potentiation of antimicrobial activity of aminoglycosides by carnosol from *Salvia officinalis*. *Biol Pharm Bull.* 2007; 30: 287-290.
- 9 **Ahmad I**, Aqil F. *In vitro* efficacy of bioactive extracts of 15 Medicinal plants against ES β L-producing multidrug-resistant enteric bacteria. *Microbiol Res.* 2007; 162: 264-275.
- 10 **Betoni JE**, Mantovani RP, Barbosa LN, Di Stasi LC, Junior AF. Synergism between plant extract and antimicrobial drugs used on *Staphylococcus aureus* diseases. *Mem Inst Oswaldo Cru Rio de Janeiro.* 2006; 101: 387-390.



- 11 **Esimone CO**, Iroha IR, Ibezim EC, Okeh CO, Okpana EM. *In vitro* evaluation of the interaction between tea extracts and penicillin G against *Staphylococcus aureus*. *Afr J Biotechnol.* 2006; 5: 1082-1086.
- 12 **Ibezim EC**, Esimone CO, Nnamani PO, Onyishi IV, Brown SA, Obodo CE. *In vitro* study of the interaction between some fluoroquinolones and extracts of *kola nitida* seed. *Afr J Biotechnol.* 2006; 5: 1781-1784.
- 13 **Yang ZC**, Wang BC, Yang XS, Wang Q, Ran L. The synergistic activity of antibiotics combined with eight traditional Chinese medicines against two different strains of *Staphylococcus aureus*. *Colloids Surf B Biointerfaces.* 2005; 41: 79-81.
- 14 **Aqil F**, Khan MSA, Owais M, Ahmad I. Effect of certain bioactive plant extracts on clinical isolates of β -lactamase producing methicillin resistant *Staphylococcus aureus*. *J Basic Microbiol.* 2005; 45: 106-114.
- 15 **Braga LC**, Leite AAM, Xavier KGS, Takahashi JA, Bemquerer MP, Chartone-Souza E, et al. Synergic interaction between pomegranate extract and antibiotics against *Staphylococcus aureus*. *Can J Microbiol.* 2005; 51: 541-547.
- 16 **Darwish RM**, Aburjai T, Al-Khalil S, Mahafzah A. Screening of antibiotic resistant inhibitors from local plant materials against two different strains of *Staphylococcus aureus*. *J Ethnopharmacol.* 2002; 79: 359-364.
- 17 **Nascimento GGF**, Locatelli J, Freitas PC, Silva GL. Antibacterial activity of plant extracts and phytochemicals on antibiotic-resistant bacteria. *Braz J Microbiol.* 2000; 31: 247-256.
- 18 **Yam TS**, Hamilton-Miller JM, Shah S. The effect of a component of tea (*Camellia sinensis*) on methicillin resistance, PBP2' synthesis, and beta-lactamase production in *Staphylococcus aureus*. *J Antimicrob Chemother.* 1998; 42: 211-216.
- 19 **Abu-Shanab B**, Adwan G, Abu-Safiya D, Jarrar N, Adwan K. Antibacterial activities of some plant extracts utilized in popular medicine in Palestine. *Turk J Biol.* 2004; 28: 99-102.
- 20 **Abu-Shanab B**, Adwan G, Jarrar N, Abu-Hijli A, Adwan K. Antibacterial activity of four plant extracts used in Palestine in folkloric medicine against methicillin-resistant *Staphylococcus aureus*. *Turk J Biol.* 2006; 30: 195-198.
- 21 **Adwan G**, Abu-Shanab B, Adwan K, Abu-Shanab F. Antibacterial effects of nutraceutical plants growing in Palestine on *Pseudomonas aeruginosa*. *Turk J Biol.* 2006; 30: 239-242.
- 22 **Bauer AW**, Kirby WM, Sherris JC, Truck M. Antibiotic susceptibility testing by a standardized single disk method. *Am J Clin Pathol.* 1966; 45: 493-496.
- 23 National Committee for Clinical Laboratory Standards (NCCLS). Performance standards for antimicrobial disk and dilution susceptibility tests for bacteria isolated from animals. NCCLS, Pennsylvania, USA, 1991. M31-A.
- 24 National Committee for Clinical Laboratory Standards (NCCLS). Performance Standards for Antimicrobial Disc Susceptibility Tests. Villanova, PA, USA; Approved Standard NCCLS Publication, 1993. M2-A5.
- 25 National Committee for Clinical Laboratory Standards (NCCLS). Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically. NCCLS, Pennsylvania, USA, 2000. M7-A5.
- 26 **Gibbons S**. Anti-staphylococcal plant natural products. *Nat Prod Rep.* 2004;21: 263-277.
- 27 **Sibanda T**, Okok AI. The challenges of overcoming antibiotic resistance; Plant extracts as potential sources of antimicrobial and resistance modifying agents. *Afr J Biotechnol.* 2007; 6: 2886-2896.
- 28 **Dickson RA**, Houghton PJ, Hylands PJ, Gibbons S. Antimicrobial, resistance-modifying effects, antioxidant and free radical scavenging activities of *Mezoneuron benthamianum* Baill, *Securinega virosa* Roxb. and Willd. and *Microglossa pyrifolia* Lam. *Phytother Res.* 2006; 20: 41-45.
- 29 **Gibbons S M**, Oluwatuyi M, Veitch NC, Gray AI. Bacterial resistance modifying agents from *Lycopus europaeus*. *Phytochemistry.* 2003; 62: 83-87.
- 30 **Duke JA**, Jobogenschutz-Godwin M, DuCellier J, Duke PAK. CRC Handbook of Medical Plant. Boca Raton: CRC Press, FL. 2003. 348.