

RESEARCH ARTICLE

Which Biology Marker Can be a Predictor for Higher Syntax Score?Linda Lison^{1,2}, Irawan Yusuf^{1,3}, Bambang Sutrisna^{4,5}, Peter Kabo^{1,6}, Teguh Santoso^{2,7},
Ferry Sandra^{8,9,10,11}*¹Postgraduate Program in Medicine, Faculty of Medicine, Hasanuddin University, Jl. Perintis Kemerdekaan Km.10, Makassar, Indonesia²Medistra Hospital, Jl. Jendral Gatot Subroto Kav. 59, Jakarta, Indonesia³Department of Physiology, Faculty of Medicine, Hasanuddin University, Jl. Perintis Kemerdekaan Km.10, Makassar, Indonesia⁴Faculty of Medicine, University of Tarumanagara, Jl. Letjen S. Parman No. 1, Grogol, Jakarta, Indonesia⁵Faculty of Public Health, University of Indonesia, Kampus UI, Depok, Indonesia⁶Department of Cardiology and Vascular Medicine, Faculty of Medicine, Hasanuddin University, Jl. Perintis Kemerdekaan Km.10, Makassar, Indonesia⁷Division of Cardiology, Department of Internal Medicine, Faculty of Medicine, University of Indonesia, Jl. Salemba Raya No.6, Jakarta, Indonesia⁸Department of Biochemistry and Molecular Biology, Faculty of Dentistry, Trisakti University, Jl. Kyai Tapa No.260, Jakarta, Indonesia⁹Postgraduate Program in Biomedics, Faculty of Dentistry, Trisakti University, Jl. Kyai Tapa No.260, Jakarta, Indonesia¹⁰BioCORE Laboratory, Faculty of Dentistry, Trisakti University, Jl. Kyai Tapa No.260, Jakarta, Indonesia¹¹Prodia Clinical Laboratory, Prodia Tower, Jl. Kramat Raya No.150, Jakarta, Indonesia

*Corresponding author. E-mail: ferrysandra@gmail.com

Abstract

BACKGROUND: Syntax score is a lesion-based angiographic scoring system, originally devised to grade the anatomic complexity of coronary artery disease (CAD) and thereby facilitate consensus in the study of a diagnostic angiogram between surgeons and interventional cardiologists. Suitable biology marker to predict the severity of CAD and to have a treatment decision for revascularisation by percutaneous coronary intervention (PCI) or bypass surgery, is still unknown. Which biology marker can be a predictor for higher Syntax score remains unknown as well.

METHODS: Severity of 105 patients who underwent coronary angiography, were evaluated with Syntax score. Laboratory tests were conducted for routine blood test, serum glutamic oxaloacetic transaminase (SGOT), serum glutamic pyruvic transaminase (SGPT), high-sensitivity C-reactive protein (hs-CRP), total cholesterol, low-density lipoprotein (LDL) cholesterol, high-density lipoprotein (HDL) cholesterol, triglycerides, kidney function, creatinine and hemoglobin A1c (HbA1c).

RESULTS: Ureum had significant correlation with Syntax score, $p=0.016$ and 95% confidence interval (CI): 0.05-0.55. Result of logistic regression analysis showed that HDL and haematocrit had significant correlation with Syntax score, with cut off point 34 ($p=0.02$, $p=0.0033$, respectively).

Abstrak

LATAR BELAKANG: Skor *Syntax* adalah sistem skor angiografi yang menentukan keparahan penyakit jantung koroner (PJK) secara anatomi, sehingga memfasilitasi konsensus studi angiogram diagnostik antara ahli bedah dan kardiologis intervensional. Marker biologis yang tepat dalam hal memperkirakan keparahan PJK dan menentukan pengobatan revaskularisasi dengan tindakan dilatasi dengan balon dan pemasangan *stent* atau operasi *bypass*, belum diketahui. Biologi marker mana yang bermanfaat dalam penentuan skor *Syntax* yang tinggi, juga belum diketahui.

METODE: Keparahan dari 105 pasien yang menjalani angiografi koroner, dievaluasi dengan perhitungan skor *Syntax*. Pemeriksaan laboratorium yang dilakukan adalah pemeriksaan darah rutin, *high-sensitivity C-reactive protein* (hs-CRP), total kolesterol, kolesterol *low-density lipoprotein* (LDL), kolesterol *high-density lipoprotein* (HDL), trigliserida, fungsi ginjal, kreatinin dan hemoglobin A1c (HbA1C).

HASIL: Ureum berkorelasi signifikan dengan skor *Syntax*, $p=0.016$ dan 95% *confidence interval* (CI): 0.05-0.55. Hasil analisa regresi logistik memperlihatkan HDL dan hematokrit berkorelasi signifikan dengan skor *Syntax*, dengan *cut off point* 34 ($p=0.02$, $p=0.0033$, secara berurutan). Berdasarkan

Based on multi-variate analysis, Syntax score $34 = 16$ (HDL ≤ 40 + Ureum ≥ 23) + 10 (Neutrophil/Lymphocyte ≥ 3.789) + 10 (HbA1c ≥ 7) + 13 (monocyte $\geq 9\%$).

CONCLUSION: HDL cholesterol, ureum, neutrophil-lymphocyte ratio, HbA1c and monocyte altogether can predict the higher Syntax score.

KEYWORDS: syntax, neutrophil, lymphocyte, HDL, cholesterol, ureum, HbA1c, monocyte

Indones Biomed J. 2014; 6(2): 107-12

analisa multivariat, skor Syntax $34 = 16$ (HDL ≤ 40 + Ureum ≥ 23) + 10 (Neutrofil/Limfosit ≥ 3.789) + 10 (HbA1c ≥ 7) + 13 (monosit $\geq 9\%$).

KESIMPULAN: Kolesterol HDL, ureum, rasio neutrofil-limfosit, HbA1C dan monosit secara bersama dapat menentukan skor *Syntax* yang tinggi.

KATA KUNCI: *syntax*, netrofil, limfosit, HDL, kolesterol, ureum, HbA1c, monosit

Introduction

Coronary artery disease (CAD) is a disease pertaining to plaque atherosclerosis which could tighten as well as block the blood vessel of artery coronary (1-3). Myocardium will not have enough blood supply as well as oxygen for myocardium contraction, leading to ischemic, heart attack and mortality. To prevent this, revascularization will be needed either with percutaneous coronary intervention (PCI) or bypass surgery, depending on the severity of coronary disease.

Syntax score is a lesion-based angiographic scoring system, originally devised to grade the anatomic complexity of CAD and thereby facilitate consensus in the study of a diagnostic angiogram between surgeons and interventional cardiologists.(1,2) Suitable biology marker to predict the severity of CAD and to have a treatment decision for revascularization by percutaneous coronary intervention (PCI) or bypass surgery, is still unknown.(4-8)

Methods

Subjects Recruitment and Selection

Subjects were recruited from Medistra hospital who underwent coronary angiography. Patients with infectious diseases, severe liver, renal disease and revascularization history, were excluded. Upon explanation, all subjects had to sign written informed consent. This study was approved by Ethics Committee of Faculty of Medicine, Hasanuddin University.

Subjects Examination and Calculation of Syntax Score

Subjects enrolled in this study underwent detailed clinical and angiographic examinations. Subjects were asked for

their medical history, smoking habit, hypertension, diabetes mellitus, hypercholesterolemia and family medical history. Calculation of Syntax score was conducted with online application at www.syntaxscore.com/calculation.(5)

Laboratory Test

Blood samples were collected in the morning after overnight fasting for 12 hours. Routine blood test was performed, and serum was separated not more than 2 hours after blood collection. Serum levels of serum glutamic oxaloacetic transaminase (SGOT), serum glutamic pyruvic transaminase (SGPT), ureum, creatinine, high-sensitivity C-reactive protein (hs-CRP), total cholesterol, low-density lipoprotein (LDL) and high-density lipoprotein (HDL) cholesterol and triglyceride were assayed using auto-analyser Cobas (Roche Diagnostics, Indianapolis, IN, USA).

Statistical Analysis

Statistical Analysis were performed using Strata (College Station, Tx, USA). Regression linear, logistic regression and multivariate tests were conducted with significant level at $p < 0.05$.

Results

Total 105 patients were enrolled in this study from January to December 2013, only 16 patients (15%) had asymptomatic angina. Syntax score > 34 were 27 patients (25.7%). Patients with hypertension and diabetes mellitus had higher Syntax score than patients without history of hypertension and diabetes mellitus, but not significant (hypertension, $p = 0.227$; diabetes mellitus, $p = 0.386$).

Regression linear analysis in Table 2 showed that only ureum had significant correlation with Syntax score, $p = 0.016$ and 95% confidence interval (CI): 0.05-0.55. We

Table 1. Distribution of variables.

No.	Variable	Mean	SEM*	Min	Max
1	Age (year old)	57.6	0.97	33	78
2	high (cm)	166.58	0.85	145	185
3	Body weight (kg)	73.44	1.32	47	125
4	Body mass index	26.4	0.39	18.82	41.76
5	Body surface area (m ²)	1.81	0.02	1.39	2.36
6	Haemoglobin (mg/dL)	14.05	0.15	9.8	17.4
7	Haematocrit (%)	41.02	0.51	11.7	52
8	Leukocyte	10.85	2.44	2.16	262
9	Thrombocyte	249.13	5.64	134	526
10	Basophil (%)	552381	0.12	0	9
11	Eosinophil (%)	2.72	0.25	0	11
12	Band neutrophil (%)	0.29	0.1	0	5
13	Segmented neutrophil (%)	62.98	0.97	43	89
14	Lymphocyte (%)	26.44	0.83	7	47
15	Monocyte (%)	7.36	0.38	0	38
16	Total cholesterol (mg/dL)	193.62	4.34	105	324
17	HDL (mg/dL)	44.94	1.68	25	183
18	LDL (mg/dL)	121.31	1.06	28	292
19	Triglyceride (mg/dL)	155.31	6.65	39	405
20	Ureum (mg/dL)	26.8	0.97	9	74
21	Creatinine (mg/dL)	0.94	0.02	0.04	1.6
22	Uric acid (mg/dL)	6.28	0.23	2.9	24
23	SGOT (U/L)	24.49	1.49	7	152
24	SGPT (U/L)	27.56	1.54	6	104
25	Fasting blood sugar (mg/dL)	107.99	3.41	28	352
26	LV-EF (%)	56.5	1.03	22	78
27	HbA1c (%)	6.81	0.1	4.8	10.1
28	hs-CRP (mg/L)	7.51	1.79	0.1	119
29	Syntax Score	23.77	1.25	3	62

*SEM: Standard Error Mean (Standard Deviation/ \sqrt{n})**Table 2. Regression linear test for Syntax score with cut off point 34.**

No.	Variable	F	R2	t	p*	95% CI
1	Total cholesterol	0.17	0.0016	-0.41	0.685	(-0.81; 0.53)
2	LDL cholesterol	0	0	0.01	0.991	(-0.06; 0.06)
3	HDL cholesterol	0.69	0.0066	-0.83	0.409	(-0.206; 0.084)
4	Creatinin	0.18	0.0017	0.42	0.676	(-8.56; 13.15)
5	Ureum	5.94	0.0545	2.44	0.016	(0.05; 0.55)
6	hs-CRP	0.18	0.0017	-0.42	0.672	(-0.16; 0.107)
7	SGOT	0.61	0.0059	-0.78	0.436	(-0.22; 0.09)
8	SGPT	1.22	0.0117	-1.1	0.272	(-0.246; 0.07)
9	Body mass index	1.31	0.0125	-1.14	0.255	(-0.978; 0.262)
10	Neutrophil-lymphocyte ratio	0.25	0.0024	0.5	0.619	(-0.02; 3.61)

*Regression Linear test

Table 3. Logistic Regression test for Syntax score with cut off point 34.

No.	Variable	p	Odd Ratio	95% CI
1	Total cholesterol	0.927	1.0004	0.99; 1.01
2	LDL cholesterol	0.35	1.005	0.99; 1.015
3	HDL cholesterol	0.02	0.936	0.885; 0.989
4	Left ventricle ejection fraction	0.243	0.975	0.93; 1.01
5	hs-CRP (impute)	0.957	1.009	0.72; 1.41
6	Creatinine	0.255	0.327	0.048; 2.23
7	Ureum (impute)	0.092	1.038	0.99; 1.08
8	Hypertension	0.598	1.26	0.52; 3.04
9	Age	0.324	1.02	0.97; 1.07
10	Hypercholesterolemia	0.523	0.657	0.18; 2.38
11	Smoker	0.412	1.51	0.56; 4.04
12	Diabetes Mellitus	0.248	1.696	0.69; 4.15
13	Family history	0.662	1.48	0.255; 8.57
14	Triglyceride (impute)	0.961	0.975	0.36; 2.63
15	Body mass index	0.311	0.94	0.83; 1.05
16	Neutrophil-lymphocyte ratio	0.214	1.13	0.93; 1.38
17	Basophil	0.147	0.173	0.016; 1.84
18	Haemoglobin	0.072	0.76	0.57; 1.02
19	Haematocrit	0.033	0.903	0.82; 0.99
20	Leukocyte	0.248	1.07	0.95; 1.21
21	Thrombocyte	0.329	1.003	0.99; 1.01
22	Eosinophil	0.576	0.95	0.79; 1.13
23	Band neutrophil	0.654	1.09	0.73; 1.65
24	Segmented neutrophil	0.437	1.01	0.97; 1.06
25	Lymphocyte	0.295	0.97	0.92; 1.02
26	Monocyte	0.235	1.07	0.95; 1.19
27	Uric acid	0.434	0.91	0.72; 1.15
28	SGOT	0.65	0.99	0.95; 1.02
29	SGPT	0.989	1.0001	0.97; 1.02
30	Fasting blood sugar	0.401	1.005	0.99; 1.01
31	HbA1c	0.053	1.51	0.99; 2.3
32	Chest pain	0.541	1.31	0.54; 3.17

Table 4. Multivariate predictor Syntax Score >34.

Variable	Cut Off	p	Odd Ratio	Coefficients	z	Score
HDL	≤40	0.062	2.51	0.9225703	1.87	35
Ureum	≥23	0.021	4.15	1.423749	2.31	43
Neutrophil/lymphocyte	≥3.789	0.186	2.24	0.805509	1.32	25
HbA1c	≥7	0.598	1.32	0.2834972	0.53	10
Monocyte	≥9%	0.116	2.31	0.8379034	1.57	29








Figure 1. Roctab of variables (HDL cholesterol, ureum, neutrophil/lymphocyte ratio, HbA1c and Monocyte) with Syntax score 34. A: HDL cholesterol: Cut off point 40; Sensitivity 40.74%; Specificity 37.18%; Likelihood ratio +0.6485; Likelihood ratio -1.5939. **B:** Urem: Cut off point 23 mg/dl; Sensitivity 85.19%; Specificity 41.03%; Likelihood ratio +1.44; Likelihood ratio -0.3611. **C:** Neutrophil-lymphocyte ratio: Cut off point 3.79; Sensitivity 29.63%; Specificity 85.9%; Likelihood ratio +2.1; Likelihood ratio -0.819. **D:** HbA1c: Cut off point 6.5 %; Sensitivity 74.9%; Specificity 41.03%; Likelihood ratio +1.256; Likelihood ratio -0.6319. **E:** Monocyte: Cut off point 9; Sensitivity 37.04; Specificity 80.77%; Likelihood ratio +1.9259; Likelihood ratio -0.7795.

used Syntax Score 34 as cut off point as reported previously. (9)

Result of logistic regression analysis showed that HDL and haematocrit had significant correlation with Syntax score with cut off point 34 ($p=0.02$, $p=0.0033$, respectively)

(Table 3).

Based on multi-variate analysis, Syntax score 34 = 16 (HDL ≤ 40 + Urem ≥ 23) + 10 (Neutrophil/Lymphocyte ≥ 3.789) + 10 (HbA1c ≥ 7) + 13 (monocyte $\geq 9\%$).




Figure 2. Roctab scoring to predict Syntax score >34. Score model can predict 73.53%; Cut off point 70; Sensitivity 70.37%; Specificity 65.38%; Likelihood ratio +2.0329; Likelihood ratio -0.4532; Risk 67%.

Discussion

Based on our current result, we found that there was not a significant correlation between neutrophil-lymphocyte ratio and Syntax score. This could be related to the ongoing treatment performed to our subjects, such as antiplatelet, aspirin, clopidogrel, also statins that decrease lipid level.

For neutrophil-lymphocyte ratio, different with our results, other reports (10-17) showed that acute coronary syndrome patients with ST elevation nor without ST elevation, will have neutrophil-lymphocyte ratio as an independent factor for the mortality(18-20). In accordance to our results, Arbel *et al.* reported patients with >3 neutrophil-lymphocyte ratio, will have more severity than the patients with ratio <3.(20) In our study, only 30% subject with angina pectoris class III and 20% angina pectoris class IV.

In our study, we found that history of hypertension and diabetic mellitus were positively correlated with syntax score. Although the correlation was not significant, but relative higher in patient with syntax score ≥ 34 . Therefore patients with the history of hypertension and diabetic mellitus, will have higher syntax scores. These results are in accordance to reports showing that hypertension and diabetes mellitus are risk factors. Eventually the risks can induce dysfunction of coronary arteries endothelium and activate LDL cholesterol in the blood vessel wall, leading to plaque formation, then tightness and blockage of coronary arteries will be occurred (1,21-23). In our results, there was no correlation between

Syntax score and hypercholesterolemia, since our patients has already been treated with lipid reducing medicine. We are aware that there are limitations in our study, therefore further research should be pursued, especially in functional Fractional Flow Reserve (FFR), Euro Score and Syntax score II (24).

Conclusion

Marker biology of HDL cholesterol, Ureum, neutrophil/lymphocyte ratio, HbA1c and Monocyte together will be predictor of higher syntax score.

Acknowledgement

We thanks to Medistra hospital for the support in this study.

References

1. Packard RR, Libby P. Inflammation in atherosclerosis: from vascular biology to biomarker discovery and risk prediction. *Clin Chem.* 2008; 54: 24-38.
2. Rose R. Atherosclerosis - An Inflammatory Disease. *N Eng J Med.* 1999; 340: 115-26.
3. Avanzas P, Arroyo-Espiguero R, Cosin-Sales J, Aldama G, Pizzi C, Quiles J, *et al.* Markers of inflammation and multiple complex stenoses (pancoronary plaque vulnerability) in patients with non-ST segment elevation acute coronary syndromes. *Heart.* 2004; 90: 847-52.
4. Ong AT, Serruys PW, Mohr FW, Morice MC, Kappetein AP, Holmes DR Jr, *et al.* The SYnergy between percutaneous coronary intervention with TAXus and cardiac surgery (SYNTAX) study: design, rationale, and run-in phase. *Am Heart J.* 2006; 151: 1194-204.
5. Garg S, Sarno G, Garcia-Garcia HM, Girasis C, Wykrzykowska J, Dawkins KD, *et al.* A new tool for the risk stratification of patients with complex coronary artery disease: the Clinical SYNTAX Score. *Circ Cardiovasc Interv.* 2010; 3: 317-26.
6. Serruys PW, Morice MC, Kappetein AP, Colombo A, Holmes DR, Mack MJ, *et al.* Percutaneous coronary intervention versus coronary-artery bypass grafting for severe coronary artery disease. *N Engl J Med.* 2009; 360: 961-72.
7. Girasis C, Garg S, Räber L, Sarno G, Morel MA, Garcia-Garcia HM, *et al.* SYNTAX score and Clinical SYNTAX score as predictors of very long-term clinical outcomes in patients undergoing percutaneous coronary interventions: a substudy of SIRolimus-eluting stent compared with pacliTAXel-eluting stent for coronary revascularization (SIRTAX) trial. *Eur Heart J.* 2011; 32: 3115-27.
8. Yadav MI, Palmerini T, Caixeta A, Madhavan MV, Sanidas E, Kirtane AJ, *et al.* Prediction of coronary risk by SYNTAX and derived scores: synergy between percutaneous coronary intervention with taxus and cardiac surgery. *J Am Coll Cardiol.* 2013; 62: 1219-30.

9. Capodanno D, Capranzano P, Di Salvo ME, Caggegi A, Tomasello D, Cincotta G, *et al.* Usefulness of SYNTAX score to select patients with left main coronary artery disease to be treated with coronary artery bypass graft. *JACC Cardiovasc Interv.* 2009; 2:731-8.
10. Duffy BK, Gurm HS, Rajagopal V, Gupta R, Ellis SG, Bhatt DL. Usefulness of an elevated neutrophil to lymphocyte ratio in predicting long-term mortality after percutaneous coronary intervention. *Am J Cardiol.* 2006; 97: 993-6.
11. Collier BS. Leukocytosis and ischemic vascular disease morbidity and mortality: is it time to intervene? *Arterioscler Thromb Vasc Biol.* 2005; 25: 658-70.
12. Sahin S, Sarikaya S, Akyol L, Altunkas F, Karaman K. Evaluation of Neutrophil to Lymphocyte ratio as an Indicator of Presence of Coronary Artery Disease in Diabetic Patients. *Natl J Med Res.* 2013; 3: 300-3.
13. Zazula AD, Précoma-Neto D, Gomes AM, Krukalis H, Barbieri GF, Forte RY, *et al.* An assessment of neutrophils/lymphocytes ratio in patients suspected of acute coronary syndrome. *Arq Bras Cardiol.* 2008; 90: 31-6.
14. Poludasu S, Cavusoglu E, Khan W, Marmur JD. Neutrophil to lymphocyte ratio as a predictor of long-term mortality in African Americans undergoing percutaneous coronary intervention. *Clin Cardiol.* 2009; 32: E6-10. doi: 10.1002/clc.20503.
15. Shen XH, Chen Q, Shi Y, Li HW. Association of neutrophil/lymphocyte ratio with long-term mortality after ST elevation myocardial infarction treated with primary percutaneous coronary intervention. *Chin Med J (Engl).* 2010; 123: 3438-43.
16. Arbel Y, Finkelstein A, Halkin A, Birati EY, Revivo M, Zuzut M, *et al.* Neutrophil/lymphocyte ratio is related to the severity of coronary artery disease and clinical outcome in patients undergoing angiography. *Atherosclerosis.* 2012; 225: 456-60.
17. Chen J, Chen MH, Li S, Guo YL, Zhu CG, Xu RX, *et al.* Usefulness of the Neutrophil-to-Lymphocyte Ratio in Predicting the Severity of Coronary Artery Disease: A Gensini Score Assessment. *J Atheroscler Thromb.* 2014; [Epub ahead of print].
18. Jurewitz DL, Pessegueiro A, Zimmer R, Bhatia R, Tobis J, Lee MS. Preprocedural white blood cell count as a predictor of death and major adverse cardiac events in patients undergoing percutaneous coronary intervention with drug-eluting stents. *J Invasive Cardiol.* 2009; 21: 202-6.
19. Horne BD, Anderson JL, John JM, Weaver A, Bair TL, Jensen KR, *et al.* Which white blood cell subtypes predict increased cardiovascular risk? *J Am Coll Cardiol.* 2005; 45: 1638-43.
20. Akpek M, Kaya MG, Lam YY, Sahin O, Elcik D, Celik T, *et al.* Relation of neutrophil/lymphocyte ratio to coronary flow to in-hospital major adverse cardiac events in patients with ST-elevated myocardial infarction undergoing primary coronary intervention. *Am J Cardiol.* 2012; 110: 621-7.
21. Arbel Y, Finkelstein A, Halkin A, Birati EY, Revivo M, Zuzut M, *et al.* Neutrophil/lymphocyte ratio is related to the severity of coronary artery disease and clinical outcome in patients undergoing angiography. *Atherosclerosis.* 2012; 225: 456-60.
22. Muller WA, Weigl SA, Deng X, Phillips DM. PECAM-1 is required for transendothelial migration of leukocytes. *J Exp Med.* 1993; 178: 449-60.
23. Springer TA, Cybulsky MI. Traffic signals on endothelium for leukocytes in health, inflammation, and atherosclerosis. In: Fuster V, Ross R, Topol EJ, editors. *Atherosclerosis and Coronary Artery Disease.* Philadelphia: Lippincott-Raven; 1996. p. 511-38.
24. Farooq V, van Klaveren D, Steyerberg EW, Meliga E, Vergouwe Y, Chieffo A, *et al.* Anatomical and clinical characteristics to guide decision making between coronary artery bypass surgery and percutaneous coronary intervention for individual patients: development and validation of SYNTAX score II. *Lancet.* 2013; 381: 639-50.