


Contents lists available at [ScienceDirect](http://www.sciencedirect.com)

Asian Pacific Journal of Reproduction

journal homepage: www.apjr.netOriginal research <http://dx.doi.org/10.1016/j.apjr.2016.07.015>Effects of *Labisia pumila* on oxidative stress in rat model of post-menopausal osteoporosisNurdiana Nurdiana^{1*}, Nelly Mariati², Noorhamdani Noorhamdani³, Bambang Setiawan⁴, Nicolaas Budhiparama⁵, Zairin Noor⁶¹Department of Pharmacology, Faculty of Medicine, University of Brawijaya, Malang, East Java, Indonesia²Midwifery Master Study Programme, Faculty of Medicine, Brawijaya University, Malang, East Java, Indonesia³Department of Microbiology, Faculty of Medicine, University of Brawijaya, Malang, East Java, Indonesia⁴Research Center for Toxicology, Cancer, and Regenerative Medicine, Department of Medical Chemistry and Biochemistry, Faculty of Medicine, University of Lambung Mangkurat, Banjarmasin, South Kalimantan, Indonesia⁵Budhiparama Institute of Hip and Knee Research and Education Foundation for Arthroplasty, Sports Medicine and Osteoporosis, Jakarta, Indonesia⁶Research Center for Osteoporosis, Department of Orthopaedic and Traumatology, Ulin General Hospital, Faculty of Medicine, University of Lambung Mangkurat, Banjarmasin, South Kalimantan, Indonesia

ARTICLE INFO

Article history:

Received 3 Jul 2016

Received in revised form 14 Jul 2016

Accepted 14 Jul 2016

Available online 6 Aug 2016

Keywords:

Ovariectomy

Oxidative stress

Oxidative damage

Lipid peroxidation

Antioxidants

ABSTRACT

Objective: To analyze the effects of *Labisia pumila* (*L. pumila*) extract on the markers of oxidative stress in ovariectomized rats.**Methods:** Twenty-five female Wistar rats were divided into five treatment groups ($n = 5$): the control group, the ovariectomized group, the ovariectomized group treated with *L. pumila* extract of various doses (10 mg/kg, 20 mg/kg, and 40 mg/kg). *L. pumila* extract was administered daily for 8 weeks.**Results:** The levels of malondialdehyde and superoxide dismutase (SOD) were subjected to spectrophotometric analysis. Serum levels of malondialdehyde and SOD were significantly higher in the group of rat model of post-menopausal osteoporosis than that of the control group ($P < 0.05$). All doses of *L. pumila* administered significantly decreased the serum levels of malondialdehyde relative to the group of rat model of post-menopausal osteoporosis, reaching the levels comparable to those of the control group ($P < 0.05$). The second dose of *L. pumila* significantly decreased the levels of SOD relative to those of the group of rat model of post-menopausal osteoporosis, although it did not reach the levels of SOD of the control group ($P < 0.05$).**Conclusion:** The ethanol extract of *L. pumila* normalizes lipid peroxidation in the rat model of post-menopausal osteoporosis via the mechanism of endogenous antioxidant replacement.

1. Introduction

Osteoporosis is characterized by the loss of bone mass and micro-architectural deterioration of bones, leading to bone fragility and increased risk of fracture [1]. Pathophysiology of the ovary-related loss of bone mass is complex; it cannot be explained simply by the increase in bone resorption or decreased bone formation [2]. Bone homeostasis is affected by a reduction in antioxidant defenses against oxidative stress along with increased reactive oxygen species. At the cellular level, a

defect in bone remodeling caused by oxidative stress is associated with reduction in osteoblasts and osteoclasts as well as reduction in bone formation accompanied by increased apoptosis of osteoblasts and osteocytes [3,4]. The model of bilateral ovariectomy resembles accelerated bone loss in postmenopausal women, which is underlain by estrogen deficiency [5]. Bilaterally ovariectomized animals showed an increase in osteoclastic bone resorption and reactive osteoblastic bone formation with the net result of a loss of bone mass [6]. Previous studies demonstrated that the ovariectomized rats had an increase in lipid peroxidation and hydrogen peroxide and a decrease in antioxidants and antioxidant cofactors [7–9].

Labisia pumila (*L. pumila*) is a plant commonly found in Southeast Asia. In Southeast Asia communities this plant

*Corresponding author: Nurdiana Nurdiana, Department of Pharmacology, Faculty of Medicine, University of Brawijaya, Malang, East Java, Indonesia.

Tel: +628123388428

E-mail: farmakodes@gmail.com

Peer review under responsibility of Hainan Medical College.

decoction is used as a health supplement by boiling the roots, leaves, or whole parts of the plant for irregular menstruation, menstrual pain, inducing labor and a tonic for the vagina walls after childbirth [10,11]. Other studies have evaluated the antioxidant contents and benefits of *L. pumila* in rat model of ovariectomy [12–14], but no studies on the control of oxidative stress in the rat model of osteoporosis due to estrogen deficiency. Hence, the purpose of the present study was to analyze the effects of *L. pumila* extract on oxidative stress in ovariectomized rats.

2. Material and methods

2.1. Subjects

Twenty-five female Wistar rats, aged 3–4 months and body weight of 150–200 g, were used as the subjects of this study. After acclimatization to the laboratory conditions, they were randomly divided into five treatment groups: the control group, the ovariectomized group and the ovariectomized group administered with *L. pumila* extract of various doses (10 mg/kg, 20 mg/kg, and 40 mg/kg). *L. pumila* extract was administered for 8 weeks. Dosage was determined on the basis of previous studies stating that the dose with no adverse effects for *L. pumila* was 50 mg/kg [15]. In the course of the study rats were reared under conditions of 12 h of light and 12 h of dark at room temperature of 24 °C. In addition, rats were given standard laboratory feed and access to water *ad libitum*. All research procedures were performed under the ethical guidelines for experimental animals and passed the ethical review of the Research Ethics Committee, Medicine Faculty of Brawijaya University Malang, East Java, Indonesia.

2.2. Procedures for ovariectomy

After anesthesia with ketamine (50 mg/kg) and xylazine (8 mg/kg), twenty rats of the ovariectomized group were subjected to bilateral ovariectomy through a ventral incision. In addition, five rats were subjected to sham surgery [16]. After ovariectomy, wound care was carried out for 10 d and then treatment was given.

2.3. Extract preparation

L. pumila in dry conditions was cut into small pieces and then crushed in a blender and sieved to separate the rough from fine particles. The extract of *L. pumila* was prepared by using the maceration method in which 100 g of *L. pumila* powder is soaked in 900 mL of 96% ethanol for 5–7 d while stirring occasionally. The soaked extract of *L. pumila* was put into the rotavapor until the ethanol solution separates from the active substance. The concentrated extract was then weighed and diluted with distilled water [17]. The ethanol extract of *L. pumila* was administered in accordance with the dosage using an oral feeding tube to the ovariectomized rats.

2.4. Serum collection

At the end of the study, all the rats were anesthetized and their blood was collected from their hearts by cardiac puncture. Furthermore, the blood was centrifuged to obtain serum.

2.5. Analysis of malondialdehyde (MDA) and superoxide dismutase (SOD)

Analysis of serum levels of MDA and SOD was performed according to the protocol in previous studies [18].

2.6. Ethics

This study passed the ethics review of the Ethics Committee of the Faculty of Medicine, Brawijaya University Malang of East Java, Indonesia.

2.7. Statistical analysis

All data were presented in mean \pm SD. Differences in the levels among treatment groups were analyzed by ANOVA using SPSS 16.0 statistical package. A *P* value of <0.05 was considered as statistically significant difference.

3. Results

Figure 1 presents the serum levels of MDA in each experimental group. Serum levels of MDA were significantly higher in the group of rat model of post-menopausal osteoporosis compared to those of the control group ($P < 0.05$). All doses of *L. pumila* significantly reduced the serum levels MDA relative to those of the group of rat model of post-menopausal osteoporosis, reaching the levels comparable to those of the control group ($P < 0.05$).

Figure 2 presents the serum levels of SOD in each experimental group. SOD levels were significantly higher in the group of rat model of post-menopausal osteoporosis than those of the control group ($P < 0.05$). Of the three doses, the second dose of *L. pumila* significantly lowered SOD levels relative to those of the group of rat model of post-menopausal osteoporosis, although it did not reach the levels of SOD of the control group ($P < 0.05$).




Figure 1. Malondialdehyde level in serum of control and experimental groups.

Note: ^a $P < 0.05$ in comparison with control (untreated) group; ^b $P < 0.05$ in comparison with ovariectomized rats (OVX) group; OVX + LP: ovariectomized rats treated by *Labisia pumila* (ng/μL).




Figure 2. Superoxide dismutase level in serum of control and experimental groups.

^a $P < 0.05$ in comparison with control (untreated) group; ^b $P < 0.05$; in comparison with OVX group; OVX + LP: ovariectomized rats treated by *Labisia pumila* (U/mL).

4. Discussion

Under normal conditions, reactive oxygen species (ROS) will be neutralized efficiently by cellular antioxidant defense mechanisms. In many conditions, there is an imbalance of ROS production and antioxidant defenses, leading to cellular destruction and dysfunction [19]. The present study showed that the levels of MDA, the marker of oxidative damage, increased significantly in the groups of rat model of post-menopausal osteoporosis relative to those of the control group ($P < 0.05$). This suggests that ovariectomy induces lipid peroxidation caused by a systemic increase in oxidative stress. All doses of *L. pumila* administered significantly decreased the serum levels of MDA relative to the group of rat model of post-menopausal osteoporosis, reaching the levels comparable to those of the control group ($P < 0.05$). This shows that *L. pumila* is capable of repairing oxidative damage [19,20].

The levels of SOD were significantly higher in the group of rat model of post-menopausal osteoporosis than those of the control group ($P < 0.05$). This finding indicates that under conditions of osteoporosis-induced oxidative stress, there is endogenous antioxidant compensation in the form of increased levels of SOD. Depending on the levels of ROS, a variety of transcription factors sensitive to changes in the redox status will be activated and will coordinate certain biological responses. Low levels of oxidative stress induce Nrf2, a transcription factor implicated in transactivation of genes that encode enzymatic antioxidant activity [21]. Of the three doses, the second dose of *L. pumila* significantly lowered SOD levels relative to those of the group of rat model of post-menopausal osteoporosis, although it did not reach the levels of SOD of the control group ($P < 0.05$). We speculated that the antioxidant levels of *L. pumila* will replace SOD to reduce oxidative stress. This mechanism will restore SOD levels to the basal levels. Previous studies showed that *L. pumila* acts as an antioxidant, which is played by flavonoids, ascorbic acid, beta-carotene, anthocyanin and phenolic compounds [19,20]. Additionally, previous studies also demonstrated the activity of SOD in *L. pumila* [22].

In conclusion, the ethanol extract of *L. pumila* normalizes lipid peroxidation in the rat model of post-menopausal

osteoporosis through the mechanism of endogenous antioxidant replacement.

Conflict of interest statement

The authors declare that they have no competing interest.

References

- [1] Zhao X, Wu Z, Zhang Y, Gao M, Yan Y, Cao P, et al. Locally administered perindopril improves healing in an ovariectomized rat tibial osteotomy model. *PLoS One* 2012; **7**(3): e33228; <http://dx.doi.org/10.1371/journal.pone.0033228>.
- [2] Zhang J, Lazarenko OP, Blackburn ML, Shankar K, Badger TM. Feeding blueberry diets in early life prevent senescence of osteoblast and bone loss in ovariectomized adult female rats. *PLoS One* 2011; **6**(9): e24486; <http://dx.doi.org/10.1371/journal.pone.0024486>.
- [3] Wauquier F, Leotoing L, Coxam V, Guicheux J, Wittrant Y. Oxidative stress in bone remodelling and disease. *Trends Mol Med* 2009; **15**: 468-477.
- [4] Kular J, Tickner J, Chim SM, Xu J. An overview of the regulation of bone remodelling at the cellular level. *Clin Biochem* 2012; **45**: 863-873.
- [5] Folwarczna J, Zych M, Trzeciak HI. Effects of curcumin on the skeletal system in rats. *Pharmacol Rep* 2010; **62**: 900-909.
- [6] de Freitas PHL, Hasegawa T, Takeda S, Sasaki M, Tabata C, Oda K, et al. Eldecacitol, a second-generation vitamin D analog, drives bone minimodeling and reduces osteoclastic number in trabecular bone of ovariectomized rats. *Bone* 2011; **49**: 335-342.
- [7] Lean JM, Davies JT, Fuller K, Jagger CJ, Kirstein B, Partington GA, et al. A crucial role for thiol antioxidants in estrogen-deficiency bone loss. *J Clin Invest* 2003; **112**: 915-923.
- [8] Ha BJ. Oxidative stress in ovariectomy menopause and role of chondroitin sulfate. *Arch Pharm Res* 2004; **27**: 867-872.
- [9] Karimi E, Jaafar HZ, Ahmad S. Phytochemical analysis and antimicrobial activities of methanolic extracts of leaf, stem and root from different varieties of *Labisia pumila* Benth. *Molecules* 2011; **16**: 4438-4450.
- [10] Fazliana M, Nazaimoon WMW, Gu HF. *Labisia pumila* extract regulates body weight and adipokines in ovariectomized rats. *Maturitas* 2009; **62**: 91-97.
- [11] Ali Z, Khan IA. Chemical constituents of *Labisia pumila* (Kacip Fatimah). *Planta Med* 2009; **75**: 40-45.
- [12] Fathilah SN, Shuid AN, Mohamed N, Muhammad N, Soelaiman IN. *Labisia pumila* protects the bone of estrogen-deficient rat model: a histomorphometric study. *J Ethnopharmacol* 2012; **142**(1): 294-299.
- [13] Shuid AN, Ping LL, Muhammad N, Mohamed N, Soelaiman IN. The effects of *Labisia pumila* var. *alata* on bone markers and bone calcium in a rat model of post-menopausal osteoporosis. *J Ethnopharmacol* 2011; **133**(2): 538-542.
- [14] Sing GD, Ganjoo M, Youssouf MS, Koul A, Sharma R, Singh S, et al. Sub-acute toxicity evaluation of an aqueous extract of *Labisia pumila*, a Malaysian herb. *Food Chem Toxicol* 2009; **47**(10): 2661-2665.
- [15] Yalin S, Sagir O, Comelekoglu U, Berkoz M, Eroglu P. Strontium ranelate treatment improves oxidative damage in osteoporotic rat model. *Pharmacol Rep* 2011; **63**: 396-402.
- [16] Fard SG, Shamsabadi FT, Emadi M, Meng GY, Muhammad K, Mohamed S. Ethanolic extract of *Eucheuma cottonii* promotes *in vivo* hair growth and wound healing. *J Anim Vet Adv* 2011; **10**(5): 601-605.
- [17] Draper H, Hadley M. Malondialdehyde determination as index of lipid peroxidation. *Methods Enzymol* 1990; **186**: 421-431.
- [18] Halliwell B, Gutteridge JMC. *Free radical in biology and medicine*. 3rd ed. Oxford: University Press; 1999.

- [19] Mohamad N, Mahmood M, Mansor H. Antioxidative properties of leaf extracts of a popular Malaysian herb, *Labisia pumila*. *J Med Plants Res* 2009; **3**: 217-223.
- [20] Huang J, Zhang H, Shimizu N, Takeda T. Triterpenoid saponins from *Arsidia mamillata*. *Phytochemistry* 2000; **54**: 817-822.
- [21] Glorie G, Legrand-Poels S, Piette J. NF- κ B activation by reactive oxygen species: fifteen years later. *Biochem Pharmacol* 2006; **72**: 1493-1505.