



PEDIATRIC UROLOGY CASE REPORTS

ISSN 2148-2969

<http://www.pediatricurologycasereports.com>

A rare form of crossed nonfused renal ectopia, multicystic dysplasia: A case report

**Carolina Talini¹, Leticia Alves Antunes¹, Bruna Cecília Neves de Carvalho¹,
Jéssica Romanelli², André Luis Fortes Alves¹, Rafaela Galli³, Karin Barszcz³,
Ricardo Possagno³, Even Mol³, Luís Fernando Garbuio³, Ayrton Alves Aranha
Junior^{1,2,3}**

¹*Pediatric Surgery, Hospital Pequeno Príncipe, Curitiba/PR, Brasil*

²*Universidade Federal do Paraná, Curitiba/PR, Brasil*

³*Universidade Estadual de Ponta Grossa, Brasil*

ABSTRACT

We report a 6 months-old male who presented a rare phenomenon of nonfused crossed renal ectopia associated with multicystic dysplasia. Male, prenatally diagnosed with cystic lesion in right hypocondria, presented as palpable mass in right upper abdominal quadrant at birth. Detailed radiologic evaluation including laparotomy confirmed a diagnosis of multicystic dysplasia in a crossed non-fused left ectopic kidney and leading to partial right ureteral obstruction, treated by surgical removal of the lesion.

Key Words: Crossed non-fused renal ectopia; multicystic dysplasia.

Copyright © 2017 pediatricurologycasereports.com

Corresponding Author: Dr. Ayrton Alves Aranha Junior. *Pediatric Surgery, Hospital Pequeno Príncipe, Curitiba/PR, Brasil*

Universidade Federal do Paraná, Curitiba/PR, Brasil

E mail: aranha43@hotmail.com

Accepted for publication: 11 February 2017

Introduction

Crossed renal ectopia is a congenital anomaly consisting of the transposition of a kidney to the opposite side [1], simple renal ectopia represents a defective ascension of the affected kidney during embryogenesis [2]. Less than 10% of the crossed ectopic kidneys are non-

fused [2]. Additionally, multicystic dysplasia in a crossed, fused or non-fused, ectopic kidney is rare [3].

The majority of pediatric patients are asymptomatic but non-specific symptoms such as abdominal pain, palpable mass, hematuria and dysuria can be present [4]. In some reported cases the patient develops additional symptoms such as; infection, renal calculi or urinary obstruction [1]. Up to 30% of cases can be incidentally diagnosed [5].

Herein is reported a rare form of nonfused crossed left renal ectopia where the patient

presented a multicystic dysplastic ectopic kidney.

Case report

A 6 months-old male was referred to the Pediatric Urology Department for evaluation of a multicystic dysplastic left kidney. The past story of the patient was as follows: He was diagnosed with a cystic lesion in the left abdomen during 22 weeks gestational ultrasound. As he presented with no other associated anomaly and normal amniotic fluid level, gestation was taken to term. Physical examination after birth was demonstrated a palpable mass in right abdomen, painless and with low mobility. Urinary volume during the first 48 hours of life was normal, as well as renal function laboratory evaluation.

Ultrasound scan of the Urinary Tract was confirmed a multicystic lesion in all right hemiabdomen associated with minimal right pyelocaliceal dilatation. The left kidney was not identified. In order to clarify the anatomy as well as to plan any surgical intervention, magnetic resonance (MR) urography was performed. MR urography showed multicystic left kidney associated with rotation anomaly crossing the midline plan. The left kidney was functionally excluded and was also compressing the proximal right ureter causing a partial obstructive renogram pattern.

At laparotomy, the right kidney was normal and the left kidney was accessed via transperitoneal. The retroperitoneal cystic mass, which was not fused with the right kidney, was resected [Fig. 1]. The procedure presented uneventful and the patient was discharged within 24 h. Histopathological examination of the mass revealed multicystic dysplasia of the kidney. Six months after surgery he is asymptomatic, with a control

ultrasound showing no dilatation in the right kidney or ureter.




Fig. 1. Multicystic ectopic kidney and orthotopic ureter presenting discrete dilation cause by multicystic crossed kidney.

Discussion

Crossed renal ectopia can be anatomically differentiated into four groups: A) Crossed renal ectopia with fused kidneys (the most common type – 90%), B) Crossed renal ectopia without fusion (rare), C) solitary Crossed renal ectopia (very rare) and D) unfused bilateral Crossed renal ectopia (also very rare) [5]. The ectopic kidney is usually located below the orthotopic kidney and occurs more frequently in males. Left-to-right ectopia represent 50-75% of cases [6].

Diverse associated urological malformations have been described, most commonly vesicoureteral reflux (VUR), but also ureteropelvic junction obstruction, megaureter, hypospadias, cryptorchism, urethral valve and cystic dysplasia [2]. So far, the presence of an ectopic kidney demands investigation for other anomalies, specially VUR [1,4].

Multicystic dysplasia in a crossed-fused or non-fused ectopic kidney has been rarely described in the literature [3,7]. In this particular case, not only this rare association was detected, but also a partial obstruction of the orthotopic pyelourethral system.

Imaging investigation is extremely important during the investigation and surgical planning of complex anomalies [8,9]. In this very case, ultrasound study was initially used, followed by voiding urethrocytogram, in order to exclude VUR. A DMSA scintigraphy was able to detect renal exclusion of the multicystic kidney. The use of MR imaging was considered to clearly define anatomy and improve surgical planning. Not used so frequently, especially because of mandatory general anesthesia and contraindication in patient with compromised renal function [4,9], in our opinion this imaging modality was very helpful for the case.

Besides the lack of consensus about surgical treatment of multicystic dysplastic kidney [10], the patient required resection because of the risk for the orthotopic kidney.

Conclusion

Crossed renal ectopia is a rare malformation, varying from 1:20000 to 1:7000 live births. Its association with dysplastic multicystic kidney is even rarer. This paper presents, in our view, the first case with association of both malformations leading to contralateral partial ureter obstruction.

Acknowledgements

The author(s) declare that they have no competing interest and financial support.

References

- [1] Ratola A, Almiro MM, Lacerda Vidal R, Neves N, Bicho A, Figueiredo S. Crossed Renal Ectopia without Fusion: An Uncommon Cause of Abdominal Mass. *Case Rep Nephrol.* 2015;2015:679342.
- [2] van den Bosch CM, van Wijk JA, Beckers GM, van der Horst HJ, Schreuder MF, Bökenkamp A. Urological and nephrological findings of renal ectopia. *J Urol.* 2010;183(4):1574-8.
- [3] Evans WP, Sumner TE, Lorentz WB Jr, Resnick MI. Association of crossed fused renal ectopia and multicystic kidney. *J Urol.* 1979;122(6):821-2.
- [4] Siegel RL, Rosenfeld DL, Leiman S. Complete regression of a multicystic dysplastic kidney in the setting of renal crossed fused ectopia. *J Clin Ultrasound.* 1992. 20(7):466-469.
- [5] Ramaema DP, Moloantoa W, Parag Y. Crossed Renal Ectopia without Fusion-An Unusual Cause of Acute Abdominal Pain: A Case Report. *Case Rep Urol.* 2012;2012:728531.
- [6] Bhojwani N, Hartmann JB, Ahmed M, Morgan BS, Davidson JC. Management of ureteral obstruction in crossed fused renal ectopia: a case report. *Can Urol Assoc J.* 2014; 8(9-10):52-754.
- [7] Nussbaum AR, Hartman DS, Whitley N, McCauley RG, Sanders RC. Multicystic dysplasia and crossed renal ectopia. *AJR Am J Roentgenol.* 1987;149(2):407-10.
- [8] Mugeiren MMA. Association of multicystic dysplasia and crossed nonfused renal ectopia: a case report. *Saudi J Kidney Dis Transplant.* 1997; 8(2): 148-151.
- [9] Tang L, Koshy J, Spevak MR, Benson JE, Bosemani T. Diagnosis of rare association of orthotopic multicystic dysplasia with

crossed fused renal ectopia. *Case Rep Urol*. 2014;2014:140850.

- [10] Tang L, Koshy J, Spevak MR, Benson JE, Bosemani T. Diagnosis of rare association of orthotopic multicystic dysplasia with crossed fused renal ectopia. *Case Rep Urol*. 2014;2014:140850.

