



A Review on Fungal Fermented Cassava Pulp as a Cheap Alternative Feedstuff in Poultry Ration

Sugiharto Sugiharto

Department of Animal Science, Faculty of Animal and Agricultural Sciences, Diponegoro University, Semarang, Central Java, 50275, Indonesia

Corresponding author's Email: sgh_undip@yahoo.co.id

Received: 07 Jan 2019

Accepted: 11 Feb 2019

ABSTRACT

In order to reduce the production cost, cassava pulp has been incorporated in chicken diets as an energy source. However, the use of such agro-industrial by-product may be confined by its higher fibre and lower protein contents. Improving the nutritional characteristics (lowering fibre and increasing protein content) through fungal solid state fermentation may be conducted to increase the inclusion level of cassava pulp in chicken rations. Apart from an energy source, fungal fermented cassava pulp (FCP) may also exert a beneficial effect on intestinal health of chickens, although further studies are needed to explore the functional benefit of FCP on chicken health.

Keywords: Chicken diet, Energy source, Fermented tapioca by-product, Fungal solid-state fermentation

INTRODUCTION

Feed cost has been a major component in chicken production, representing about 70% of the total cost of production. Generally, chicken diets are composed of several feedstuffs, and that energy-rich feedstuffs such as maize constitute more than half of the diets. Owing to this fact, any increase in the price of maize can potentially increase the chicken production cost and consequently, reduce the profit margin of farmers. Nutritionists are now exploring for unconventional energy-rich feedstuffs that may be used to reduce the proportion of maize in chicken rations. Among the alternative feedstuffs, cassava pulp has frequently been incorporated in chicken rations as a source of energy. On the other hand, the incorporation of such agro-industrial by-product in chicken rations may be limited by the high and low contents of fibre and protein, respectively (Khempaka et al., 2009). Fungal solid state fermentation could be one of the methods to improve the nutritional qualities and thus increase the inclusion level of cassava pulp in chicken rations (Khempaka et al., 2014; Sugiharto et al., 2017a). This present mini review elaborates the potential of fungal Fermented Cassava Pulp (FCP) as an energy-rich alternative feed ingredient for chickens.

Cassava pulp as an energy-rich alternative feedstuff for chickens

As a response to the increased price of maize, nutritionists are currently searching some cheap energy-rich alternative to partially replace the use of maize in chicken diets. As a by-product of tapioca industry, cassava pulp is quite cheap and abundantly available throughout the year, particularly in tropical countries. Cassava pulp contains true metabolizable energy (TME) of about 2,484 kcal/kg energy, therefore, it may be potential as an energy source in chicken rations (Khempaka et al., 2009). On the other hand, it should be noticed that the nutritional qualities of cassava pulp may vary depending on the varieties of cassava, manufacturing process to produce tapioca, the process of drying and contamination (with fibre, sand and soil) (Chauynarong et al., 2015).

Dried cassava pulp has been incorporated as an energy source both in broiler and laying hen diets (Table 1). Nevertheless, its maximum inclusion level differs in broiler and laying hen diets. Dried cassava pulp may be incorporated at higher levels in laying hen (15-20%) as compared to that in broiler diets (Maximum 11%). Similar with this case, King and Zeidler (2004) documented that

dried tomato pomace (contains 31% fibre) may be incorporated at 15% in laying hen diets, while such pomace may only be included in broiler rations at the maximum of 5%. The better capability of laying hens in degrading fibre than of broilers seems to be the reason for the higher fibre digestibility (and thus intake of fibre-rich feedstuffs) in laying hens. Moreover, laying hens seem to have lower energy requirement compared to broiler chickens. This makes laying hens more tolerant to cassava pulp (Diarra and Devi, 2015). Within each broiler and laying hens, the variations in the optimal dietary levels of cassava pulp also existed. In such case, several factors may determine the variation, including the nature (e.g., nutritional composition) of cassava pulp, dietary composition, energy to protein ratio of rations, strains of chickens and other experimental conditions.

Aside from the high energy content, cassava pulp generally contains high and low contents of fibre (13.6%) and protein (1.98%), respectively (Khempaka *et al.*, 2009). According to Morgan and Choct (2016), most of the fibre

in cassava pulp is counted as insoluble fibre. The latter properties may, therefore, limit the use of cassava pulp in poultry rations, as poultry has a very limited capacity in degrading the insoluble fibre. High fibre fraction in cassava pulp may also increase the bulkiness of the diet and therefore limit the capacity of the digestive tract of chickens. This condition may eventually reduce palatability and feed intake in chickens (Khempaka *et al.*, 2009). To elevate the dietary inclusion level of cassava pulp, supplementation of enzymes have been conducted. Khempaka *et al.* (2018) have recently supplemented dried cassava pulp with mixed enzymes containing cellulase, glucanase and xylanase in laying hens. Such enzyme supplementation could increase the inclusion level of dried cassava pulp from 20% (Khempaka *et al.*, 2016) to 30% (Khempaka *et al.*, 2018) in laying hen diets. The activity of enzymes in degrading the fibre fraction of cassava pulp may increase the digestibility of such feed ingredient and thus increase the feed intake of chickens (Khempaka *et al.*, 2018).

Table 1. The use of dried cassava pulp in chicken diets

References	Findings and recommendations
Khempaka <i>et al.</i> (2009)	Dried cassava pulp should be limited to 8% or less, as higher inclusion level may compromise the growth performance of broiler chickens.
Kumsri <i>et al.</i> (2009)	Dietary inclusion of 10% dried cassava pulp reduced weight gain of broiler chickens.
Ali-Mursyid <i>et al.</i> (2010)	Dried cassava pulp may be incorporated in broiler rations at a maximum of 11%, higher inclusion level may be detrimental for the growth of broilers.
Tang <i>et al.</i> (2012)	Feeding dried cassava pulp at a level of 25% resulted in poor growth performance in broiler chickens.
Triprugsachart <i>et al.</i> (2007)	Feeding dried cassava pulp up to 15% represented no detrimental impact on egg production of laying hens.
Chauynarong <i>et al.</i> (2010)	Dietary incorporation of dried cassava pulp up to 15% had no deleterious effect on egg production of laying hens.
Khempaka <i>et al.</i> (2016)	Dried cassava pulp can be included up to 20% in laying hen rations with no negative impact on productive performance, nutrient digestibility, and egg quality.
Khempaka <i>et al.</i> (2018)	Dried cassava pulp supplemented with mixed enzymes (cellulase, glucanase and xylanase) may be included in laying hen feeds up to 30% with no detrimental effects on nutrient digestibility, productive performance and egg quality.

Fungal fermentation to improve the nutritional characteristics of cassava pulp

Fermentation is a simple process using microorganisms to break down the complex substrates into simpler components (Sugiharto and Ranjitkar, 2019). Eventually, the degraded compounds maybe utilized maximally by the chickens. Depending on the microorganisms involved, supplements added, duration of the fermentation process and other fermentation conditions, there is a slight variation in the improvement

of the nutritional quality of cassava pulp particularly with regard to protein and fibre contents. It appears from the documented studies that fungi (Filamentous fungi and yeast) are the most common microorganisms employed to ferment cassava pulp. The definite reason for such preference is not specifically known. There are several traits belong to fungi that may be exploited to improve the nutritional properties of cassava pulp, one of which is its fibrinolytic activity. A study by Mustafa *et al.* (2016) noticed that treatment with fungi was capable of degrading

the insoluble fibre (lignin and hemicellulose) resulting in reduced fibre content of the substrates. Such fibre degradation may be facilitated by the activity of extracellular cellulases produced by the fungi (Bhardwaj et al., 2017). The cellulolytic activity of the fungi may also transform the cellulosic compounds into protein, and therefore increase the protein content of materials. In such case, the conversion of fibre into protein-rich fungal biomass may be responsible for the increased protein content of the fungal fermented products (Asadollahzadeh et al., 2018). In addition, Bayitse et al. (2015) suggested that simple sugars may also be metabolized to protein resulting in an increase in protein content of the fermented products. Apart from the improved fibre and protein contents, fermentation has been known to result in the reduction of Hydrogen Cyanide (HCN) (Diarra and Devi, 2010).

To increase the protein content of FCP, supplementation using urea during fermentation has commonly been conducted (Table 2). During the fermentation process, urea may be used as a nitrogen source for the fungal growth (Bayitse et al., 2015). Such an increase in fungal biomass may thereby increase the protein content of the fungal fermented products. In most cases, fungal fermentation of cassava pulp has been carried out according to the solid-state fermentation method. This fermentation method is characterized by the low content of moisture in the substrates (Sugiharto and Rajitkar, 2019). Indeed, there is no specific reason on why solid state fermentation is more attracted to be employed in the fungal fermentation of cassava pulp. Yet, Gowthaman et al. (2001) suggested that solid-state fermentation may better support the fungi to grow on complex natural solid substrates without substantial pretreatment.

Table 2. Nutritional characteristics of fungal fermented cassava pulp

References	Microorganisms involved in fermentation	Supplement in fermentation	Characteristics of fermented cassava pulp
Lubis et al. (2007)	<i>Aspergillus niger</i>	None	Crude protein content increased from 2.21 to 3.58% and crude fibre increased from 11.2 to 17.0%
Thongkratok et al. (2010)	<i>Aspergillus oryzae</i>	Urea	Protein and amino acid contents increased by 17.4 and 15.1%, respectively
Animashahun et al. (2013)	<i>Penicillium</i> spp.	None	Crude protein increased from 2.39 to 3.25% and crude fibre decreased from 11.4 to 9.63% after 7 days of fermentation
Khempaka et al. (2014)	<i>A. oryzae</i>	Urea	Crude protein increased from 2.02% to 11.8%, whereas crude fibre decreased from 14.6% to 10.6%
Bayitse et al. (2015)	<i>Trichoderma pseudokoningii</i> (ATCC 26801)	Urea and ammonium sulphate	Protein content increased by 48.1% and 36.9% with supplementation of urea and ammonium sulphate, respectively
Sugiharto et al. (2015)	<i>Acremonium charticola</i>	None	Crude fibre decreased from 18.4% to 16.9%, while crude protein did not significantly change (compared to unfermented cassava pulp)
Sugiharto et al. (2015)	<i>Rhizopus oryzae</i>	None	Crude fibre decreased from 18.4% to 17.6%, while crude protein did not significantly change (compared to unfermented cassava pulp)
Sugiharto et al. (2016)	<i>A. charticola</i>	Urea	Crude protein increased from 2.14% to 11.3%, while crude fibre decreased from 25.6% to 20.8%
Sugiharto et al. (2016)	<i>R. oryzae</i>	Urea	Crude protein increased from 2.14% to 12.8%, while crude fibre decreased from 25.6% to 22.7%
Sengxayalth and Preston (2017)	<i>Saccharomyces cerevisiae</i>	Urea and di-ammonium phosphate	Crude protein increased from 9.5 to 18.4% and true protein increased from 2 to 12% dry matter
Okathok et al. (2018)	<i>A. oryzae</i>	Urea	Crude protein increased from 1.98% to 13.3%, true protein increased from 0.98% to 12.4% and crude fibre decreased from 13.6% to 10.7%
Yafetto (2018)	<i>A. niger</i>	Ammonium nitrate	Crude protein increased by 22.61%

The use of fungal FCP in chicken diets

Fungal fermentation has been attributed to the improved nutritional qualities of cassava pulp. As a consequence, fermentation can increase the inclusion

levels of cassava pulp in chicken rations. As shown in Table 3, FCP may be included in broiler and laying hen rations greater than that of unfermented cassava pulp. However, the levels of FCP inclusion may vary from study

to studies depending on the nutritional qualities of FCP, dietary composition and other experimental conditions. In most cases, there is a positive correlation between nutrient digestibility and feed intake in chickens (Sundu et al., 2006). In light with this, the improved nutritional qualities in FCP (especially the reduced fibre content) may be attributed to the reduced bulkiness and increased digestibility and thereby increased feed intake in chickens (Khempaka et al., 2014). Fermentation has been suggested to improve the palatability of products (Supriyati et al., 2015). In this respect, better palatability of FCP may be one of the reasons for the increased FCP intake in chickens when compared with the intake of unfermented cassava pulp. The toxicology of liver due to urea

supplementation (during fermentation process) and the presence of toxic compounds such as HCN in cassava pulp may be a crucial point of consideration when using FCP as a dietary ingredient for chickens. In conjunction with Khempaka et al. (2014), we assessed the activities of aspartate aminotransferase (AST) and alanine aminotransferase (ALT) as indicators of liver health and found no change in the activities of these enzymes when feeding FCP as compared to control (Sugiharto et al., 2017b). The bioconversion of urea to fungal biomass protein (Bayitse et al., 2015) and the destruction of HCN during the fermentation process (Diarra and Devi, 2010) may implicate in safe inclusion of FCP in chicken diets.

Table 3. The levels of FCP inclusion in chicken rations

References	Recommended levels of FCP inclusion
Lubis et al. (2007)	<i>A. niger</i> -FCP-urea-zeolite may be used in diets up to 15% without negative impacts on growth performance and health of broilers.
Ali-Mursyid et al. (2010)	FCP could be included in diets up to 16.5% without detrimental effects on growth performance and nutrient digestibility of broiler chicks.
Khempaka et al. (2014)	<i>A. oryzae</i> -FCP can be included up to 16% in the rations with no deleterious effects on nutrient digestibility and retention, final body weight, carcass traits and biochemical parameters.
Sugiharto et al. (2017a)	<i>A. charticola</i> -FCP can be included up to 16% with no adverse effects on final body weight, digestibility and carcass characteristics of broiler chickens
Okmathok et al. (2018)	<i>A. oryzae</i> -FCP was safe to be included up to 24% in diets without deleterious effects on nutrient digestibility, egg production and quality and physiological conditions of laying hens

Functional properties of FCP as a dietary component in chicken diets

Fermented products have been attributed to functional properties such as higher lactic acid bacteria (LAB) and organic acids contents. These properties make FCP beneficial for the health of the gastrointestinal tract of chickens (Sugiharto and Ranjitkar, 2019). Regarding to the effect of FCP on chicken health, the published data are still scarce. A recent study by our research group revealed that *A. charticola*-FCP decreased coliform bacteria count in the ileum and increased butyric and propionic acid concentrations in cecal contents of broiler chickens (Sugiharto et al., 2017a). The capacity of LAB and organic acids in controlling the proliferation of potentially pathogenic bacteria such as coliform may be responsible for the reduced population of such pathogenic bacteria in the intestine of chickens fed FCP. To increase the functionality of fermented products, fermentation using probiotic microorganisms (as starter inoculum) has been conducted (Sugiharto and Ranjitkar, 2019). In this respect, fermented products may not only have improved

nutritional qualities, but also contain higher numbers of probiotic microorganisms. Due to the limited data, the study on the functional effect of FCP on chickens needs to be extensively conducted.

CONCLUSION

Solid-state fermentation using fungi can be a simple method to improve the nutritional qualities of cassava pulp and thus increase the inclusion level of such cheap agro-industrial by-product in chicken rations. Further studies are needed to explore the functional benefit of FCP on chicken health.

DECLARATIONS

Acknowledgments

Author acknowledges the support by Diponegoro University, Semarang Central Java, Indonesia through non-tax state revenue.

Competing interest

I have no conflict of interest.

Consent to publish

I gave my consent prior to publication of this article.

REFERENCES

- Ali-Mursyid WM, Bachruddin Z, Zuprizal and Nur-Cahyanto M (2010). Corn substitution using fermented solid cassava-waste on broiler chicken. *Journal of the Indonesian Tropical Animal Agriculture*, 35 (1): 9-15. DOI: 10.14710/jitaa.35.1.9-15
- Animashahun RA, Omoikhoje SO, Edokpayi MI and Adesina BT (2013). Nutrient values of cassava residual pulp as affected by solid-state fermentation with *Penicillium* spp. *International Journal of Applied Agricultural and Apicultural Research*, 9 (1&2): 42-47.
- Asadollahzadeh M, Ghasemian A, Saraeian A, Resalati H and Taherzadeh MJ (2018). Production of fungal biomass protein by filamentous fungi cultivation on liquid waste streams from pulping process. *BioResources*, 13 (3): 5013-5031.
- Bayitse R, Hou X, Laryea G and Bjerre A-B (2015). Protein enrichment of cassava residue using *Trichoderma pseudokoningii* (ATCC 26801). *AMB Express*, 5 (80). DOI: 10.1186/s13568-015-0166-8
- Bhardwaj V, Degrassi G and Bhardwaj RK (2017). Bioconversion of cellulosic materials by the action of microbial cellulases. *International Research Journal of Engineering and Technology*, 4 (8): 494-503.
- Chauynarong N, Bhuiyan MM, Kanto U and Iji PA (2015). Variation in nutrient composition of cassava pulp and its effects on *in vitro* digestibility. *Asian Journal of Poultry Science*. 9 (4): 203-212. DOI: [10.3923/ajpsaj.2015.203.212](https://doi.org/10.3923/ajpsaj.2015.203.212)
- Chauynarong N, Iji PA and Kanto U (2010). The optimum level of cassava pulp in diets for layers. *Proceeding of 21st Australian Poultry Science Symposium*, pp: 136-139.
- Diarra SS and Devi A (2015). Feeding value of some cassava by-products meal for poultry: A review. *Pakistan Journal of Nutrition*, 14 (10): 735-741. DOI: 10.3923/pjn.2015.735.741
- Gowthaman MK, Krishna C and Moo-Young M (2001). Fungal solid state fermentation - an overview. *Applied Mycology and Biotechnology*, 1: 305-352. DOI: 10.1016/S1874-5334(01)80014-9
- Khempaka S, Hokking L and Molee W (2016). Potential of dried cassava pulp as an alternative energy source for laying hens. *Journal of Applied Poultry Research*, 25: 359-369. DOI: 10.3382/japr/pfw020
- Khempaka S, Maliwan P, Okrathok S and Molee W (2018). Digestibility, productive performance, and egg quality of laying hens as affected by dried cassava pulp replacement with corn and enzyme supplementation. *Tropical Animal Health and Production*, 50: 1239-1247. DOI: 10.1007/s11250-018-1550-6
- Khempaka S, Molee W and Guillaume M (2009). Dried cassava pulp as an alternative feedstuff for broilers: Effect on growth performance, carcass traits, digestive organs, and nutrient digestibility. *Journal of Applied Poultry Research*, 18: 487-493. DOI: 10.3382/japr.2008-00124
- King AJ and Zeidler G (2004). Tomato pomace may be a good source of vitamin E in broiler diets. *California Agriculture*, 58 (1): 59-62. DOI: 10.3733/ca.v058n01p59
- Kumsri P, Ruangpanit Y, Attamangkune S, Songsermn O and Amornthewaphet N (2009). Effect of cassava pulp level and feed form on performance in broiler. *Proceeding of the 47th Kasetsart University Conference: Animals, Bangkok, Thailand*, pp: 132-140.
- Lubis AD, Suhartono, Darmawan B, Ningrum H, Noormasari IY and Nakagoshi N (2007). Evaluation of fermented cassava (*Manihot esculenta* Crantz) pulp as a feed ingredient for broiler. *Tropics*, 17 (1): 73-80. DOI: 10.3759/tropics.17.73
- Morgan NK and Choct M (2016). Cassava: Nutrient composition and nutritive value in poultry diets. *Animal Nutrition*, 2: 253-261. DOI: 10.1016/j.aninu.2016.08.010
- Mustafa AM, Poulsen TG and Sheng K (2016). Fungal pretreatment of rice straw with *Pleurotus ostreatus* and *Trichoderma reesei* enhance methane production under solid-state anaerobic digestion. *Applied Energy*, 180: 661-671. DOI: 10.1016/j.apenergy.2016.07.135
- Okrathok S, Pasri P, Thongkratok R, Molee W and Khempaka S (2018). Effects of cassava pulp fermented with *Aspergillus oryzae* as a feed ingredient substitution in laying hen diets. *Journal of Applied Poultry Research*, 27: 188-197. DOI: 10.3382/japr/pfx057
- Sengxayalth P and Preston TR (2017). Fermentation of cassava (*Manihot esculenta* Crantz) pulp with yeast, urea and diammonium phosphate (DAP). *Livestock Research for Rural Development* 29 (9). <http://www.lrrd.org/lrrd29/9/pom29177.html>
- Sugiharto S and Ranjitkar S (2019). Recent advances in fermented feed towards improved broiler chicken performance, gastrointestinal tract microecology and immune responses: A review. *Animal Nutrition*. DOI: 10.1016/j.aninu.2018.11.001
- Sugiharto S, Yudiarti T, Isroli I, Widiastuti E and Putra FD (2017b). Intestinal microbial ecology and hematological parameters of broiler fed cassava waste pulp fermented with *Acremonium charticola*. *Veterinary World*, 10 (3): 324-330. DOI: [10.14202/vetworld.2017.324-330](https://doi.org/10.14202/vetworld.2017.324-330)
- Sugiharto S, Yudiarti T, Isroli I, Widiastuti E and Putra FD (2017a). Effects of feeding cassava pulp fermented with *Acremonium charticola* on growth performance, nutrient digestibility and meat quality of broiler chicks. *South African Journal of Animal Science*, 47 (2): 103-138. DOI: 10.4314/sajas.v47i2.4B.
- Sugiharto S, Yudiarti T and Isroli I (2015). Functional properties of filamentous fungi isolated from the Indonesian fermented dried cassava, with particular application on poultry. *Mycobiology*, 43 (4): 415-422. DOI: [10.5941/MYCO.2015.43.4.415](https://doi.org/10.5941/MYCO.2015.43.4.415)
- Sugiharto S, Yudiarti T, Isroli I (2016). Haematological and biochemical parameters of broilers fed cassava pulp fermented with filamentous fungi isolated from the Indonesian fermented dried cassava. *Livestock Research for Rural Development*, 28 (4). <http://www.lrrd.org/lrrd28/4/sugi28053.html>

- Sundu B, Kumar A and Dingle J (2006). Response of broiler chicks fed increasing levels of copra meal and enzymes. *International Journal of Poultry Science*, 5 (1): 13-18. DOI: 10.3923/ijps.2006.13.18
- Supriyati, Haryati T, Susanti T and Susana IWR (2015). Nutritional value of rice bran fermented by *Bacillus amyloliquefaciens* and humic substances and its utilization as a feed ingredient for broiler chickens. *Asian-Australasian Journal of Animal Sciences*, 28 (2): 231-238. DOI: 10.5713/ajas.14.0039
- Tang DF, Ru YJ, Song SY, Choct M and Iji PA (2012). The effect of cassava chips, pellets, pulp and maize-based diets on performance, digestion and metabolism of nutrients for broilers. *Journal of Animal and Veterinary Advances*, 11 (9): 1332-1337. DOI: 10.3923/javaa.2012.1332.1337
- Thongkratok R, Khempaka S and Moole W (2010). Protein enrichment of cassava pulp using microorganisms fermentation techniques for use as an alternative animal feedstuff. *Journal of Animal and Veterinary Advance*, 9 (22): 2859-2862. DOI: 10.3923/javaa.2010.2859.2862
- Triprugsachart S, Ruangpanit Y, Attamanegkune S, Songserm O and Rattanatabtimtong S (2007). Effect of cassava pulp level on laying hen performance and egg quality. *Proceeding of the 47th Kasetsart University Conference: Animals, Bangkok, Thailand*. pp: 165-173.
- Yafetto L (2018). Protein enrichment of cassava pulp by solid-state fermentation using *Aspergillus niger*. *Studies in Fungi*, 3(1): 7-18. DOI: 10.5943/sif/3/1/2