



Research Article

Morphology and performance specifications of *Blaptostethus pallescens* Poppius (Heteroptera: Anthocoridae) when reared on two alternate laboratory hosts

TRIPTI GUPTA^{1*}, CHANDISH R. BALLAL² and SHASHIKALA S. KADAM²

¹Pest Control (India) Pvt. Ltd., Bio-Control Research Laboratories (BCRL), Arakere Post, Bengaluru - 562163, Karnataka, India

²ICAR-National Bureau of Agricultural Insect Resources, H. A. Farm Post, Bellary Road, Bengaluru - 560024, Karnataka, India

*Corresponding author E-mail: tripti123gupta@gmail.com

ABSTRACT: The present research paper provides detailed morphology and morphometrics of an anthocorid predator, *Blaptostethus pallescens* Poppius. To identify the alternate laboratory host to be utilized for mass production, biological parameters of the anthocorid were evaluated by rearing on UV irradiated eggs of two alternate laboratory hosts, *Corcyra cephalonica* (Stainton) and *Sitotroga cerealella* (Motsch.). Five nymphal instars were recorded on both hosts. No significant differences were observed in incubation period, nymphal duration, total developmental period and sex ratio when reared on the two hosts. However, significant differences were observed in two biological parameters *i.e.*, adult longevity and fecundity. Longer life span and higher fecundity of *B. pallescens* was recorded when reared on *C. cephalonica* eggs in comparison to *S. cerealella* eggs, indicating the suitability of *C. cephalonica* eggs for mass rearing *B. pallescens* in the laboratory. *B. pallescens* could be reared continuously for 7 generations on *C. cephalonica* eggs.

KEY WORDS: *Blaptostethus pallescens*, biological parameters, *Corcyra cephalonica*, fecundity, generations, incubation, morphology, morphometrics, *Sitotroga cerealella*

(Article chronicle: Received: 16-02-2018; Revised: 12-11-2018; Accepted: 15-12-2018)

INTRODUCTION

Pest management through utilization of chemical pesticides in agriculture is facing several obstacles including, development of pesticide resistance in pest population and human health hazards caused due to pesticide residues in food items. At this juncture biological control could emerge as an alternative method in controlling pests in different crop ecosystems. Several anthocorids species were reported to provide control several pests (Ballal and Yamada, 2016). *Blaptostethus pallescens* Poppius, a general predator with wide host range has been reported on maize, rose, castor, grapes, bamboo (Jalali and Singh, 2002; Ballal *et al.*, 2003b; 2009; Gupta, 2009) in India. *Blaptostethus pallescens* has been identified as a potential bio-control agent for the management of eggs and larvae of lepidopteran pest, sucking pests like mites, thrips, mealybugs, aphids and stored insect pests (Ballal *et al.*, 2009; 2012; Gupta and Ballal, 2011; Kaur and Singh, 2011; Sobhy *et al.*, 2014; Lessando *et al.*, 2015). However, natural population of this predator is insufficient to maintain the target pest population below economic injury levels on several crops. Therefore, mass production and augmentative releases of natural enemies may help to maintain some of the target pests below economic injury levels.

Knowledge on host acceptance behaviour and predatory potential of bio agents are prerequisites to optimise their mass production. Detailed information on morphometrics, biology and suitable alternative laboratory host for mass production are necessary to improve the utilization of this predator in biological control.

The objective of the present work was to study the morphology, morphometrics and biology to standardize an inexpensive production protocol utilizing an alternate laboratory host. We also aim to evaluate the effect of continuous laboratory rearing for several generations on the suitable and superior alternate laboratory host eggs.

MATERIALS AND METHODS

The study was conducted at the Mass Production Laboratory, National Bureau of Agricultural Insect Resources, Bengaluru. Morphology and morphometrics of *B. pallescens* egg, nymph and adult stages were studied by using ocular and stage micrometers. The biology of *B. pallescens* on two alternate laboratory host eggs *Corcyra cephalonica* (Stainton) and *Sitotroga cerealella* (Olivier) were studied under laboratory conditions (26±2°C and 55-65% RH).

From the lab reared *B. pallescens* culture, five pairs of adults were released into each pearl pet plastic container (500 ml). The containers were provided with UV-irradiated *C. cephalonica* eggs as feeding, bean pieces (4-5 per container) as ovipositional substrates and cotton lint to avoid cannibalism. Eight such sets were maintained. After every 24 hour period, the bean pieces with *B. pallescens* eggs were collected and observed under the microscope to record the number of eggs laid, after which they were placed in 500 ml pearl pet containers for hatching.

Number of nymphs hatched from total eggs on the bean pieces in each container was counted for calculating percent hatching. Freshly hatched nymphs were kept individually in jewel boxes provided with UV-irradiated *C. cephalonica* eggs and observed daily to record total number of instars, duration of each instar and total nymphal period.

When adults emerged, they were collected and observed under microscope to differentiate the sex. Percent adults formed were calculated based on the number of healthy adults developed from the total number of nymphs recorded in each replication. Longevity of adult male and female was recorded. Same procedure was followed to study the biology of *B. pallescens* reared on *S. cerealella* eggs.

Generation-wise progeny production was studied by initiating the experiment with field collected adults of *B. pallescens*. The experiment was initiated with two females of *B. pallescens* collected from maize field per replication and replicated five times. The field collected adults were allowed to oviposit and the nymphs which hatched from the eggs were placed in a nymphal container. The progeny produced by the field-collected females in the laboratory was considered as the first laboratory generation. The adults belonging to the first laboratory generation were placed in an ovipositional container, which was provided with feeding (*C. cephalonica* eggs) and ovipositional substrate (bean pieces). This ovipositional container was marked as the first laboratory generation. Progeny produced by the adults of each generation were recorded and kept in separate ovipositional containers marked with the generation number. The progeny production per female and per cent increase per generation were also calculated and represented graphically.

RESULTS AND DISCUSSION

Morphology and morphometrics

Eggs are bottle shaped and creamy white with an operculum at the anterior end. The posterior end is oval and inserted inside the bean pod with only the operculum visible. Some eggs are laid tangential to the surface of bean pods. At

the time of hatching, operculum opens like a lid. The mode of egg laying varies in different anthocorid species. Some anthocorid species require plant tissue for the oviposition, while some can lay on cotton strands, container walls, etc. Muraleedharan and Ananthakrishnan, (1978) reported that the eggs of *Carayonocoris indicus* Muraleedharan are inserted into the petiole of *Cassia marginata* L. *Blaptostethus pallescens* eggs are deposited singly with the operculum facing upward. Beans pods can be used successfully as ovipositional substrate for rearing *B. pallescens* (Ballal *et al.*, 2003b; Sobhy *et al.*, 2014). Kodakkadan Srikumar *et al.* (2017) reported that *B. pallescens* laid eggs on tea shoots. The average length, breadth of egg and diameter of operculum of *B. pallescens* egg are 0.82 ± 0.02 mm, 0.38 ± 0.00 mm and 0.25 ± 0.25 mm, respectively (Table 1).

Five nymphal instars were reported in present study. Morphological characteristics of each instar were as follows

- First instar: Just after hatching, the head, thorax and posterior margin of abdomen is slightly pink in colour with dark red eyes. The rostrum is three segmented and pale yellow in colour. The antenna is four segmented and greyish in colour. Dark red scent glands are clearly visible on the 3rd, 4th and 5th abdominal segments. Abdominal apex has long white bristles and legs are pale yellow in colour. The average length and greatest width 0.93 ± 0.02 and 0.38 ± 0.00 mm, respectively.
- Second instar: Uniformly pink in colour. First antennal segment is transparent and the other three segments are greyish in colour. First and second antennal segments are stout and 3rd and 4th segments are slender and filiform. It measures 1.07 ± 0.02 mm in length and 0.45 ± 0.02 mm in width (greatest).

Table 1. Morphometrics of *Blaptostethus pallescens*

Stage	Length (mm)	Greatest width (mm)
Egg	0.82 ± 0.02	0.38 ± 0.0
Nymphal instars		
I	0.93 ± 0.02	0.38 ± 0.00
II	1.07 ± 0.02	0.45 ± 0.02
III	1.66 ± 0.06	0.69 ± 0.04
IV	1.97 ± 0.11	0.88 ± 0.02
V	2.60 ± 0.14	0.90 ± 0.02
Adult		
Male	2.70 ± 0.06	0.92 ± 0.08
Female	2.90 ± 0.24	0.98 ± 0.03




Fig. 1. Day wise fecundity of *Blaptostethus pallescens* on two alternate laboratory host eggs.

- c) Third instar: Acquires uniform reddish orange colour and is darker in colour than second instar. Wing pads are visible. Average length is 1.66 ± 0.06 mm and greatest width 0.69 ± 0.04 mm.
- d) Fourth instar: Body colour changes to dark reddish brown. Scent glands are not very distinct as in earlier instars. Wing pads are more developed. Length and greatest width measured 1.97 ± 0.11 and 0.88 ± 0.02 mm, respectively.
- e) Fifth instar: Reddish black in colour. Rostrum red. Wing pads extend beyond thorax. Basal antennal segment is darker than rest of the segments. Average length 2.6 ± 0.14 mm and greatest width 1.0 ± 0.02 mm.

Tawfik and Husseini (1971), Ballal *et al.* (2003b), Sobhy *et al.* (2014) and Kodakkadan Srikumar *et al.* (2017) also reported five nymphal instars in *B. pallescens*. Muraleedharan and Ananthakrishnan (1978) reported five nymphal instars in anthocorid species like *C. indicus*, *Montandoniolamoraguesi* (Puton), *Xylocoris clarus* (Distant) and *Scoloposcelis paralellus* (Motschulsky).

Fifth instar moults to become adult which is black in colour with fully developed wings. Clear sexual dimorphism is visible in *B. pallescens*. Abdomen is broader in female, with two copulatory tubes. The tip of abdomen in male is slightly twisted towards its left side when observed ventrally. A bunch of setae are visible on dorsal

side in both sexes. Fore femora is with 5-6 strong peg like teeth on distal side and fore tibiae with a lateral row of teeth almost throughout their length. Mid and hind tibiae are with irregular spines on the surface and more spines are present in the distal half. Female is larger than male. Similar observations were made by Rajasekhara (1973) in *B. kumbi* and Ballal *et al.* (2003b) in *B. pallescens*. Average length and greatest width of female is 2.9 ± 0.24 and 0.98 ± 0.03 mm, respectively and that of male 2.7 ± 0.06 mm and 0.92 ± 0.08 mm, respectively (Table 1).

Comparative Biology of *Blaptostethus pallescens* on *Corcyra cephalonica* and *Sitotroga cerealella*

There was no significant difference in the incubation period of *B. pallescens*, when reared on *C. cephalonica* and *S. cerealella* host eggs. It ranged from 3-5 (mean 4.5 ± 0.22) and 4-6 days (mean 4.60 ± 0.33), respectively (Table 2). There was no significant difference in the total nymphal period when fed on *S. cerealella* and *C. cephalonica*, mean values being 16.50 ± 1.02 and 16.3 ± 0.62 days, respectively (Table 2). Duration of first to fifth nymphal instars was observed to be 3.6 ± 0.62 , 2.2 ± 0.20 , 2.2 ± 0.37 , 2.4 ± 0.40 and 6.0 ± 0.55 days, respectively (Table 3). First and fifth nymphal instars were recorded to be a longer nymphal duration which has also been reported by Sobhy *et al.* (2014) and Kodakkadan Srikumar *et al.*, (2017).

No significant difference was observed in total developmental time of female and male when reared on *C. cephalonica* and *S. cerealella* eggs (Table 2). Female survived for a longer duration than male in both the treatments.

Table 2. Comparative biology of *Blaptostethus pallescens* on eggs of *Sitotroga cerealella* and *Corcyra cephalonica*

Stage	<i>S. cerealella</i>	<i>C. cephalonica</i>	CD at P ≤ 0.05
	Mean ± SEM (Range)	Mean ± SEM (Range)	
Egg period (days)	4.60±0.33 (4-6)	4.5± 0.22 (3-5)	NS
Total nymphal Period (days)	16.50±1.02 (14-20)	16.3±0.62(14-16)	NS
Male Developmental period	23.00±1.52 (20-25)	21.0± 0.50(21-20)	NS
Female Developmental period	19.60±0.03(19-20)	19.8± 2.30(15-23)	NS
Longevity male (days)	31.25±2.05 (26-32)	42.4± 5.82 (38-68)	4.05
Longevity female (days)	47.40±4.55 (36-58)	58.2± 5. 06 (44-78)	4.77
Eggs/female	91.25±8.43 (74-109)	132.8± 22.4 (84-199)	30.09
Nymphs/female	75.75±8.86 (55-91)	119.6± 18.3 (78-175)	8.67
Adults/female	58.50±8.70 (40-82)	100.4±17.46 (67-160)	6.92
Percent eggs hatched	82.59±3.01 (75-89)	90.9± 2.18 (84.61-97.5)	2.65
Percent adults developed from total number of nymphs	77.05± 5.79(63-90)	83.46±2.41(77.31-90.28)	NS
Sex ratio (F: M) (% female)	1.44: 1 (59)	1.5: 1 (60)	NS

Table 3. Duration of nymphal instars of *Blaptostethus pallescens* on *Corcyra cephalonica* eggs (the more suitable host)

Instar	Duration in days	
	Mean ± SEM	Range
I	3.6± 0.62	2-6
II	2.2± 0.20	2-3
III	2.2± 0.37	2-3
IV	2.4± 0.40	2-4
V	6.0± 0.55	4-7

Longevity of adult female and male was significantly higher when reared on *C. cephalonica* eggs in comparison to when reared on *S. cerealella* eggs. On *C. cephalonica* eggs, female longevity ranged from 44-78 days (mean 58.2±5.06) and male longevity was 38-68 days (mean 42.4±5.82) (Table 2). When *B. pallescens* was provided with *S. cerealella* eggs, females lived for a mean of 47.40±4.55 (range 36-58) and males survived for 31.25±2.05 (range 26-32) days. Longer life span of females in other anthocorid species like *Cardiastethus exiguus* Poppius and *Orius tantillus* (Motsch.) were also observed by Ballal *et al.* (2003a) and Gupta and Ballal (2006).

Adults mated just after emergence. Significant difference was observed in the number of eggs laid by a female when reared on *S. cerealella* eggs in comparison to that when reared on *C. cephalonica* eggs. Fecundity varied from 74-109 eggs per female (mean 91.25±8.43), when reared on eggs of *S. cerealella* and 84-199 eggs per female (mean 132.8±22.4),

when reared on eggs of *C. cephalonica*. Egg laying was observed from the first day after mating and continued till mortality of the adult female. The number of eggs laid per day varied from 0 to 6. Higher peaks were observed on 5th, 12th, 16th and 30th day (approximately 6 eggs per day) when fed on *C. cephalonica* eggs (Fig. 1). Intermittently, zero progeny production was recorded. After 50 days, progeny production gradually reduced. When *S. cerealella* eggs were provided, higher peaks were observed on 2nd, 15th, 19th, 20th and 43rd day.

Per cent hatching was 90.9 and 83.46% of nymphs formed adults when reared on *C. cephalonica* eggs. On *S. cerealella* eggs 82.59% eggs hatched and 77.05% nymphs developed into adults. Sex ratio was 1.44: 1(Female: Male) on *S. cerealella* and on *C. cephalonica* 1.5:1.00 (Female: Male). A female biased sex ratio was recorded in *B. pallescens* when reared on both host eggs. This agrees with the studies of Tawfik and Husseini (1971) and Ballal *et al.* (2003b) who also observed percent female progeny was more than male. Generally a female biased sex ratio is considered to be positive attribute for a biological control agent, making it amenable to rearing in the insectary and a good field performer.

Generation-wise progeny production of *Blaptostethus pallescens* on UV-irradiated *Corcyra cephalonica* eggs

When *B. pallescens* was reared on *C. cephalonica* eggs in the laboratory, seven generations could be reared in a period of nine months. Nine adults were produced in the first generation from one field collected *B. pallescens* female. Progeny production in the second generation was recorded

as 13 adults per female. In the third generation there was an increase in total progeny production i.e., 126 adults but the percentage increase was only 62. In the fourth, fifth and sixth generations, the progeny production increased by 67, 90 and 127 percent respectively, with the number of progeny produced recorded as 211, 400 and 669 adults in the respective generations. In the seventh generation a total of 932 adults were produced but the percent increase was only 39 (Fig. 2). Almost six generations overlapped during the study period because of the prolonged longevity of the adult females. Progeny production was high in the first laboratory generation compared to the subsequent laboratory generations. Bio-deterioration in the laboratory cultures of parasitoids and predators was also observed by Penn *et al.* (1998), PDBC- ICAR (1999), Ballal *et al.* (2001; 2003a) on continuous laboratory rearing of *Trichogramma* parasitoids, *Scymnus coccivora* Ayyar, *Campoletis chlorideae* Uchida and *Cardiastethus exiguus* Poppius, respectively. Though the total number of adults obtained per generation kept increasing, the number of progeny produced per female ranged between 2.48- 4.46 from second to seventh laboratory generations (Fig. 2).

The remarkable increase in the total *B. pallescens* adults obtained from each production unit in each generation

could be attributed to the high fecundity and longevity of *B. pallescens* on *C. cephalonica* eggs. However, the lower progeny production per female could be due to the rearing conditions provided and continuous rearing in confinement. This agrees with the statement of Boller (1972) that the role of insect behavior and its effects under selection in the insectaries is a major factor influencing the quality of reared insect.

Rearing of anthocorid predators on target hosts like thrips or mites may be extremely cumbersome. In the present study the biological parameters of *B. pallescens*, with high longevity and fecundity indicates that it is a potential bio agent which can be effectively reared on eggs of alternate laboratory host *C. cephalonica* eggs. It was also observed that field collected *B. pallescens* could be reared successfully up to seven generations in the laboratory, rejuvenation of the laboratory culture may be required after seventh generations by bringing in the wild culture of the anthocorid predator from field and mixing with the lab culture.

ACKNOWLEDGEMENTS

The authors wish to thank Dr. Sunil Joshi, Principal Scientist and Dr. Y. Lalitha, Assistant Chief Technical Officer,




Fig. 2. Generation wise progeny production of *Blaptostethus pallescens* on *Corcyra cephalonica* eggs (figures given in brackets are rate of increase/ decrease from the previous generation)

ICAR-National Bureau of Agricultural Insect Resources, for the continuous supply of eggs of *Sitotroga cerealella* and *Corcyra cephalonica* for conducting the experiments.

REFERENCES

- Ballal CR, Joshi S, Rao NS. 2001. Bio-deterioration of laboratory cultures of *Campoletis chlorideae* Uchida. pp. 257. In: Verghese A, Reddy RP (Eds.) *IPM in Horticultural crops: Emerging trends in the New Millennium*. Proceeding of the Second National Symposium on Integrated pest Management (IPM) in Horticultural Crops New molecules, Bio-pesticides and Environment. Bangalore, 257 pp. PMCid:PMC95474
- Ballal CR, Singh SP, Poorani J, Tripti Gupta. 2003a. Feasibility of mass multiplication and utilization of *Cardiastethus exiguus* Poppius, a potential anthocorid predator of *Opisina arenosella* Walker (Lepidoptera: Oecophoridae). pp. 29-33. In: Tandon PL, Ballal CR., Jalali S. K. (Eds.). *Biological Control of Lepidopteran pests*, 354 pp. PMID:15025331
- Ballal CR, Singh SP, Poorani J, Gupta T. 2003b. Biology and rearing requirements of an anthocorid predator, *Blaptostethus pallescens* Poppius (Heteroptera: Anthocoridae). *J Biol Control* **17**: 29-33.
- Ballal CR, Gupta T, Sunil Joshi, Chandrashekhar K. 2009. Evaluation of an anthocorid predator, *Blaptostethus pallescens* against two-spotted spider mite, *Tetranychus urticae*. *Bull IOBC/WPRS* **49**: 127-132.
- Ballal CR, Gupta T, Joshi S. 2012. Morphometry and biology of a new anthocorid *Montandoniola indica*, a potential predator of *Gynaikothrips uzeli*. *Bull IOBC/WPRS* **80**: 79-84 pp.
- Ballal CR, Yamada K. 2016. Anthocorid predator. In: pp. 329-366. Omkar (Ed.). *Ecofriendly pest management for food security* Elsevier, London 727pp.
- Boller E. 1972. Behavioural aspects of mass-rearing of insects. *Entomophaga* **17**(1): 9-25. <https://doi.org/10.1007/BF02371070>
- Kodakkadan Srikumar, Smitha S. Suresh kumar B, Radakrishnan B. 2017. Biology and feeding efficacy of the anthocorid, *Blaptostethus pallescens* Poppius on *Oligonychus coffeae* in tea. *J Biol Control* **31**(4): 198-200. <https://doi.org/10.18311/jbc/2017/18157>
- Gupta T, Ballal CR. 2006. Biology and Feeding Potential of an Anthocorid Predator, *Orius tantillus* (Motsch.) (Heteroptera: Anthocoridae) on *Sitotroga cerealella*. *Ind J Plant Protect.* **34**(2): 168-172.
- Gupta T. 2009. *Studies on production and utility of anthocorid predators- with special reference to Orius spp. and Blaptostethus spp.* Ph.D thesis submitted to Mysore University, 207.
- Gupta T, Ballal CR. 2011. Preferential feeding potential of an anthocorid predator *Blaptostethus pallescens* Poppius on different stages of cotton mealybug. *J Environ Entomol.* **33**(4): 423-428.
- Jalali SK, Singh SP. 2002. Seasonal activity of stem borers and their natural enemies on fodder maize. *Entomon* **27**(2): 137-146.
- Kaur R, Singh VJ. 2011. *Blaptostethus pallescens* Poppius and *Xylocoris flavipes* (Reuter) in the suppression of *Corcyra cephalonica* Stainton in stored rice grain. *J Biol Control* **25**(4): 329-332.
- Lessando M. Gontijo, Daiane Celestino, Obiratanea S. Queiroz, Raul Narciso C. Guedes, Marcelo C. Picanço. 2015. Impacts of azadirachtin and chlorantraniliprole on the developmental stages of pirate bug predators (Hemiptera: Anthocoridae) of the tomato pinworm *Tuta absoluta* (Lepidoptera: Gelechiidae). *Florida Entomologist* **98**(1). <https://doi.org/10.1653/024.098.0111>
- Muraleedharan N, Ananthakrishnan TN. 1978. Bioecology of four species of Anthocoridae (Hemiptera: Insecta) predaceous on thrips, with key to genera of anthocorid from India. Occasional paper. *Records Zool Surv India* **11**: 1-32.
- Penn SL, Ridgway RL, Scriven GT, Inscoc MN. 1998. Quality assurance by the commercial producers of arthropod natural enemies. pp. 202-230. In: Ridgway RL, Hoffmann MP, Inscoc MN, Glenistar CS (Eds.). In: *Mass reared natural enemies: Application, Regulation and Needs*. Entomological Society of America, Baltimore, PDBC- ICAR. 1999. Annual Report of the Project Directorate of Biological Control, Bangalore, 1998-99, 217 pp.
- Sobhy I, Hamid AMA, Sarhan AA, Shoukry AA, Mandour NS, Reitz SR. 2014. Life history traits of *Blaptostethus*

pallescens (Hemiptera: Anthocoridae), a candidate for use in augmentative biological control in Egypt. *Appl Entomol Zool.* **49**: 315-324. <https://doi.org/10.1007/s13355-014-0252-4>

Tawfik MFS, El-Husseini MM. 1971. The life history of the anthocorid predator *Blaptostethus piceus* Fieber var. *pallescens* Poppius (Hemiptera: Anthocoridae). *Bulletin de la Societe Entomologique d'Egypte* **55**: 239-252.