

# A transgenic monkey model for the study of human brain evolution

Lei Shi<sup>1,2</sup>, Bing Su<sup>1,2,3,\*</sup>

<sup>1</sup>State Key Laboratory of Genetic Resources and Evolution, Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming Yunnan 650223, China

<sup>2</sup>Primate Research Center, Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming Yunnan 650223, China

<sup>3</sup>Center for Excellence in Animal Evolution and Genetics, Chinese Academy of Sciences, Kunming Yunnan 650223, China

Why humans have large brains with higher cognitive abilities is a question long asked by scientists. However, much remains unknown, especially the underlying genetic mechanisms. With the use of a transgenic monkey model, we showed that human-specific sequence changes of a key brain development gene (Primary microcephaly 1, *MCPH1*) could result in detectable molecular and cognitive changes resembling human neoteny, a notable characteristic developed during human evolution. This study was published in *National Science Review* (Shi et al., 2019).

Neoteny has been hypothesized to help explain why we differ from our close relatives, such as the chimpanzee (*Pan troglodytes*) (Liu et al., 2012). Also called juvenilization, neoteny is the delay in or slowing down of physiological development of a species (Skulachev et al., 2015). Neoteny in humans is the retention of juvenile features into adulthood and is exaggerated compared to that in non-human primates. The neotenuous human brain provides an extended window for the plasticity of the neural network, and therefore a longer time for learning, which may be crucial for the formation of human cognition (Bucifer et al., 2011).

In the past two decades, comparative genomic studies have identified many candidate genes that carry human-specific sequence changes potentially contributing to human brain evolution (Bustamante et al., 2005; Clark et al., 2003). However, functional dissection of these candidate genes, particularly the phenotypic consequences of human-specific mutations, is lacking. As a model organism, transgenic mice (*Mus musculus*) have been used in the study of human brain function; for example, the study of *Foxp2* that carries two human-specific amino acid changes (Enard et al., 2009). However, compared to the human brain, the mouse brain is dramatically different in size, structure, developmental pattern, and function, and therefore is not an ideal model to study human brain evolution. In contrast, the rhesus monkey (*Macaca mulatta*), an Old-World primate species widely used in biomedical research, is a better animal model due to its high sequence similarity with humans (>93% for protein coding genes) (Yan et al., 2011;

Zhang et al., 2014), yet relatively large phylogenetic distance (~25 million years of divergence from humans), which alleviates certain ethical concerns (Coors et al., 2010). These advantages have been demonstrated in the use of transgenic monkey models of human diseases (Chen et al., 2017; Liu et al., 2016; Luo et al., 2016; Qiu et al., 2019; Yang et al., 2008; Zhang et al., 2018).

*MCPH1* is a key gene regulating brain development in humans, and its truncate mutations can cause primary microcephaly (Jackson et al., 2002). Similarly, *MCPH1* knockout in mice and monkeys can recapitulate the microcephaly phenotype (Gruber et al., 2011; Ke et al., 2016), highlighting its critical role in mammalian brain development. In previous research, we demonstrated accelerated protein sequence changes in *MCPH1* during primate evolution, especially in the lineage leading to the origin of our own species (Wang & Su, 2004). Furthermore, *in vitro* functional analysis showed that human-specific *MCPH1* changes resulted in functional divergence between human and non-human primates (Shi et al., 2013). Such lines of evidence raise the question as to what phenotypic changes would appear following the introduction of human *MCPH1* copies into the rhesus monkey genome.

Using lentivirus delivery, we successfully generated 11 transgenic monkeys carrying 2–9 copies of human *MCPH1*. Brain image analysis showed that the transgenic monkeys had similar brain volume and cortex thickness as the controls, but their relative brain volume and gray matter percentage were higher. Importantly, the transgenic monkeys exhibited delayed cortex gray matter development compared to the

Received: 04 April 2019; Accepted: 09 April 2019; Online: 11 April 2019

Foundation items: This study was supported by the Strategic Priority Research Program of the Chinese Academy of Sciences (XDB13000000), the National Natural Science Foundation of China (31730088) and the Youth Innovation Promotion Association of CAS

\*Corresponding author, E-mail: sub@mail.kiz.ac.cn

DOI: 10.24272/j.issn.2095-8137.2019.031

controls. Furthermore, the transgenic monkeys had more immature and fewer mature neurons and glial cells than the controls, which, according to the transcriptome data, was likely caused by the suppression of many neuron maturation- and neuron differentiation-related genes. This observation was further illustrated by detailed analysis of developing brain laminae at the neurogenesis peak during fetal development.

Finally, to test whether the observed delay in brain development had any impact on behavior and cognition, we tested working memory in the transgenic monkeys using delayed matching-to-sample tasks. Notably, the transgenic monkeys exhibited better working memory and shorter reaction time than the controls, suggesting that neoteny brain development in the transgenic monkeys was beneficial for the formation of cognitive abilities, confirming the proposed evolutionary advantage of human neoteny. In our future work, we will: (1) examine the molecular pathway explaining how human *MCPH1* copies delay brain development and neural maturation, thus identifying more neoteny-related genes; and (2) apply more sophisticated tests to further understand how the neoteny brain affects cognitive formation in transgenic monkeys.

This study represents the first attempt to utilize a transgenic monkey model to study human brain evolution. It also provides the first molecular genetic evidence showing neoteny changes during brain development caused by human-specific mutations of a single gene. The results highlight the great potential of non-human primate models in studying human evolution and may pave the way for future studies to explore the genetic mechanisms of human-specific traits and elucidate the etiology of human brain disorders such as autism and Alzheimer's disease.

## COMPETING INTERESTS

The authors declare that they have no competing interests.

## AUTHORS' CONTRIBUTIONS

L.S. and B.S. wrote the manuscript. All authors read and approved the final version of the manuscript.

## REFERENCES

Bufile E, Agustí J, Blesa R. 2011. Human neoteny revisited: The case of synaptic plasticity. *American Journal of Human Biology*, **23**(6): 729–739.

Bustamante CD, Fedel-Alon A, Williamson S, Nielsen R, Hubisz MT, Glanowski S, Tanenbaum DM, White TJ, Sninsky JJ, Hernandez RD, Civallo D, Adams MD, Cargill M, Clark AG. 2005. Natural selection on protein-coding genes in the human genome. *Nature*, **437**(7062): 1153–1157.

Chen Y, Yu J, Niu Y, Qin D, Liu H, Li G, Hu Y, Wang J, Lu Y, Kang Y, Jiang Y, Wu K, Li S, Wei J, He J, Wang J, Liu X, Luo Y, Si C, Bai R, Zhang K, Liu J, Huang S, Chen Z, Wang S, Chen X, Bao X, Zhang Q, Li F, Geng R, Liang A, Shen D, Jiang T, Hu X, Ma Y, Ji W, Sun Y. E. 2017. Modeling rett syndrome using TALEN-Edited MECP2 mutant cynomolgus monkeys. *Cell*, **169**(5): 945–955.

Clark AG, Glanowski S, Nielsen R, Thomas PD, Kejariwal A, Todd MA, Tanenbaum DM, Civallo D, Lu F, Murphy B, Ferreira S, Wang G, Zheng X, White TJ, Sninsky JJ, Adams MD, Cargill M. 2003. Inferring nonneutral evolution from human-chimp-mouse orthologous gene trios. *Science*, **302**(5652): 1960–1963.

Coors ME, Glover JJ, Juengst ET, Sikela JM. 2010. The ethics of using transgenic non-human primates to study what makes us human. *Nature Reviews Genetics*, **11**(9): 658–662.

Enard W, Gehre S, Hammerschmidt K, Hölter SM, Blass T, Somel M, Brückner MK, Schreiweis C, Winter C, Sohr R, Becker L, Wiebe V, Nickel B, Giger T, Müller U, Groszer M, Adler T, Aguilar A, Bolle I, Calzada-Wack J, Dalke C, Ehrhardt N, Favor J, Fuchs H, Gailus-Dumer V, Hans W, Hölzlwimmer G, Javaheri A, Kalaydjiev S, Kallnik M, Kling E, Kunder S, Mossbrugger I, Naton B, Racz I, Rathkolb B, Rozman J, Schrewe A, Busch DH, Graw J, Ivandic B, Klingenspor M, Klopstock T, Ollert M, Quintanilla-Martinez L, Schulz H, Wolf E, Wurst W, Zimmer A, Fisher SE, Morgenstern R, Arendt T, de Angelis MH, Fischer J, Schwarz J, Pääbo S. 2009. A humanized version of Foxp2 affects cortico-basal ganglia circuits in mice. *Cell*, **137**(5): 961–971.

Gruber R, Zhou Z, Sukchev M, Joerss T, Frappart PO, Wang ZQ. 2011. MCPH1 regulates the neuroprogenitor division mode by coupling the centrosomal cycle with mitotic entry through the Chk1-Cdc25 pathway. *Nature Cell Biology*, **13**(11): 1325–1334.

Jackson AP, Eastwood H, Bell SM, Adu J, Toomes C, Carr IM, Roberts E, Hampshire DJ, Crow YJ, Mighell AJ, Karbani G, Jafri H, Rashid Y, Mueller RF, Markham AF, Woods CG. 2002. Identification of microcephalin, a protein implicated in determining the size of the human brain. *American Journal of Human Genetics*, **71**(1): 136–142.

Ke Q, Li W, Lai X, Chen H, Huang L, Kang Z, Li K, Ren J, Lin X, Zheng H, Huang W, Ma Y, Xu D, Chen Z, Song X, Lin X, Zhuang M, Wang T, Zhuang F, Xi J, Mao FF, Xia H, Lahn BT, Zhou Q, Yang S, Xiang AP. 2016. TALEN-based generation of a cynomolgus monkey disease model for human microcephaly. *Cell Research*, **26**(9): 1048–1061.

Liu X, Somel M, Tang L, Yan Z, Jiang X, Guo S, Yuan Y, He L, Oleksiak A, Zhang Y, Li N, Hu Y, Chen W, Qiu Z, Paabo S, Khaitovich P. 2012. Extension of cortical synaptic development distinguishes humans from chimpanzees and macaques. *Genome Research*, **22**(4): 611–622.

Liu Z, Li X, Zhang JT, Cai YJ, Cheng TJ, Cheng C, Wang Y, Zhang CC, Nie YH, Chen ZF, Bian WJ, Zhang L, Xiao J, Lu B, Zhang YF, Zhang XD, Sang X, Wu JJ, Xu X, Xiong ZQ, Zhang F, Yu X, Gong N, Zhou WH, Sun Q, Qiu Z. 2016. Autism-like behaviours and germline transmission in transgenic monkeys overexpressing MeCP2. *Nature*, **530**(7588): 98–102.

Luo X, Li M, Su B. 2016. Application of the genome editing tool CRISPR/Cas9 in non-human primates. *Zoological Research*, **37**(4): 214–219.

Qiu P, Jiang J, Liu Z, Cai Y, Huang T, Wang Y, Liu Q, Nie Y, Liu F, Cheng J, Li Q, Tang YC, Poo MM, Sun Q, Chang HC. 2019. BMAL1 knockout macaque monkeys display reduced sleep and psychiatric disorders. *National Science Review*, **6**(1): 87–100.

Shi L, Li M, Lin Q, Qi X, Su B. 2013. Functional divergence of the brain-size regulating gene MCPH1 during primate evolution and the origin of humans. *BMC Biology*, **11**: 62.

Shi L, Luo X, Jiang J, Chen Y, Liu C, Hu T, Li M, Lin Q, Li Y, Huang J, Wang H, Niu Y, Shi Y, Styner M, Wang J, Lu Y, Sun X, Yu H, Ji W, Su B. 2019. Transgenic rhesus monkeys carrying the human MCPH1 gene copies show

- human-like neoteny of brain development. *National Science Review*, doi: 10.1093/nsr/nwz043.
- Skulachev VP, Holtze S, Vyssokikh MY, Bakeeva LE, Skulachev MV, Markov AV, Hildebrandt TB, Sadovnichii VA. 2017. Neoteny, prolongation of youth: from naked mole rats to “naked apes” (humans). *Physiological Reviews*, **97**(2):699–720.
- Wang YQ, Su B. 2004. Molecular evolution of microcephalin, a gene determining human brain size. *Human Molecular Genetics*, **13**(11): 1131–1137.
- Yan G, Zhang G, Fang X, Zhang Y, Li C, Ling F, Cooper DN, Li Q, Li Y, van Gool AJ, Du H, Chen J, Chen R, Zhang P, Huang Z, Thompson JR, Meng Y, Bai Y, Wang J, Zhuo M, Wang T, Huang Y, Wei L, Li J, Wang Z, Hu H, Yang P, Le L, Stenson PD, Li B, Liu X, Ball EV, An N, Huang Q, Zhang Y, Fan W, Zhang X, Li Y, Wang W, Katze MG, Su B, Nielsen R, Yang H, Wang J, Wang X, Wang J. 2011. Genome sequencing and comparison of two nonhuman primate animal models, the cynomolgus and Chinese rhesus macaques. *Nature Biotechnology*, **29**(11): 1019–1023.
- Yang SH, Cheng PH, Banta H, Piotrowska-Nitsche K, Yang JJ, Cheng EC, Snyder B, Larkin K, Liu J, Orkin J, Fang ZH, Smith Y, Bachevalier J, Zola SM, Li SH, Li XJ, Chan AWS. 2008. Towards a transgenic model of Huntington's disease in a non-human primate. *Nature*, **453**(7197): 921–924.
- Zhang W, Wan H, Feng G, Qu J, Wang J, Jing Y, Ren R, Liu Z, Zhang L, Chen Z, Wang S, Zhao Y, Wang Z, Yuan Y, Zhou Q, Li W, Liu GH, Hu B. 2018. SIRT6 deficiency results in developmental retardation in cynomolgus monkeys. *Nature*, **560**(7720): 661–665.
- Zhang XL, Pang W, Hu XT, Li JL, Yao YG, Zheng YT. 2014. Experimental primates and non-human primate (NHP) models of human diseases in China: current status and progress. *Zoological Research*, **35**(6): 447–464.