

Meta-Analysis

Asian Pacific Journal of Reproduction

Journal homepage: www.apjr.net



doi: 10.4103/2305-0500.270102

Effect of cognitive behavioral therapy on anxiety and depression of infertile women: A meta-analysis

Hossein Mashhadi Abdolahi¹, Morteza Ghojzadeh², Salman Abdi³, Mahdieh Abbasalizad Farhangi⁴, Zeinab Nikniaz⁵, Leila Nikniaz¹✉¹Tabriz Health Services Management Research Center, Tabriz University of Medical Sciences, Tabriz, Iran²Research Center for Evidence-Based Medicine, Tabriz University of Medical Sciences, Tabriz, Iran³Clinical Psychiatry Research Center, Tabriz University of Medical Sciences, Tabriz, Iran⁴Drug Applied Research Center, Tabriz University of Medical Sciences, Tabriz, Iran⁵Liver and Gastrointestinal Diseases Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

ARTICLE INFO

Article history:

Received 7 April 2019

Revision 20 June 2019

Accepted 8 September 2019

Available online 7 November 2019

Keywords:

Cognitive behavioral therapy

Anxiety

Depression

Infertile women

Meta-analysis

ABSTRACT

Objective: To investigate the effect of cognitive behavioral therapy on anxiety and depression in infertile women.**Methods:** This study was performed in 2019 by searching Google Scholar, Scopus, PubMed/MEDLINE, Cochrane library, EMBASE, Proquest, ISI Web of Science, MagIran, SID and IranMedex. Eligible studies were selected by two reviewers and outcomes of interest were extracted. The meta-analysis was performed using the random effect models. I² - square statistic test was used for heterogeneity analysis. Presence of publication bias was also checked.**Results:** Eleven studies were included in this review. Between-group differences (cognitive behavioral therapy and control groups) in risk ratio for infertile women's depression was $d = -1.36$; 95% $CI = -1.81, -0.90$; $P < 0.001$. For infertile women's anxiety, between-group differences in risk ratio was $d = -0.83$; 95% $CI = -1.18, -0.47$; $P < 0.001$.**Conclusions:** Cognitive behavioral therapy is effective in the reduction of depression and anxiety in patients with or without *in vitro* fertilization/intracytoplasmic sperm injection treatment. However, the lack of high-quality studies makes it challenging to make a solid and precise conclusion. Well-designed studies should be undertaken in the future to confirm these results.

1. Introduction

Infertility is defined as the failure to achieve a clinical pregnancy after 1 year or more of unprotected intercourse[1]. The prevalence of infertility is 5% to 10% among couples worldwide[2]. Most couples are able to cope with infertility, but others present challenging emotional reactions such as depression, anxiety, frustration, interpersonal problems, feelings of inferiority, suppressed anger, and unconscious feelings of guilt[3–5]. Some studies showed a relationship between depression and infertility, but it has remained a

controversial issue[6,7]. Due to the high prevalence of emotional conditions in infertile women, both early detection and treatment of these disorders are very important[8]. Behavioral therapy is a worthy way to help infertile women for adapting with the emotional problem[9]. Based on the results of a study conducted by Terzioglu, lower anxiety and depression scores were observed by supporting infertility treatment[10]. Other studies showed that leaning

This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 License, which allows others to remix, tweak and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

©2019 Asian Pacific Journal of Reproduction Produced by Wolters Kluwer-Medknow.

All rights reserved.

How to cite this article: Abdolahi HM, Ghojzadeh M, Abdi S, Farhangi MA, Nikniaz Z, Nikniaz L. Effect of cognitive behavioral therapy on anxiety and depression of infertile women: A meta-analysis. *Asian Pac J Reprod* 2019; 8(6): 251-259.

✉ Corresponding author: Leila Nikniaz, Ph.D., Tabriz Health Services Management Research Center, Tabriz University of Medical Sciences, Tabriz, Iran.
E-mail: nikniazleila@gmail.com

confronting methods by infertile women increases psychological health and reduces hopelessness[11–13]. In addition, studies showed that the number of treatment cycles for being pregnant could be reduced by proposing psychological interventions in advance of fertilization treatments[14].

Cognitive-behavioral treatment (CBT) is a psychotherapeutic method based upon a combination of basic behavioral and cognitive research. This treatment has led to positive consequences by improving coping capacities and reducing catastrophic thinking about pain[15]. It is believed that CBT is an effective treatment for different mental problems; however, in the field of infertility, few studies have assessed the success of this type of psychosocial intervention. There is only one systematic review and one meta-analysis reviewed the efficacy of different psychological interventions in infertile couples and the results showed that different types of psychological treatments had no effect on mental health in this population[13,16]. Based on the recommendation of the above studies, research should especially focus on specific types of psychological interventions and specific target population. Therefore, this meta-analysis aimed to evaluate the effect of CBT on anxiety and depression of infertile women.

2. Materials and methods

2.1. Data sources

Published studies until April 2019 were collected for a systematic review of cognitive behavioral therapy on anxiety and depression of infertile women by searching PubMed/MEDLINE, Scopus, EMBASE, Cochrane Library, ISI Web of Science, Proquest and Google scholar. In addition, Iranian databases including Scientific Information Database, Iranian Research Institute for Information Science and Technology, Magiran, IranMedex and Barakat knowledge network system were searched. Keywords were selected based on medical subject heading terms and included (but not limited to): “anxiety OR depression OR mental health OR stress OR psychological health OR quality of life AND infertile women OR infertile female OR infertile couple OR infertile patient AND cognitive behavioral therapy OR cognitive behavioural therapy OR cognitive-behavioral treatment OR CBT”. Also, the Persian equivalence of these key words was searched. The references of recent reviews and other eligible articles were manually searched for additional studies not identified by the electronic search.

2.2. Criteria

2.2.1. Types of studies

Randomized clinical trials or quasi-experimental studies in English or Persian language had surveyed CBT effectiveness. Studies were excluded if performed on samples other than infertile women, other languages rather than Persian and English, articles with poor quality, all articles related to the organizations’ reports, letters to the editor, abstracts presented at conferences, review articles, thesis or suggestions and article by using invalidated questionnaires.

2.2.2. Interventions

The intervention group was restricted to CBT and the control group did not receive CBT.

2.2.3. Outcome

Anxiety and depression of infertile women were reported separately for the CBT group and the control group through the most widely used and validated instruments in psychology as follow: general health questionnaire, Beck depression inventory, Hamilton depression rating scale, depression anxiety stress scales, state-trait anxiety inventory, and Cattell anxiety scale.

The results of included studies should be reported by *P*-value, *t*-value or *F*-value and the sample size of the study population should accurately represent the average effect size of 0.50 with a power of 0.80.

2.3. Data extraction and quality assessment

Two independent reviewers screened all titles and abstracts of the retrieved papers to decide on their inclusion. In addition, full articles of the potentially relevant studies were retrieved and independently screened for eligibility by two reviewers. Disagreements were resolved through discussion, and the reasons for exclusion were recorded for each of the excluded full-text articles. Two researchers also conducted data extraction independently, with disagreements being resolved through discussion.

The following information was subtracted from the studies: the first author, publishing year, sample size, duration of infertility, duration of intervention, dwelling, intervention, and anxiety and depression measurement before and after the intervention. For quality assessment of included trials, Cochrane Collaboration’s tool was applied and selection bias, performance bias, attrition bias, reporting bias, and other bias were examined. Any discrepancies were resolved through discussion and by consulting a third reviewer.

2.4. Quantitative synthesis

All analyses were carried out by using the Comprehensive Meta-Analysis version 2.0. The mean changes of depression and anxiety in both CBT and control groups before and after the study were extracted and if it was not reported, the value was calculated based on the *t*-value between and within groups, the *F*-value or *P*-value. The mean difference between the treatment and control groups was selected to represent the difference in continuous outcomes [with 95% confidential interval (*CI*)].

2.5. Statistical analysis

Heterogeneity was determined by using the *Q* statistic and *I*². A significance level of *P*=0.05 for Cochran’s *Q* test or *I*²>50% was considered as statistically important heterogeneity[17]. Funnel plot and adjusted rank correlation test (methods of Begg and Mazumdar and Eggers) were applied for assessing publication bias.

2.6. Subgroup analysis

We performed a subgroup analysis considering the type of medical treatment. The subgroups were defined as getting assisted reproductive treatment or not (assisted reproductive technology: *in vitro* fertilization (IVF)/intracytoplasmic sperm injection (ICSI) versus no IVF/ICSI).

3. Results

3.1. Selection of eligible studies and study characteristics

The search strategy resulted in 269 titles, which were reduced to 168 following the deletion of duplicates by using the Endnote 7.2.1 literature manager software. Initial assessment of the titles and abstracts reduced the number of papers to 24 (unrelated papers, other

study types =144) and further reduced to 11 on the closer assessment of the full text based on the inclusion and exclusion criteria. In this systematic review, 3 studies due to the use of unknown questionnaires, 1 study due to the lack of pre-test, 5 studies due to the lack of mean and standard deviation report, 2 studies due to the poor credit and lack of mean and standard deviation report, 1 study which was published both in Persian and English, and one study due to lack of suitable interpretation of the questionnaire were excluded. Finally, 11 studies were included in this meta-analysis. A flow chart of literature retrieval is presented in Figure 1.

In the eligible studies, the participants were infertile women who participated in CBT session for handling depression or anxiety. Questionnaires were completed for all patients before and after study. The basic characteristics of the included studies and the mean scores of depression and anxiety were summarized in Table 1. As for publishing year, the included studies were published from 2002 to 2015. And the treatment duration in all studies was 12-24 h.




Figure 1. Flow diagram for selection of articles.

Quality assessment of the included studies was presented in Table 2. All studies had one or more domains characterized as high risk of bias. No study enjoyed a better quality in terms of having more low-risk domains than high-risk ones out of 7 domains. So, all of the included studies were of low quality. Low quality studies were rated as high risk in relation to selection bias, due to a non-random selection of participants, performance and detection bias.

3.2. Participant characteristics

Of the included studies, 10 were conducted in Iran and one in Portugal. In the selected studies, 527 infertile women were enrolled. Among them, 256 and 271 participants were in the CBT group and control group, respectively. All cognitive-behavioral interventions were applied as group psychotherapy (Table 1).

3.3. Meta-analysis results

In most of the eligible studies, several questionnaires were used to measure anxiety and depression. Therefore, each questionnaire was entered as a separate study to meta-analysis. Accordingly, 20 studies (questionnaires) were analyzed.

3.3.1. Depression

Heterogeneity was determined by using the Q statistic and I^2 which was not significant ($Q=38.02$, $P<0.001$, $I^2=76.33$). So, the random

effect model was used. The forest plot was shown in Figure 2. As shown in Figure 2, the between-group differences (CBT and control groups) in risk ratio for infertile women's depression were significant ($d=-1.36$; 95% $CI=-1.81$, -0.90 ; $P<0.001$).

As shown in Figure 3, regarding depression, in both study groups where patients did ($d=-1.39$; 95% $CI=-0.19$, -2.25 ; $P=0.002$) or did not ($d=-1.35$; 95% $CI=-1.91$, -0.79 ; $P<0.001$) receive IVF/ICSI treatment, CBT resulted in a significant decrease in depression.

3.3.2. Anxiety

According to the identification of statistical heterogeneity ($Q=42.86$, $P<0.001$, $I^2=79.00$), the random effect model was used. The forest plot is shown in Figure 4. As shown in Figure 4, the between-group differences (CBT and control groups) in risk ratio for infertile women's anxiety were significant ($d=-0.83$; 95% $CI=-1.18$, -0.47 ; $P<0.001$).

As shown in Figure 5, regarding anxiety, in both study groups where patients did ($d=-0.84$; 95% $CI=-1.32$, -0.35 ; $P=0.001$) or did not ($d=-0.82$; 95% $CI=-1.41$, -0.22 ; $P=0.007$) receive IVF/ICSI treatment, CBT resulted in a significant decrease in anxiety.

3.3.3. Publication bias

Results of the funnel plot and Egger test showed no possibility of publication bias (depression: $t=1.6$, $P=0.14$, anxiety: $t=0.8$, $P=0.40$) (Figure 6).

Table 1. Basic characteristics of included studies and mean scores of depression and anxiety.

Author	Country	Sample size		Mean age (year)	Duration of infertility (year)	Number of sessions (minute)	Questionnaire	Control Mean difference		CBT Mean difference	
		CBT group	The control group					Before study	After study	Before study	After study
Heydari et al. (2002) [18]	Iran	55	55	26.70±4.70	6.10±3.50	12 (90)	STAI-state	43.50±9.70	42.20±11.90	46.40±9.90	31.60±8.10
							STAI-trait	44.10±7.10	42.50±8.80	47.10±11.10	40.60±9.90
Garaee et al. (2003) [19]	Iran	30	30	-	-	15 (60)	STAI-state	43.00±11.20	38.20±10.80	48.60±13.50	39.80±9.50
							STAI-trait	44.80±10.80	41.30±10.30	47.00±10.10	41.70±7.90
Nilforooshan et al. (2006) [20]	Iran	15	15	27.00±4.38	8.00±2.70	6 (120)	BDI	15.30±11.14	20.80±11.91	18.00±9.16	10.66±7.33
Faramarzi M et al. (2008) ^{1,2} [21, 22]	Iran	29	30	28.30±5.30	5.40±3.90	10 (90)	GHQ (Anxiety)	7.80±11.00	6.50±3.90	8.00±4.00	3.20±2.00
							GHQ (depression)	7.20±4.20	7.50±3.90	7.70±4.20	3.60±2.70
							BDI	19.80±8.50	19.70±8.40	20.10±7.90	7.70±4.80
							CAS	38.10±9.10	39.90±8.80	42.30±8.60	30.60±8.60
Hasani et al. (2008) [23]	Iran	10	10	27.70±2.83	5.20±2.28	10 (90)	BDI	14.50±7.61	16.00±9.20	20.40±11.57	10.60±12.52
Hamid (2011) [24]	Iran	20	20	33.63±5.16	11.13±7.41	12 (90)	BDI	61.42±5.83	62.12±13.09	60.21±8.41	26.18±5.91
CAS	21.16±3.27	19.12±4.63	20.32±4.83	14.17±2.12							
Mosalanajad et al. (2012) ¹ [25]	Iran	15	16	35.20	-	15 (90)	DASS-depression	6.72±4.37	5.66±3.72	13.11±4.76	6.41±3.26
							DASS-anxiety	7.72±3.96	7.16±4.60	11.11±4.45	7.17±3.84
Mosalanajad et al. (2012) ² [26]	Iran	32	33	35.20	1.00-1.33	15 (90)	DASS-depression	8.87±3.54	7.80±3.15	14.00±2.38	8.00±2.62
							DASS-anxiety	9.41±3.45	9.25±3.26	13.96±2.59	8.06±2.63
Galhardo et al. (2013) [27]	Portugal	55	37	34.87± 4.20	1.23±0.81	10 (90)	BDI	11.35±9.27	10.22±8.79	11.02±7.05	6.18±4.05
							STAI-state	47.57±15.22	47.57±14.22	47.82±13.09	43.02±8.70
Talaie et al. (2014) [28]	Iran	10	10	21.37	2-5	10 (90)	BDI	17.50±5.20	18.23±6.30	20.00±7.37	14.50±6.94
							HDRS	19.60±6.10	21.00±8.30	21.90±7.23	16.00±10.83

-: unavailable. 1,2= different studies related to one author. STAI: state-trait anxiety inventory; BDI: Beck depression inventory; GHQ: general health questionnaire; CAS: Cattell anxiety scale; DASS: depression anxiety stress scales; HDRS: Hamilton depression rating scale.




Figure 2. Difference between the cognitive-behavioral treatment group and the control group in risk ratio for depression. 1,2 = different studies related to one author; a,b = different questionnaires were used in one study. Std diff in means: Standard difference in means; CI: Confidential interval.




Figure 3. Difference between the cognitive-behavioral treatment group and the control group in risk ratio for depression in IVF/ICSI-treated group and not IVF/ICSI-treated group. IVF: *in vitro* fertilization; ICSI: intracytoplasmic sperm injection. 1,2 = different studies related to one author; a,b = different questionnaires were used in one study; Std diff: Standard difference in means; CI: Confidential interval.




Figure 4. Difference between the cognitive-behavioral treatment group and the control group in risk ratio for anxiety. 1,2 = different studies related to one author; a,b = different questionnaires were used in one study. Std diff in means: Standard difference in means; CI: Confidential interval.




Figure 5. Difference between the cognitive-behavioral treatment group and the control group in risk ratio for anxiety in IVF/ICSI-treated group and not IVF/ICSI-treated group. IVF: *in vitro* fertilization; ICSI: intracytoplasmic sperm injection. 1,2 = different studies related to one author; a,b = different questionnaires were used in one study. Std diff in means: Standard difference in means; CI: Confidential interval.




Figure 6. Funnel plot to evaluate possibility of publication bias. A: depression; B: anxiety. The number of circles is related to the different questionnaires used in 11 studies. At all, 20 questionnaires were analysed. Std diff in means: Standard difference in means.

4. Discussion

Infertility has mental and social as well as reproductive consequences. Assessing the effectiveness of CBT on anxiety and depression of infertile women has been investigated in 11 randomized clinical trials or quasi-experimental studies. These studies comprised 527 infertile women. Of the included studies, 10 were conducted in Iran and one in Portugal. The quality assessment suggested that the overall study quality was fair and no significant publication bias was detected. Despite the subgroup analysis for applying assisted reproductive technology, studies have heterogeneity. Using different questionnaires in different studies caused a heterogeneity which was significant in the case of anxiety. Subgroup analysis was not possible in the case of questionnaires because of the low number of studies. Overall, the results of the included studies do support the use of CBT in the treatment of

infertile women’s depression and anxiety. In contrast to the results of this study, Hammerli *et al*[29] in a meta-analysis study showed that psychological interventions for infertile patients had no effect on mental health. For justifying this discrepancy, we could mention that in the study of Hammerli *et al*, different types of psychological interventions were applied in both men and women by using a wide variety of assessment tools. Also, the results of a systematic review showed no effect of different psychological treatments on infertile couples’ stress and depression; however, these type of treatments had a positive effect on reducing anxiety[16]. Based on the recommendation of the above studies, future research should be focused on the specific target population and specific psychological interventions. It should be considered that men and women are differentially responded to psychological treatments[20]. One reason for these different responses is that women experience higher levels of mental distress than men[30,31]. Regarding the higher levels of

mental distress in infertile women, two explanations have been discussed. One is that women must bear the main burden of medical treatment for infertility and their desire for a child is more important to their life's plan[32,33]. The other is that infertile women feel higher blame for infertility and have a major role in the beginning and following the treatments[34], they have greater emotional investment than men[35] and they spend more time to do the practice. Therefore, it seems that women are more dynamic and responsive in using psychological programs.

The results of some other studies in infertile women supported our finding, which showed that CBT and receiving daily information and support could lower the anxiety and depression scores of infertile women than controls[10,36]. As infertile women experience prolonged, procedural and profound process, cognitive-therapeutic methods are considered as one of the best options due to their direct focus on thoughts and conscious imaginations of individuals. CBT avoids 'waiting stress' so it could be valuable for infertile couples to manage their stress[10,37]. This effect occurs through muscle relaxation and respiratory rate reduction, which results in better controlling of the muscle tension, decreasing oxygen expenditure and carbon dioxide excretion, diminishing heart rate and blood pressure, reducing energy expenditure and at last reducing anxiety and mental stress. These physiological changes result from the reduction of sympathetic nervous system activity[38].

In addition, the mental health of infertile women could be recovered by CBT through the causal cognition model. Depraved outlooks and negative vision are involved in the depression onset and maintenance. Primarily, the negative automatic view would be recovered by a cognitive intervention that mediates depressive signs, and then negative approaches would be replaced with more adaptive ones[39].

It seems that group interventions were significantly more effective than individual counseling interventions[11]. Infertile women feel lonely and suppose that their problem is unique so they cannot communicate their feelings and problems. As a result, it seems that groups supply the safest situation to talk about the unexpressed feelings and consequently psychological discharge[38] as well as for distributing their experiences[40].

The present meta-analysis also found that CBT is effective in the reduction of depression and anxiety in patients with or without IVF/ICSI treatment. This result may be confounded by the duration of infertility and interventions. Women in both groups at different stages of their medical were handling with a variety of indications regarding the medical procedure, which can be confounders. Based on the results of this study and considering the limitations, it seems that setting up counseling centers inside the gynecological clinics would help infertile women to reduce psychological distress and increase rate of successful treatment.

Low risk of subjective data selection was the advantage of our systematic review. Predefined criteria were used for searching databases, screening and assessing articles. Nevertheless, this study had some limitations. First, this study is not registered in

PROSPERO. Also, our analysis had some limitations as follows: negative findings are less likely to be published so publication bias cannot be excluded. In addition, systematic reviews are only as good as the included studies. In this review, only some of the included random controlled trials seemed methodologically sound. Unclear or inadequate allocation concealment, inadequate information about controlling confounders, and applying different questionnaires for evaluating depression and anxiety were the most important problems. Also, most of the studies were conducted in one country which could distort the results.

Based on the relationship between depression and infertility and the effect of counseling based on the CBT approach on the treatment of depression and anxiety, using counseling approaches are recommended. It is suggested that the counseling centers should be set up inside the medical center so that these programs can be applied to help infertile couples with psychiatric problems to promote mental health. Further high-quality investigations on the effect of CBT approach on infertile couples at different stages of their medical handling, different infertility problems and modalities are needed.

Conflict of interest statement

None of the authors had a conflict of interest with respect to this manuscript.

Acknowledgments

We would like to appreciate the editors and referees for considering our study and thanks for absolute support of Research Development and Cooperation Centre and Iranian Centre for Evidence-Based Medicine, Tabriz University of Medical Sciences.

References

- [1] Macaluso M, Wright-Schnapp TJ, Chandra A, Johnson R, Satterwhite CL, Pulver A, et al. A public health focus on infertility prevention, detection, and management. *Fertil Steril* 2010; **93**(1): 16. e1-10.
- [2] Oladokun A, Oladokun R, Morhason-Bello I, Arulogun O, Bamgboye A, Adewole I, et al. Attitude of infertile women to child adoption in Nigeria. *Int J Gynecol Obstet* 2009; **107**: S296.
- [3] Ramazanzadeh F, Noorbala AA, Abedinia N, Naghizadeh MM. Emotional adjustment in infertile couples. *Int J Reprod BioMed* 2009; **7**(3): 97-103.
- [4] Faramarzi M, Pasha H, Esmailzadeh S, Kheirkhah F, Heidary S, Afshar Z. The effect of the cognitive behavioral therapy and pharmacotherapy on infertility stress: A randomized controlled trial. *Int J Fertil Steril* 2013; **7**(3): 199-206.

- [5] Brandes M, Van Der Steen J, Bokdam S, Hamilton C, De Bruin J, Nelen W, et al. When and why do subfertile couples discontinue their fertility care? A longitudinal cohort study in a secondary care subfertility population. *Hum Reprod* 2009; **24**(12): 3127-3135.
- [6] Domar AD, Seibel MM, Benson H. The mind/body program for infertility: A new behavioral treatment approach for women with infertility. *Fertil Steril* 1990; **53**(2): 246-249.
- [7] Domar AD, Clapp D, Slawsby EA, Dusek J, Kessel B, Freizinger M. Impact of group psychological interventions on pregnancy rates in infertile women. *Fertil Steril* 2000; **73**(4): 805-811.
- [8] Noorbala AA, Ramezanzadeh F, Abedinia N, Naghizadeh MM. Psychiatric disorders among infertile and fertile women. *Soc Psychiat Epidemiol* 2009; **44**(7): 587-591.
- [9] Podolska Magdalena Z, Mariola B. Infertility as a psychological problem. *Ginekol Pol* 2011; **82**: 44-49.
- [10] Terzioğlu F. Investigation into effectiveness of counseling on assisted reproductive techniques in Turkey. *J Psychosom Obstet Gynecol* 2001; **22**(3): 133-141.
- [11] Boivin J. A review of psychosocial interventions in infertility. *Soc Sci Med* 2003; **57**(12): 2325-2341.
- [12] De Liz T, Strauss B. Differential efficacy of group and individual/couple psychotherapy with infertile patients. *Hum Reprod* 2005; **20**(5): 1324-1332.
- [13] Hämmerli K, Znoj H, Barth J. The efficacy of psychological interventions for infertile patients: A meta-analysis examining mental health and pregnancy rate. *Hum Reprod Update* 2009; **15**(3): 279-295.
- [14] Campagne DM. Should fertilization treatment start with reducing stress? *Hum Reprod* 2006; **21**(7): 1651-1658.
- [15] Kashikar-Zuck S, Sil S, Lynch-Jordan AM, Ting TV, Peugh J, Schikler KN, et al. Changes in pain coping, catastrophizing, and coping efficacy after cognitive-behavioral therapy in children and adolescents with juvenile fibromyalgia. *J Pain* 2013; **14**(5): 492-501.
- [16] Ying L, Wu LH, Loke AY. The effects of psychosocial interventions on the mental health, pregnancy rates, and marital function of infertile couples undergoing *in vitro* fertilization: A systematic review. *J Assisted Reprod Genet* 2016; **33**(6): 689-701.
- [17] Higgins JP, Green S. *Cochrane handbook for systematic reviews of interventions*. Chichester: John Wiley & Sons; 2011.
- [18] Heydari P, Latifnezhad R, Sahebi A, Jahanian M, Mazloom SR. Effects of cognitive behavior therapy on treatment of anxiety in infertility patients under IUI. *J Reprod Infertil* 2002; **11**: 40-51.
- [19] Garaee V, Mazaheri ML, Sahebi A, Peivandi S, Agahoseini M. Effects of cognitive behavior therapy on treatment of anxiety in infertility patients under GIFT and ZIFT. *J Reprod Infertil* 2003; **5**: 170-180.
- [20] Nilforooshan P, Ahmadi A, Abedi MR, Ahmadi M. Studying the effect of cognitive-behavioral counseling based on interacting cognitive subsystems on depression of infertile couples. *Middle East Fertil Soc J* 2006; **11**(1): 43-47.
- [21] Faramarzi M, Alipour A, Esmaelzadeh S, Kheirkhah F, Poladi K, Pash H. Treatment of depression and anxiety in infertile women: Cognitive behavioral therapy versus fluoxetine. *J Affect Dis* 2008; **108**(1-2): 159-164.
- [22] Faramarzi M, Kheirkhah F, Esmaelzadeh S, Alipour A, Hjjahmadi M, Rahnama J. Is psychotherapy a reliable alternative to pharmacotherapy to promote the mental health of infertile women? A randomized clinical trial. *Eur J Obstet Gynecol Reprod Biol* 2008; **141**(1): 49-53.
- [23] Hassani F, Navabinejad SH, Nooranipoor R. Comparison of the efficacy of cognitive behavioral and emotion-focused therapy on depression of infertile couples. *Women Res* 2008; **6**(3): 61-83.
- [24] Hamid N. Effectiveness of stress management on depression, anxiety and infertility in women. *J Behav Sci* 2011; **5**(1): 55-60.
- [25] Mosalanejad L, Khodabakhshi Koolae A, Morshed Behbahani B. Looking out for the secret wound: The effect of e-cognitive group therapy with emotional disclosure on the status of mental health in infertile women. *Int J Fertil Steril* 2012; **6**(2): 87-94.
- [26] Mosalanejad L, Khodabakhshi Koolae A, Jamali S. Effect of cognitive behavioral therapy in mental health and hardiness of infertile women receiving assisted reproductive therapy (ART). *Iran J Reprod Med* 2012; **10**(5): 483-488.
- [27] Galhardo A, Cunha M, Pinto-Gouveia J. Mindfulness-based program for infertility: Efficacy study. *Fertil Steril* 2013; **100**(4): 1059-1067.
- [28] Talaei A, Kimiaei SA, Borhani M, Mohareri F, Talaei A, Kanegaei R. The effect of cognitive behavioral therapy on depression in infertile women. *IJOGI* 2014; **17**(94): 1-9.
- [29] Haemmerli K, Znoj H, Berger T. Internet-based support for infertile patients: A randomized controlled study. *J Behav Med* 2010; **33**(2): 135-146.
- [30] Wischmann T. Psychosocial aspects of fertility disorders. *Der Urologe Ausg A* 2005; **44**(2): 185-194.
- [31] Covington SN, Burns LH. *Infertility counseling: A comprehensive handbook for clinicians*. Cambridge: Cambridge University Press; 2006.
- [32] Abbey A, Halman LJ, Andrews FM. Psychosocial, treatment, and demographic predictors of the stress associated with infertility. *Fertil Steril* 1992; **57**(1): 122-128.
- [33] Edelmann RJ, Humphrey M, Owens DJ. The meaning of parenthood and couples' reactions to male infertility. *Br J Med Psychol* 1994; **67**(3): 291-299.
- [34] Pasch LA, Dunkel-Schetter C, Christensen A. Differences between husbands' and wives' approach to infertility affect marital communication and adjustment. *Fertil Steril* 2002; **77**(6): 1241-1247.
- [35] Hirsch AM, Hirsch SM. The effect of infertility on marriage and self-concept. *J Obstet Gynecol Neonat Nurs* 1989; **18**(1): 13-20.
- [36] Hansson M, Bodlund O, Chotai J. Patient education and group counselling to improve the treatment of depression in primary care: A randomized controlled trial. *J Affect Disorders* 2008; **105**(1): 235-240.
- [37] Kupka MS, Dorn C, Richter O, Schmutzler A, van der Ven H, Kulczycki A. Stress relief after infertility treatment—spontaneous conception, adoption and psychological counselling. *Eur J Obstet Gynecol Reprod Biol* 2003; **110**(2): 190-195.
- [38] Hamzehpour T. The effect of cognitive behavioural therapy on anxiety in infertile women. *Eur J Exp Biol* 2014; **4**(1): 415-419.
- [39] McNaughton-Cassill ME, Bostwick JM, Arthur NJ, Robinson RD, Neal GS. (eds). Efficacy of brief couples support groups developed to manage the stress of *in vitro* fertilization treatment. *Mayo Clin Proc* 2002; **77**: 1060-1066.
- [40] Frits M, Speroff L. *Clinical gynecologic endocrinology & infertility*. Philadelphia: Lippincott Williams & Wilkins; 2011.