



# Utilization of phosphate solubilizing bacteria (PSB) for sustainable agriculture

Dalvi Sanjay M<sup>1\*</sup> and Rakh RR<sup>2</sup>

<sup>1</sup>Department of Botany, Shri Guru Buddhiswami Mahavidyalaya, Purna (Jn.) – 431511

<sup>2</sup>Department of Microbiology, Shri Guru Buddhiswami Mahavidyalaya, Purna (Jn.) – 431511

\*Email: [sanjaydalvi4u@gmail.com](mailto:sanjaydalvi4u@gmail.com) | [drrrrakh@gmail.com](mailto:drrrrakh@gmail.com)

## Manuscript details:

Received: 10.11.2019  
Accepted: 05.12.2019  
Published: 30.12.2019

### Cite this article as:

Dalvi Sanjay M and Rakh RR (2019) Utilization of phosphate solubilizing bacteria (PSB) for sustainable agriculture, *Int. J. of Life Sciences*, Volume 7(4): 691-699.

**Copyright:** © Author, This is an open access article under the terms of the Creative Commons Attribution-Non-Commercial - No Derives License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

Available online on  
<http://www.ijlsci.in>  
ISSN: 2320-964X (Online)  
ISSN: 2320-7817 (Print)

## ABSTRACT

After Nitrogen Phosphorus is the second most required macronutrient for plants. It is required from molecular level to physical development of plants. Most of the soils contain high levels of Phosphorus. However P forms many insoluble complexes with Calcium, Iron, and Aluminium. It makes the nutrient a paradox. It is reported to be critical factor of many crop production systems due to limited plant available P forms in soil. 80% Phosphorus from soil remains unutilized. Phosphate solubilizing bacteria provide an eco-friendly alternative to convert insoluble phosphate into soluble forms. Species of PSB like Bacillus, Rhizobium and Pseudomonas have ability to release metabolites such as organic acids to carry out mineral phosphate solubilization. The present review is focused on an urgent need of shifting towards a more sustainable agriculture by using PSB.

**Keywords:** Rhizosphere, Phosphorus, Phosphate Solubilizing Bacteria.

## INTRODUCTION

After Nitrogen Phosphorus (P) is the second most essential macronutrient, for growth and development of plants as it is involved in important metabolic pathways like photosynthesis, respiration, biological oxidation, nutrient uptake, cell division and cell building (Pathak *et al.*, 2017; Gupta *et al.*, 2012). Soils contain high levels of P, but, a greater part of soil Phosphorus, approximately 95-99%, is present in the form of insoluble phosphates and hence cannot be utilized by the plants (Muhammad and Maran, 2012; Kannapiram and Sri Ramkumar, 2011). Unlike the case of Nitrogen there is no large atmospheric source of Phosphorus that can be made Biologically available to plants (Karpagam and Nagalaxmi 2014). Conventional farming system relies on heavy application of chemical phosphorus fertilizers to maintain optimum level of phosphorus in agricultural soils (Ahmed and Kiber, 2014). However, major portions (around 75% in some soils) of the soluble phosphorus are

rapidly immobilized in soil which makes it unavailable for plants (Richardson and Simpson, 2011). Also the repeated use of chemical fertilizers deteriorates soil quality (Gyaneshwar et al, 2002).

The soil surrounding the germinating seed is known as spermosphere (Barillot *et al.*, 2013). This ecological niche is the first habitat used by any developed microorganism by seed inoculation and support microbial activity (Ranjan *et al.*, 2013). This in turn influences plant growth. As the plant grows older the growing roots are surrounded by strongly adhering soil particles and this soil-root interaction zone is the rhizosphere (Kumar *et al.*, 2018; Antoun, 2012). The rhizosphere are said to be the areas of very high biological diversity teaming with many different organisms, as the plant roots provide food, shelter and energy (Barillot *et al.*, 2013). These active microbial population can either exert beneficial or detrimental effects or they can be neutral (Sunder-Rao and Sinha, 1963). The success in the use of beneficial microorganisms requires an excellent understanding between the different components of the complex plant-soil-microorganisms (Gyaneshwar *et al.*, 2002)

Natural solubilization of mineral phosphates is an important mechanism exhibited by different rhizospheric microorganisms, known as phosphate solubilizing microorganisms (PSM) (Selvi *et al.*, 2017). Bacteria are the predominant microorganisms that solubilize mineral phosphate in nature, as compared to other microorganisms (Paul and Sinha, 2017; Debojyoti *et al.*, 2015). Phosphate solubilizing bacteria (PSB) play an important role in biogeochemical phosphorus cycling in both terrestrial and aquatic environments (Das *et al.*, 2007). Application of phosphate solubilizing bacteria increases soil fertility due to their ability to convert insoluble P to soluble P by releasing organic acids, chelation and ion exchange (Muhamad and Maran, 2012; Selvi *et al.*, 2017; Tarafdar and Classon, 1988). The present review emphasises the role of Phosphorus in plant nutrition, PSB, mechanism of phosphate solubilization and utilization of PSB for sustainable agriculture.

#### **ROLE OF PHOSPHORUS IN PLANT NUTRITION**

Phosphorus (P) is a most important growth limiting nutrient for plants (Richardson and Simpson, 2011). Unlike the case of Nitrogen, there is no large distinctive source that can be naturally managed for crops accessibility (Sharma *et al.*, 2013). P, being an

important constituent element of nucleic acids, enzymes, coenzymes, nucleotides and phospholipids, is involved in the transformation of energy, transfer of hereditary characters and cell organization in plants (Rodriguez and Fraga, 1999; Goldstein, 1986). Nearly all phases of plant cycle including root growth, stalk and stem strength, photosynthesis and respiration, flowering and anthesis, seed formation, crop maturity and production N-fixation in legumes, crop quality and resistant to plant diseases are the attributes associated with P nutrition (Balemi and Negisho, 2012; Satyaprakash *et al.*, 2017). Its deficiency causes stunted growth and severe yield losses. Its concentration in soil solution is very low, because soluble forms of P are fixed by soil solid phase, making less than 0.01% of total P available to plants (Trolov *et al.*, 2003). Phosphorous is therefore, one of the least mobile nutrients in soil (Balemi and Negisho, 2012; Gyaneshwar, 2012). Worldwide soils are supplemented with inorganic P as chemical fertilizers to support crop production, but repeated use of fertilizers deteriorates soil quality (Vessey, 2003). In India about 98% of soils including fertile ones are deficient in P, as the concentration of free P (available form for plants) is not more than 10µm even at pH 6.5 (Debojyoti *et al.*, 2015). Thus, most of the Indian soils are poor and marginal in phosphorus and require adequate P-fertilization to sustain high productivity and profitability (Debojyoti *et al.*, 2015). Phosphorus mainly occurs in three forms (i) soluble inorganic-P which occurs in soil solution (ii) insoluble inorganic P occurring as primary orthophosphate ( $H_2PO_4$ ) and (iii) secondary orthophosphate ( $HPO_4^{2-}$ ) (Goldstein, 1986). The form of phosphate taken up by the plant from soil solution is phosphate anions mainly  $H_2PO_4^-$  and  $HPO_4^{2-}$  (Richardson, 2001). Many agricultural soils surround huge deposition of total P, generally from 200 to 5000 mg P kg<sup>-1</sup> of soil by mean value of 600 mg P kg<sup>-1</sup> of soil and its accumulation based on frequent use of inorganic fertilizers and sludge from the treated wastewaters (Rodriquez and Fraga, 1999). Much of the P which is present in soil and provided to the crops through inorganic fertilizers become unavailable through precipitation by reacting with  $Fe^3$  and  $Al^{3+}$  in acidic and with  $Ca^{2+}$  in calcareous soils, to form Aluminium Phosphate ( $AlPO_4$ ) and Ferrous Phosphate ( $FePO_4$ ) which are sparingly soluble precipitates (Sharpley, 1985) and ultimately resulting in the low concentration of P in soils making it unavailable to plants (Pikovskaya, 1948). Therefore, crop plants can utilise only a fraction of applied P and this make them

poor in performance (Abbasi *et al.*, 2015). The concentration of P in soil solution is very low, varying from 0.001mg L<sup>-1</sup> in very poor soils to 1mg L<sup>-1</sup> in heavily fertilized soils (Seema *et al.*, 2013)

### PHOSPHATE SOLUBILIZING BACTERIA (PSB)

The fact that certain soil microbes are capable of dissolving relatively insoluble phosphatic compounds has opened the possibility for inducing microbial solubilization of phosphates in soil (Ahmed and Kiber, 2014). Phosphate solubilizing microorganisms are the one which solubilize the insoluble form of phosphate in the soil to make it available for the plant growth (Eivazi and Tabatabai, 1977). Phosphate solubilizing microorganisms include several Bacteria, Fungi, Actinomycetes, Yeast and Cyanobacteria (Saxena *et al.*, 2016; Yazdani *et al.*, 2009; Chen *et al.*, 2006; Khan and Zaidi, 2006). But bacteria are the predominant microorganisms that solubilize mineral phosphate in nature, as compared to other microorganisms (Paul and Sinha, 2017; Zaidi and Khan, 2007).

PSB play a vital role in mobilizing Phosphate for the use of plants from the native soil Phosphorous pools as well as rock phosphates (Sharma *et al.*, 2013; Trolov *et al.*, 2003). Several phosphate solubilizing bacteria can utilize insoluble phosphate sources such as tri-calcium phosphate (TCP), hydroxyapatite, flourapatite, ferric, Aluminium and magnesium phosphate, bone meal and rock phosphates and convert them into soluble forms (Muhamad and Maran, 2012; Gaur *et al.*, 1973; Eivazi and Tabatabai, 1977). The important genera of P-solubilizing bacteria include *Achromobacter*, *Aerobacter*, *Alcaligenes*, *Azotobacter*, *Bacillus*, *Escherichia*, *Pseudomonas*, *Serratia* and *Xanthomonas* (Linu *et al.*, 2009; Afzal and Bano, 2008; El-Tarabily *et al.*, 2008; Premono *et al.*, 1996), *Azospirillum* (Mehmet *et al.*, 2005). Among these, bacteria belonging to genera *Pseudomonas*, *Bacillus* and *Rhizobium* and fungi belonging to genera *Penicillium* and *Aspergillus* possess the greater ability to solubilize the insoluble phosphates (Singh and Siddiqui, 2015, Chhabot *et al.*, 1996). The efficiency of these strains depends on various factors such as temperature, soil, pH, type of insoluble phosphate etc. (Ahmed and Kiber 2014, Vessey, 2003; Eivazi and Tabatabai, 1977). The phosphate solubilizing bacteria can be isolated from different sources such as soil (Barillot *et al.*, 2013 Chen *et al.*, 2006; Mehta *et al.*, 1954), rhizosphere (Dalvi *et al.*, 2019; Gyaneshwar *et al.*, 2002), rootnodules (Vikram and Hamzehzarghani, 2008) and compost

(Zalate and Padmani, 2009; Panwar *et al.*, 2002)). Inoculation with phosphate solubilizing organisms in multilocational trials conducted with wheat, paddy, chickpea, maize, greengram, lentils and potato have shown increased yields (Trolov *et al.*, 2003).

### MECHANISM OF PHOSPHATE SOLUBILIZATION

Bacteria are the predominant microorganisms that solubilize mineral phosphate in nature, as compared to other microorganisms (Paul and Sinha, 2017). Phosphate solubilizing bacteria (PSB) play an important role in biogeochemical phosphorus cycling in both terrestrial and aquatic environments (Das *et al.*, 2007). Application of phosphate solubilizing bacteria increases soil fertility due to their ability to convert insoluble P to soluble P by releasing organic acids, chelation and ion exchange (Omar, 1998; Narula *et al.*, 2000.).

Phosphate solubilization takes place through various microbial processes/mechanisms including organic acid production and proton extrusion, the principal mechanism being the lowering of soil pH by microbial production of organic acids or the release of protons and mineralization by producing acid phosphatases (Tarafdar and Claasen, 1988), ultimately resulting in P availability in soil (Whitelaw, 2000). Soil phosphates mainly of phosphatic fertilizers under alkaline conditions are fixed in the form of insoluble phosphates. Many of the calcium phosphates, including rock phosphate ores (fluoroapatite, francolite), are insoluble in soil. Their solubility increases with a decrease of soil pH ( Gyaneshwar *et al.*, 2002).

Phosphobacteria have been found to produce some organic acids such as monocarboxylic acid (acetic, formic), monocarboxylic hydroxy (lactic, glucenic, glycolic), monocarboxylic, ketoglucenic, decarboxylic (oxalic, succinic), dicarboxylic hydroxy (malic, maleic) and tricarboxylic hydroxy (citric) acids in order to solubilize inorganic phosphate compounds (Muhamad and Maran, 2012; Eivazi and Tabatabai, 1977). The type of organic acid produced and their amounts differ with different organisms (Karpagam and Nagalaxhmi, 2014). Among them, gluconic acid and 2-ketogluconic acid seems to be the most frequent agent of mineral phosphate solubilization (Deubel *et al.*, 2000; Song *et al.*, 2008). The organic acids are the products of the microbial metabolism, mostly by oxidative respiration or by fermentation of organic carbon sources like glucose (Trolove *et al.*, 2003).

Phosphate solubilization is the result of combined effect of pH decrease and organic acids production (Fankem *et al.*, 2006, Suranga and Kumar, 1993). PSB decrease the soil pH by producing organic acids (Sardina *et al.*, 1986). These organic acids compete with the P binding sites in the soil (Nahas, 1996). Through Organic acid production PSB mobilizes P from sparingly soluble phosphates and convert into soluble forms mainly by the chelation mediated mechanism (Whitelaw, 2000). Acidification of the surroundings of microbial cell releases proton through the production of organic acids (Villegas and Fortin, 2002). These metabolite organic acids as a result of anion exchange of P by an acid anion can either directly dissolve the mineral P or can chelate  $Al^{3+}$ ,  $Fe^{3+}$ ,  $Ca^{2+}$  ions associated with inorganic P (Gupta *et al.*, 1993). The carboxyl and hydroxyl groups chelate the cation bound to inorganic P and convert it into soluble forms (Sagoe, 1998). As the soil pH increases the divalent and trivalent forms of inorganic P,  $HPO_4^{2-}$  and  $HPO_4^{3-}$  occur in the soil (Suranga and Kumar, 1993). The organic acids released by PSB into the soil consequently decreases the soil pH and acidifies the surrounding environment and lead to the release of P ions by  $H^+$  substitution for the cation bound to phosphate (Goldstein, 1986) making the P available to plants. PSB mineralize soil organic P by the production of acid phosphatases. Release of organic anions, and production of Siderophores and acid phosphatase by microbes, hydrolyze the soil organic P or split P from organic residues resulting in P availability (Dodor and Tabatabai, 2003).

#### UTILIZATION OF PHOSPHATE SOLUBILIZING BACTERIA (PSB) FOR SUSTAINABLE AGRICULTURE

As a result of energy crisis, environmental hazards and depleting soil fertility, man needs to shift towards a more sustainable agriculture (Kumar *et al.*, 2018). Chemical fertilizers should be replaced with biofertilizers, the latter being eco-friendly, productive and accessible (Sheraz *et al.*, 2010). PSB play a crucial role in bio-geo phosphorus cycle as they affect the transformation of the soil phosphates (Sharpley, 1985). They increase nutrient uptake from soil, thus reducing the need for fertilizers and preventing accumulation of phosphates in soil. The concentration and number of PSB in rhizosphere can be inadequate for the complete supply of inorganic phosphates to the plants (Sunder-Rao and Sinha, 1963). Therefore inoculation of the plants with these phosphate

solubilizers is essential for efficient phosphate solubilization (Karpagam and Nagalaxmi, 2014).

Several workers concluded that application of phosphate-solubilizing microorganisms increases soil fertility and results in growth promotion of crops (Omar 1998; Narula *et al.*, 2000). The positive effect of P solubilizers has been reported on food and fodder crops (Dalvi *et al.*, 2019; Santana *et al.*, 2016; Gupta *et al.*, 2012). Phosphate solubilizing bacteria are being used as biofertilizers since 1940's (Gerretsen, 1948; Krasilinikov, 1957; Vssey, 2003). Inoculation of seeds with PSB is a promising technique which alleviates the deficiency of P (Qureshi *et al.*, 2012). Inoculation with PSB such as *Pseudomonas* (Husen *et al.*, 2009), *Bacillus* (Turan *et al.*, 2007; Mehejibin and Patel, 2007), *Rhizobium* (Afzal and Bano, 2008; Chabot *et al.*, 1996), *Micromonospora* (El-Tarabily *et al.*, 2008), *Burkholderia* (Minaxi and Saxena, 2010), *Azotobacter* (Naseri and Mirzai, 2010), *Acinetobacter* (Gulati *et al.*, 2010), *Azospirillum* (Nesri and Mirzai, 2010; Mehmet *et al.*, 2005) and *Gluconacetobacter* (Linu *et al.*, 2009) has been reported in increasing solubilization of fixed P ensuring high crop yields (Rodriguez and Fraga, 1999).

Bacteria belonging to the genus *Bacillus*, *Pseudomonas*, and *Rhizobium* have proven to be powerful PSB (Saxena *et al.*, 2016; Noori and Saud, 2012; Turan *et al.*, 2007, Mhejibin *et al.*, 2007). Savankumar and Samiyappan, 2006, studied ACC deaminase from *Pseudomonas fluorescens* mediated saline resistance in groundnut. Bhatia *et al.*, 2008, Showed beneficial effects of fluorescent Pseudomonads on growth promotion and suppression of charcoal rot in groundnut. Studies on *Pseudomonas* and *Trichoderma* proved phosphate solubilisation and antagonistic activities against *Fusarium oxysporum* and *Rhizoctonia solani* (Jay *et al.*, 2014). *Rhodococcus* sp, *Pseudomonas* sp. and *Arthrobacter nicotinovorans* when inoculated with *Zea mays* and grown in P deficient soils amended with tricalcium phosphate enhanced the plant growth (Sofia and Paula, 2014). *Pseudomonas* sp., *Serratia marcescens* and *Bacillus cereus* were known to reduce the bacterial wilt caused by *Ralstonia solanacearum* in tomato (Sing and Siddiqui, 2015) and hence proved to be effective biocontrol agents (Henok and Kerstin, 2013). Bano and Mussarat, 2003, Naseby *et al.*, 2001 and Haas and Keel, 2003, also observed biocontrol properties of *Pseudomonas* strains.

Several workers reported potency of PSB in solubilising Phosphorous and increasing yield in crops as Tomato (Pathak *et al.*, 2017; Sharon *et al.*, 2016; Lamsal, 2013 ; Poonia and Dhaka, 2012; Hariprasad and Niranjana, 2009), Maize (Mohamed *et al.*, 2017; Hussain *et al.*, 2013; Gholami *et al.*, 2009 ; Yazdani *et al.*, 2009; Nadeem *et al.*, 2010), Wheat (Sachdev *et al.*, 2009; Afzal and Bano, 2008; Babana and Antoun, 2006; Mehmet *et al.*, 2005; Khalid *et al.*, 2004; Nerula *et al.*, 2000), Soyabean (Husen *et al.*, 2009; El-Tarabily *et al.*, 2008), Moog bean (Walpolia and Yoon, 2013; Jha *et al.*, 2012; Minaxi and Saxena, 2010; Vikram *et al.*, 2008), Cicer (Sharma *et al.*, 2015; Zaidi and Khan, 2007), Groundnut (Zalate and Padmani, 2009; Bhatia *et al.*, 2008; Panwar *et al.*, 2002), Cotton Qureshi *et al.*, (2012), Sunflower (Zehara, 2010), Safflower (Zhang *et al.*, 2019), Sugarcane (Sundara *et al.*, 2002), Sorghum (Gopalkrishna and Humayun, 2011) , Chilli (Abbasi *et al.*, 2015) and Mash bean (Niazi *et al.*, 2015).

Some workers noticed additional features of PSB to facilitate plant growth and development in the presence of various stresses. *Pseudomonas* sp. enhanced plant growth in salt stress (Nadeem *et al.*, 2010; Mohammad *et al.*, 1998). *Acinetobacter* (Gulati *et al.*, 2010), *Pseudomonas putida* (Pandey *et al.*, 2006) and *Pseudomonas corrugate* (Trivedi and Sa, 2008) are observed to be cold tolerant. *Pseudomonas* sp. (Sandhya *et al.*, 2010), *Arthrobacter* sp., *Bacillus* sp. (Banerjee *et al.*, 2010) showed tolerance towards drought stress. *Pseudomonas fluorescens* *Bacillus* and *Hallobacillus* showed tolerance towards salinity stress (Saravanakumar and Samiyappan, 2006).

The review concludes that phosphate solubilizing bacteria promise a better alternative to the chemical fertilizers, the latter being costly, deteriorating the agriculture soils and environmental concern. Phosphate solubilizing biofertilizers will prove eco-friendly and cost effective agro technology to improve crop production. More research in this field is needed to minimize the use of chemical fertilizers and make use of biofertilizers in large scale for sustainable agriculture.

#### Acknowledgement:

The authors are grateful for the financial support provided by Swami Ramanand Teerth Marathwada University, Nanded, through the research project of Rajiv Gandhi Science and Technology Commission (RGSTC), Government of Maharashtra, sanctioned to

Dr. Sanjay M. Dalvi, Department of Botany, Shri Guru Buddhiswami Mahavidyalaya, Purna (Jn.).

#### REFERENCES

- Abbasi MK, Musa N and Manzoor M (2015) Phosphorus release capacity of soluble P fertilizers and insoluble rock phosphate in response to phosphate solubilizing bacteria and poultry manure and their effect on plant growth promotion and P utilization efficiency of chilli (*Capsicum annum* L.), *Biogeosci. Discuss*, 12:1839-1873.
- Afzal A and Bano A (2008) *Rhizobium* and phosphate solubilising bacteria improve the yield and phosphorus uptake in wheat (*Triticum aestivum*), *Int J Agri Biol*, 10: 85-88.
- Ahemad M and Kiber M (2014) Mechanisms and applications of plant growth promoting rhizobacteria: Current perspective, *J. King Saud Uni. Sci.* 26: 1-20.
- Antoun H (2012) Beneficial Microorganisms for the Sustainable Use of Phosphates in Agriculture. *Procedia Engineering*, 46: 62 – 67.
- Babana AH and Antoun H (2006) Effect of Tlesmi phosphate rock solubilizing microorganisms on phosphorus uptake and yield of field grown wheat (*Triticum aestivum* L.) in Mali, *Plt Soil*, 287(1-2): 51-58.
- Balemi T and Negisho K (2012) Management of soil phosphorus and plant adaptation mechanisms to phosphorus stress for sustainable crop production: A review, *J. Soil Sci. Plant Nutr.* 12 (3): 547-561.
- Banerjee S, Palit R, Sengupta C and Standing D (2010) Stress induces phosphate solubilisation by *Arthrobacter* sp. and *Bacillus* sp. isolated from tomato rhizosphere, *Aust. J. Crop Sci.*, 4: 378-383.
- Bano N and Musarrat J (2003) Characterization of a new *Pseudomonas aeruginosa* strain NJ-15 as a potent Biocontrol agent, *Curr Microbiol*, 46: 324-327.
- Barillot CDC, Sarde CO, Bert V, Tarnaud E and Cochet N (2013) A standardized method for the sampling of rhizosphere and rhizoplane soil bacteria associated to a herbaceous root system, *Ann. Microbiol.*, 63(2): 471-476.
- Bhatia S, Maheshwari DK, Dubey RC, Arora DS, Bajpai VK and Kang SC (2008) Beneficial effects of fluorescent *Pseudomonads* on seed germination, growth promotion and suppression of charcoal rot in Groundnut (*Arachis hypogea* L.) *J Microbiol Biotechnol*, 18(9): 1578-1583.
- Chabot R, Anton H and Cescas MP (1996) Growth promotion of maize and lettuce by phosphatesolubilizing *Rhizobium leguminosarum* biovar. *Phaseoli*, *Plant and Soil* 184: 311-321.

- Chen YP, Rekha PD, Arunshen AB, Lai WA and Young CC (2006) Phosphate solubilizing bacteria from subtropical soil and their tricalcium phosphate solubilizing abilities, *Appl. Soil Ecol.*, 34: 33-41.
- Dalvi SM, Rakh RR, Kadam VN and Nagthane V (2019) Screening for phosphate solubilizing bacteria (PSB) from rhizospheric soil, *Ajanta*, 8(1): 130-136.
- Das S, Lyla PS and Khan SA (2007) Biogeochemical processes in the continental slope of Bay of Bengal: I. bacterial solubilization of inorganic phosphate, *Rev. Biol. Trop.* 55: 1 – 9.
- Debojyoti R, Paul M and Banerjee S (2015) Isolation Identification and Characterization of Phosphate Solubilising Bacteria from Soil and the Production of Biofertilizer, *Int. J. Curr. Microbiol. App. Sci* , 4(11): 808-815
- Deubel A, Gransee and Merbach W (2000) Transformation of organic rhizodeposits by rhizoplane bacteria and its influence on the availability of tertiary calcium phosphate, *J. Plant Nutr. Soil Sci.*, 163: 387-392.
- Dodor DE and Ali Tabatabai M (2003) Effect of cropping systems on phosphatases in soils, *J. Plant Nutr. Soil Sci.*, 166: 7–13.
- Eivazi F and Tabatabai MA (1977) Phosphatases in soils, *Soil Biology and Biochemistry* 9: 167-172.
- El-Tarabily KA, Nassar AH and Sivasithamparam K (2008) Promotion of growth of bean (*Phaseolus vulgaris* L.) in a calcareous soil by a phosphatesolubilizing, rhizosphere-competent strain of *Micromonospora endolithica*, *Applied Soil Ecology*, 39: 161-171.
- Fankem H, Nwaga D, Deubel A, Dieng L, Merbach W and Etoa FX (2006) Occurrence and functioning of phosphate solubilizing microorganisms from oil palm tree (*Elaeis guineensis*) rhizosphere in Cameroon, *African J. Biotech.*, 5: 2450-2460.
- Gaur AC, Mandan M and Ostwal KP (1973) Solubilization of phosphatic compounds by native microflora of rock phosphates. *Indian J Experi Biol*, 11: 427- 429.
- Gerretsen FC (1948) The influence of microorganisms on the phosphorus uptake by the plant, *Plant and Soil*, 1: 51-81.
- Gholami A, Shahsavani S and Nezarat S (2009) The Effect of Plant Growth Promoting Rhizobacteria (PGPR) on Germination, Seedling Growth and Yield of Maize, *Int. J. Biol. Biomol. Agric. Food Biotechnol. Eng.*, 3(1): 9–14.
- Goldstein AH (1986) Bacterial solubilization of mineral phosphates : Historical perspective and future prospects, *Am. J. Altern. Agric.*, 1(2): 51–57.
- Gopalakrishnan S and Humayun P (2011) Evaluation of bacteria isolated from rice rhizosphere for biological control of charcoal rot of sorghum caused by *Macrophomina phaseolina* (Tassi) Goid, *World J. Microbiol. Biotechnol.*, 27: 1313–1321.
- Gulati A, Sharma and Vyas P (2010) Organic acid production and plant growth promotion as a function of phosphate solubilization by *Acinetobacter rhizosphaerae* strain BIHB 723 isolated from the cold deserts of the trans-Himalayas, *Archives of Microbiology*, 192(11): 975–983.
- Gupta M, Kiran S, Gulati A, Singh V and Tewari R (2012) Isolation and identification of phosphate solubilizing bacteria able to enhance the growth and Aloiin: A biosynthesis of *Aloe barbadensis* Miller, *Microbiol Res*, 167:358-363.
- Gupta R, Shanker AB, Saxena RK and Kuhad RC (1993) Solubilization of low grade Indian rock phosphates and inorganic phosphates by *Bacillus licheniformis*, *Folia Microbiologica*, 38: 274-276.
- Gyaneshwar P, Kumar GN, Parekh LJ and Poole PS (2002) Role of soil microorganisms in improving P nutrition of plants, *Plant Soil*, 245: 83-93.
- Haas D and Keel C (2003) Regulation of antibiotic production in root-colonizing *Pseudomonas* sp. And relevance for biological control of plant disease, *Annu Rev Phytopathol*, 41: 117-153.
- Hari Prasad P and Niranjana SR (2009) Isolation and characterization of phosphate solubilizing rhizobacteria to improve plant health of tomato, *Plant Soil*, 316: 13–24.
- Henok K and Kerstin W (2013) Characterization of plant growth promoting rhizobacteria and their potential as bioprotectant against tomato bacterial wilt caused by *Ralstonia solanacearum*. *Biol. Control*, 67: 75–83.
- Husen, E, Wahyudia AT, Suwanto A and Saraswati R (2009) Soybean seedling root growth promotion by 1-aminocyclopropane-1-carboxylate deaminase producing *Pseudomonads*, *Indon J Agric Sci*, 10: 19-25.
- Hussain MI, Asghar HN, Akhtar MJ and Arshad M (2013) Impact of phosphate solubilizing bacteria on growth and yield of maize, *Soil Environ*, 32(1): 71-78.
- Jay PV, Janardan Y, Kavindra NT and Durgesh KJ (2014) Evaluation of plant growth promoting activities of microbial strains and their effect on growth and yield of chickpea (*Cicer arietinum* L.) in India, *Soil Biol. Biochem.*, 70: 33-37.
- Jha A, Sharma D and Saxena J (2012) Effect of single and dual phosphate solubilizing bacterial strain inoculations on overall growth of mung bean plants, *Arch Agr Soil Sci*, 58: 967-971.
- Kannapiran E and Sri Ramkumar V (2011) Isolation of phosphate Solubilizing bacteria from sediments of Thondi coast, Palk Strait, Southeast coast of India, *Annals of Biological Research*. 25:157-163.

- Karpagam T and Nagalaxhmi (2014) Isolation and characterization of phosphate solubilizing microbes from agricultural soil, *Int J Curr Microbiol App Sci*, 3(3): 601-614.
- Khalid A, Arshad M and Zahir ZA (2004) Screening plant growth promoting rhizobacteria for improving growth and yield of wheat, *Journal of Applied Microbiology*, 96: 473-480.
- Khan MS and Zaidi A (2006) Influence of composite inoculations of phosphate solubilizing organisms and an arbuscular mycorrhizal fungus on yield, grain protein and phosphorus and nitrogen uptake by green gram, *Arch Agron Soil Sci*. 52: 579-583.
- Krasilinikov NA (1957) On the role of soil micro-organism in plant nutrition. *Microbiologiya*, 26: 659-72.
- Kumar A, Kumar A and Patel H (2018) Role of microbes in phosphorus availability and acquisition by plants, *Int J Curr Microbiol and Appl Sci*, 7(5): 1344-1347.
- Lamsal K (2013) Biocontrol of Late Blight and Plant Growth Promotion in Tomato Using Rhizobacterial Isolates, *J. Microbiol. Biotechnol.*, 23(7) :1-8.
- Linu MS, Stephen J and Jisha MS (2009) Phosphate solubilizing *Gluconacetobacter* sp., *Burkholderia* sp. and their potential interaction with cowpea (*Vigna unguiculata* L.) Walp.), *International Journal of Agriculture Research*, 4: 79-87.
- Mahejibin K, and Patel CB (2007) Plant growth promoting effect of *Bacillus firmus* strain NARS1 isolated from Central Himalayan region of India on *Cicer arietinum* at low temperature, *African Crop Science Conf Proc*, 8: 1179-1181.
- Mehmet O, Cevdet A, Oral D and Aki SM (2005) Single and double inoculation with *Azospirillum/Trichoderma*: The effect on dry bean and wheat, *Biol Fert Soils*, 41: 262-266.
- Mehta N C, Legg JO, Goring CAL and Black CA (1954) Determination of organic phosphorus in soils. I. Extraction method, *Soil Sci Soc of America Proc*, 18: 443-449.
- Minaxi and Saxena J (2010) Disease suppression and crop improvement in moong beans (*Vigna radiate*) through *Pseudomonas* and *Burkholderia* strains isolated from semi arid region of Rajasthan. *Bio Cont*, 55(6): 799- 810.
- Mohamed HM and Almaroai YA (2017) Effect of Phosphate Solubilizing Bacteria on the Uptake of Heavy Metals by Corn Plants in a Long-Term Sewage Wastewater Treated Soil, *International Journal of Environmental Science and Development*, 8(5): 366-371.
- Mohammad M, Shibli R, Ajlouni M and Nimri L (1998) Tomato root and shoot responses to salt stress under different levels of phosphorus nutrition, *J. Plant Nutr.*, 21(8): 37-41.
- Muhammad MA and Maram G (2012) The Effect of Phosphate Solubilizing Bacteria and Organic Fertilization on Availability of Syrian Rock Phosphate and Increase of Triple Superphosphate Efficiency, *World J Agri Sci*, 8 (5): 473-478.
- Nadeem SM, Hussain I, Naveed M, Ashgar HN, Zahir ZA and Arshad M (2010) Performance of plant growth promoting rhizobacteria containing ACCdeaminase activity for improving growth of maize under salt-stressed conditions, *Pakistan J Agril Sci*, 43:114-121.
- Nahas E (1996) Factors determining rock phosphate solubilization by microorganism isolated from soil, *World J. Microb. Biotechnol.*, 12: 18-23.
- Narula N, Kumar V, Behl RK, Duebel AA, Gransee A and Merbach W (2000) Effect of P solubilizing *Azotobacter chroococcum* on NPK uptake in P responsive wheat genotypes grown under greenhouse conditions, *J Plant Nutr Soil Sci*. 163: 393-401.
- Naseby DC, Way JA, Bainton NJ and Lynch JM (2001) Biocontrol of *Pythium* in the pea rhizosphere by antifungal metabolite producing and non-producing *Pseudomonas* strains, *J. Appl Microbiol*, 90 (3): 421-429.
- Naseri R and Mirzaei A (2010) Response of yield and yield components of Safflower (*Carthamus tinctorius* L.) to seed inoculation with *Azotobacter* and *Azospirillum* and different nitrogen levels under dry land condition, *Am-Eurasian J Agric Env Sci*. 9:445-9.
- Niazi MTH, Kashif S, Asghar HN, Saleem M, Khan MY and Zahir ZA (2015) Phosphate solubilizing bacteria in combination with pressmud improve growth and yield of Mash bean, *J Animal Plant Sci*, 25(4):1049-1054.
- Noori MSS and Saud HM (2012) Potential plat growth promoting activity of *Pseudomonas* sp. isolated from paddy soil in Malaysia as Biocontrol agent, *J. Plant Pathol Microbiol*, 3(2): 120-123.
- Omar SA (1998) Role of rock-phosphate-solubilizing fungi and vesicular arbuscular-mycorrhiza (VAM) in growth of wheat plants fertilized with rock phosphate, *World J Microb Biot*, 14: 211-219.
- Pandey A, Trivedi P, Kumar B and Palni LMS (2006) Characterization of a phosphate solubilizing and antagonistic strain of *Pseudomonas putida* (B0) isolated from a sub-alpine location in the Indian Central Himalaya, *Curr. Microbiol.*, 53:102-107.
- Panwar AS, Singh N, Saxena DC and Hazarika UK (2002) Yield and quality of groundnut seed as influence by phosphorus, biofertilizers and organic manures, *Indian J Hill Farming*, 15(1): 68-71.
- Pathak R, Vipassana P, Shrestha A, Lamichhane J, Gauchan DP (2017) Isolation of phosphate solubilizing bacteria and their use for plant growth

- promotion in tomato seedling and plant, *Kathmandu Uni J Sci, Eng and Tech*, 13(2): 61-70
- Paul D and Sinha SN (2017) Isolation and characterization of phosphate solubilizing bacterium *Pseudomonas aeruginosa* KUPSB 12 with antibacterial potential from river Ganga, *Annals of Agrarian Sci*, 15: 130-136.
- Pikovskaya RI (1948) Mobilization of Phosphorus in soil in connection with vital activity of some microbial species, *Microbiology*, 17: 362-370.
- Poonia MK and Dhaka BL (2012) Effect of phosphorus solubilizing bacteria (PSB) on growth and yield in tomato, *J. Hortl. Sci.* 7(1): 104-107.
- Premono ME, Moawad A and Velk P (1996) Effect of phosphate-solubilizing *Pseudomonas putida* on the growth of maize and its survival in the rhizosphere, *Indonesian Journal of Crop Science*, 11: 13- 23.
- Qureshi MA, Ahmad ZA, Akhtar N, Iqbal A, Mujeeb F and Shakir MA (2012) Role of phosphate solubilizing bacteria (psb) in enhancing p availability and promoting cotton growth, *J. Animal & Plant Sci.*, 22(1): 204-210.
- Ranjan A, Mahalakshmi MR and Sridevi M (2013) Isolation and characterization of phosphate-solubilizing bacterial species from different crop fields of Salem, Tamil Nadu, India, *Int J Nutr Pharmacol Neurol Dis*, 3(1): 29-33.
- Richardson AE (2001) Prospectus for using microorganisms to improve the acquisition of phosphorus by plants, *Aust. J. Plant Physiol.*, 28(9): 897-906.
- Richardson AE and Simpson RJ (2011) Soil microorganisms mediating phosphorus availability, *Plant Physiol*, 156: 989-996.
- Rodriguez H & Fraga R (1999) Phosphate solubilizing bacteria and their role in plant growth promotion, *Biotechnol. Adv.*, 17: 319-339.
- Sachdev DP, Chaudhari HG, Kasture VM, Dhavale DD and Chopde BA (2009) Isolation and characterization of Indol Acetic Acid (IAA) producing *Klebsiella pneumoniae* strains from Rhizosphere of wheat (*Triticum aestivum*) and their effect on plant growth, *Indian J Exp Biol*, 47: 993-999.
- Sandhya V, Ali S, Grover G, Reddy G and Venkateswarlu B (2010) Effect of plant growth promoting *Pseudomonas* spp. on compatible solutes, antioxidant status and plant growth of maize under drought stress, *Plant Growth Regul.*, 62: 21-30.
- Santana EB, Marques ELS and Dias JCT (2016) Effects of phosphate-solubilizing bacteria, native microorganisms and rock dust on *Jatropha curcas* L. growth, *Gene Molec Res*, 15(4).
- Saravanakumar D and Samiyappan R (2006) ACC deaminase from *Pseudomonas fluorescens* mediated saline resistance in groundnut (*Arachis hypogea*) plants, *J App Microbiol*, 102: 1283-1287.
- Sardina MG, Boiardi JL and Erbola RJ (1986) Solubilization of phosphorus from low grade minerals by microbial action, *Biotechnology Letters* 8: 247-252.
- Satyaprakash M, Nikitha T, Reddi EUB, Sadhana B and Vani SS (2017) Phosphorus and phosphate solubilizing bacteria and their role in plant nutrition: A review. *Int. J Curr Microbiol App Sci*, 6(4): 2133-2144.
- Saxena J, Saini A, Kushwaha K and Arino A (2016) Synergetic effect of plant growth promoting bacterium *Pseudomonas fluorescens* and Phosphate solubilizing fungus *Aspergillus awamori* for growth enhancement of Chickpea, *Indian Journal of Biochemistry & Biophysics*, 53: 135-143.
- Seema B, Sharma, Riyaz Z, Sayyed, Mrugesh H, Trivedi and Thivakaran A (2013) Phosphate solubilizing microbes: sustainable approach for managing phosphorus deficiency in agricultural soils, *Springer Plus*, 2: 587.
- Selvi KB, Paul JAA, Vijaya V and Saraswathi K (2017) Analyzing the efficacy of phosphate solubilizing microorganisms by enrichment culture techniques, *Biochemistry and Molecular Biology Journal*, 3: 1-3.
- Sharma K, Agrawal A, Bhatnagar M and Sharma R (2015) Effect of phosphate solubilizing bacteria on the germination of Cicer arietinum seeds and seedling growth, *J. Herb. Med. Toxicol.*, 1(1): 61-63.
- Sharma SB, Sayyed RZ, Trivedi MH and Gobi TA (2013) Phosphate solubilizing microbes: sustainable approach for managing phosphorus deficiency in agricultural soils, *Springer Plus*. 2: 587-590.
- Sharon JA, Hathwaik LT, Glenn GM, Imam SH and Lee CC (2016) Isolation of efficient phosphate solubilizing bacteria capable of enhancing tomato plant growth, *J Soil Sci Plant Nutri*, 16 (2): 525-536
- Sharpley AN (1985) Phosphorus cycling in unfertilized and fertilized agricultural soil, *Soil Science Society of America Journal* 49: 905-911.
- Sheraz MS, Hassan GI, Samoon SA, Rather HA, Showkat A, Dar and Zehra B (2010) Bio-fertilizers in organic agriculture, *J. Phytol.*, 2(10): 42-54.
- Singh N and Siddiqui ZA (2015) Effect of *Bacillus subtilis*, *Pseudomonas fluorescens* and *Aspergillus awamori* on the wilt-leaf spot disease complex of Tomato, *Phtopar*, 43: 61-65.
- Singh S and Kapoor KK (1994) Solubilization of insoluble phosphates by bacteria isolated from different sources, *Environmental Ecology*, 12: 51-55.
- Sofia IA, Pereira, Paula ML and Castro (2014) Phosphate solubilizing rhizobacteria enhance Zea



- mays growth in agricultural P-deficient soils, *Ecol. Engi.*, 73: 526–535
- Song OR, Lee SJ, Lee YS, Lee SC, Kim KK and Choi YL (2008) Solubilization of insoluble inorganic phosphate by *Burkholderia cepacia* DA 23 isolated from cultivated soil, *Brazil J. Microbiol.*, 39: 151-156.
- Sundara B, Natarajan V and Hari K (2002) Influence of phosphorus solubilizing bacteria on the changes in soil available phosphorus and sugarcane and sugar yields, *Field Crops Research*, 77(1): 43–49.
- Sundera-Rao WVB and Sinha MK (1963) Phosphate dissolving microorganisms in the soil and rhizosphere, *Indian J Agri Science* 33: 272-275.
- Suranga S and Kumar N (1993) Phosphate solubilization under varying pH by Rhizobium from the legumes. *Journal of Experimental Biology* 31: 855-857.
- Tarafdar JC and Claasson N (1988) Organic phosphorus compounds as a phosphorus source for higher plants through the activity of phosphatase produced by plant roots and microorganisms, *Biology and Fertility of Soils*, 5: 308-312.
- Trivedi P and Sa T (2008) *Pseudomonas corrugate* (NRRL B-30409) mutants increased phosphate solubilisation, organic acid production, and plant growth at lower temperatures, *Curr. Microbiol.*, 56: 140–144.
- Trolove SN, Hedley MJ, Kirk GJD, Bolan NS and Loganathan P (2003) Progress in selected areas of rhizosphere research on P acquisition. *Aust. J. Soil Res.* 41: 471–499.
- Turan M Ataoglu N and Sahin F (2007) Effect of *Bacillus* FS-3 on growth of tomato (*Lycopersicon esculentum* L.) plant and availability of phosphorus in soil, *Plant Soil Environ.*, 53:58-64
- Vessey JK (2003) Plant growth promoting rhizobacteria as biofertilizers, *Plant Soil.* 255: 571–586.
- Vikram A and Hamzehzarghani H (2008) Effect of phosphate solubilizing bacteria on nodulation and growth parameters of green gram (*Vigna radiata* L. W), *Res J Microbiol*, 3: 62-68.
- Villegas J and Fortin JA (2002) Phosphorus solubilization and pH changes as a result of the interactions between soil bacteria and arbuscular mycorrhizal fungi on a medium containing NO<sub>3</sub> as nitrogen source, *Can. J. Bot.*, 80: 571-576.
- Vyas P and Gulati A (2009) Organic acid production in vitro and plant growth promotion in maize under controlled environment by phosphate-solubilizing fluorescent *Pseudomonas*, *BMC Microbiol.*, 9 (1):174-188.
- Walpola BC and Yoon M (2013) Phosphate solubilizing bacteria: Assessment of their effect on growth promotion and phosphorous uptake of mung bean (*Vigna radiata* [L.] R. Wilczek), *Chil. J. Agric. Res.* 73: 275-281.
- Whitelaw MA (2000) Growth promotion of plants inoculated with phosphate solubilizing fungi. *Adv Agron.* 69: 99–151.
- Yazdani M, Mohammad AB, Pirdashti H and Esmaili MA (2009) Effect of Phosphate Solubilization Microorganisms (PSM) and Plant Growth Promoting Rhizobacteria (PGPR) on Yield and Yield Components of Corn (*Zea mays* L.), *World Academy of Science, Engineering and Technology*, 49: 90-92
- Zaidi A and Khan MS (2007) Stimulatory effect of dual inoculations with phosphate solubilizing organisms and an arbuscular mycorrhizal fungus on chickpea, *Austral J Experim Agric.* 47: 1016-1021.
- Zalate PY and Padmani DR (2009) Effect of organic manure and biofertilizers on growth and yield attributing characters of Kharif groundnut (*Arachis hypogaea* L.), *Int J Agric Sci*, 5(2): 343-345.
- Zehra E (2010) Performance of phosphate solubilizing bacteria for improving growth and yield of sunflower (*Helianthus annuus* L.) in the presence of phosphorus fertilizer, *African J. Biotechnol.*, 9(25): 3794-3800.
- Zhang T, Feng H and Lei M (2019) Phosphate-solubilizing bacteria from safflower rhizosphere and their effect on seedling growth, *Open Life Sci.* 14: 246–254.