



Analgesic Properties of Fresh Leaf Juice of *Tridax Procumbens* Linn.

Bele Madhusudan Sudhakarrrao

Department of Zoology, Shri Vasantryao Naik Mahavidyalaya, Dharni, Dist. Amravati-444 702
Email: madhusudansbele@gmail.com

Manuscript details:

Received: 15.09.2019
Accepted: 10.12.2019
Published: 30.12.2019

Cite this article as:

Bele Madhusudan Sudhakarrrao (2019) Analgesic Properties of Fresh Leaf Juice of *Tridax Procumbens* Linn., *Int. J. of Life Sciences*, Volume 7(4): 715-721.

Copyright: © Author, This is an open access article under the terms of the Creative Commons Attribution-Non-Commercial - No Derives License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

Available online on
<http://www.ijlsci.in>
ISSN: 2320-964X (Online)
ISSN: 2320-7817 (Print)

ABSTRACT

Ayurveda is one of the traditional systems of medicine practiced in India and Sri Lanka. Medicines from Ayurveda are largely based upon herbal and herbomineral preparations. Pain is an unpleasant sensation that occurs whenever any tissue is being damaged. Most human diseases have a pain and inflammation components which lead to individuals seek medical attention. The present study was designed to investigate the Analgesic activity of oral administration of fresh leaf juice of *Tridax procumbens* L. in Swiss albino mice (*Mus musculus*). The analgesic activity was determined using acetic acid-induced writhing response. These experimental results suggest that fresh leaf juice of *Tridax procumbens* produced analgesic effect in mice.

Keywords: Analgesic, oedema, *Tridax procumbens*, mice & Ayurveda.

INTRODUCTION

Ayurveda is one of the traditional systems of medicine practiced in India and Sri Lanka and can be traced back to 6000 BC (Charak, Samhita, 1949). Medicines from Ayurveda are largely based upon herbal and herbomineral preparations and have specific diagnostic and therapeutic principles (Patwardhan and Hopper, 1992). The therapeutic efficacies of many indigenous plants, for various diseases have been described by traditional herbal medicine practitioners. Natural products are the source of synthetic and traditional herbal medicines which are still the primary health care system in some parts of the world. In the present investigation we have studied *Tridax procumbens* for its analgesic activity.

Analgesic drugs which are currently in use are either narcotics or non-narcotics which have proven toxic effects and relieve a pain without affecting its causes (Mate *et al.*, 2008). These are associated with adverse side effects on skin, gastrointestinal, renal, hepatic, central nervous system (Simon, 1995; Suleyman *et al.*, 2007). On the contrary, many medicinal plants had been used successfully since long time without any side effects (Ikram, 1983).

In this respect new drugs with improved pain management capacity and fewer side effects are being sought with urgency. Medicinal plants are believed to be an important source of new chemical substances with potential therapeutic effect (Farnsworth, 1989; Eisner, 1990). The research of plants with alleged folkloric use as pain relievers, therefore be viewed as a fruitful and logical research strategy in the search for new analgesic and anti-inflammatory drugs (Elisabetsky *et al.*, 1995).

Tridax procumbens Linn. Commonly known as 'Ekdandi' and Maxican Daisy is a native of tropical America. It also grows wildly in tropical Africa, Asia and Australia and is available in all seasons and in most parts of country (The wealth of India, 1985). It is a perennial herb belonging to the family Asteraceae. Its leaves are used in the treatment of bronchial catarrh, dysentery and diarrhea and for preventing hair loss. The juice of its leaves possesses antiseptic, insecticidal and parasiticidal properties. It possesses a large numbers of chemical constituents which have been identified and isolated in flowers as well as other aerial parts of plants. (Saxena and Albert, 2005, Chantraine *et al.*, 1998).

The people from rural area and the tribal's use fresh leaf of plant as a remedy against several ailments. The leaf juice and the whole plant juice of *Tridax procumbens* is used to treat cuts and wounds (Malhotra and Moorthy, 1973; Sharma *et al.*, 1979; Pandey *et al.*, 1981; Sharma and Malhotra, 1984; Sudhakar and Roll, 1985; Goel and Murgal, 1988), in diarrhoea and dysentery (Sebastian and Bhandari, 1984). It is also used in scorpion bite, stomachache (Jain, 1991) and toothache (Sudhakar and Rolla, 1985). From these uses it seems that it possesses either pain killing properties or antimicrobial properties or both. The bacteriostatic activity of *Tridax procumbens* was demonstrated (Deshmukh *et al.*, 2007). In the present investigations the analgesic activity of *Tridax procumbens* was studied in swiss albino mice.

Pain

Pain is an unpleasant sensation that occurs whenever any tissue is being damaged, and it causes the individual to react to remove the pain stimulus (Guyton and Hall, 2006) to prevent exacerbation of further damage. Therefore it is a protective mechanism for the body. Pain can accompany many disease processes. It is a sign for the brain that

something unwanted happening to the organs or tissue. It could be either a skin injury but it will involve some level of pain (Matosiuk *et al.*, 2007). Currently two main mechanisms of analgesic action are considered as the most useful in pain management.

Peripheral analgesic action

The first involves chain of the prostaglandins transformation with the enzyme Cox-2, specifically involved in the inflammation process, the cyclooxygenase of the second type (Cox-2) (Feldman and McMahan, 2000). Prostaglandins and bradykinin play an important role in the pain process (Hirose *et al.*, 1984). Cyclooxygenase-2 (Cox-2) is an enzyme which is involved in synthesis of prostaglandins from arachidonic acid (Phadke, 1988). Prostaglandins elicit pain by direct stimulation of sensory nerve endings and also sensitize sensory nerve ending to other pain provoking stimuli (Campbell, 1991). Substances inhibiting Cyclooxygenases (Most are acting nonspecifically on Cox-1 and Cox-2) are called Nonsteroidal Anti-inflammatory Drug (NSAID) (Uionne and Gordo, 1994).

Central analgesic action

The second mechanism involves blocking of transduction of the pain stimuli to the brain and backward to cortical cord with the use of the opioid receptors system (Besson, 1999; Curtis *et al.*, 2000). There are three categories of pain receptors or nociceptors.

- Mechanical nociceptors – respond to mechanical damage such as cutting, crushing or pinching.
- Thermal nociceptors – respond to temperature extremes, especially heat.
- Polymodal nociceptors – respond equally to all kinds of damaging stimuli, including irritating chemicals released from injured tissue.

None of the nociceptors have specialized receptor structures, they are all naked nerve endings. All nociceptors can be sensitized by the presence of prostaglandins, which greatly enhance the receptor response to noxious stimuli. The centrally acting analgents are narcotics which inhibits these receptors. Pain is centrally modulated via a number of complex processes including opiate, dopaminergic, descending noradrenergic and serotonergic systems (Bensreti and Sewell, 1983; Terman and Bonica, 2000). Most human diseases have a pain and inflammation components which lead to individuals seek medical attention

(Merskey, 1986). As a consequence analgesic and anti-inflammatory drugs are among the most prescribed drugs in clinical practice (Lim and Yap, 1999).

The flavonoids such as luteolins and quercetin are present in flowers (Ali *et al.*, 2001). The aerial parts (except flowering tops) possess various saturated and unsaturated fatty acids (Gadre and Gabhe, 1988). The plant also possesses phytosterols such as beta-sitosterol, campesterols and stigmasterol, which impart an anti-inflammatory property to it (Gadre and Gabhe, 1993).

Acetic acid induced writhing

Acetic acid induced writhing model of visceral pain (Vyklícky, 1979) is a very sensitive test for analgesic drug development. Writhes can be described as a wave of constriction and elongation passing caudally along the abdominal wall with twisting of the trunk and extension of the hind limbs in mice. This is due to nociceptive property of acetic acid (Surender and Majumdar, 1995).

Acetic acid induced pain sensation by triggering localized inflammatory response. Such pain stimulus leads to release of free arachidonic acid from tissue phospholipids (Ahmed *et al.*, 2006). The acetic acid induced writhing response is a sensitive procedure to evaluate peripherally acting analgesics. The response is thought to be mediated by peritoneal mast cells (Ronaldo *et al.*, 2000), acid sensing ion channels (Voilley, 2004) and the prostaglandin pathways (Koster *et al.*, 1959; Hossain *et al.*, 2006). The results of acetic acid induced writhing test showed highly significant decrease in writhing due to treatment of fresh leaf juice of *Tridax procumbens*. The results were parallel to Voveran treatment. These results indicate that *Tridax procumbens* is a strong analgesic drug and is acting peripherally probably interfering with prostaglandin pathways.

Phytochemical analysis of *Tridax procumbens* showed presence of flavonoids, alkaloids, tannins, steroids (Nia *et al.*, 2003; Agrawal *et al.*, 2009; Salahdeen *et al.*, 2004; and Kale *et al.*, 2008). The observed analgesic activity may be attributed to these compounds (Hasan *et al.*, 2009). The role of tannins in antinociception was reported by Vanu *et al.*, (2006); Salawu *et al.*, (2008). Flavonoids have been found to play an important role in the peripheral analgesic property through inhibition of Cox-2 the enzyme involved in prostaglandin

synthesis (Hossinzadeh *et al.*, 2002, Salawu *et al.*, 2008, Hasan *et al.*, 2009, Zakaria *et al.*, 2007, Gupta *et al.*, 2008). Flavonoids have antioxidant property. They inhibit NO (Olszanecki *et al.*, 2002) and cause vasodilation which is one of the important mechanisms of antinociception (Naseri *et al.*, 2005). These evidences suggest that *Tridax procumbens* is an effective analgesic drug inhibiting pain sensation through peripheral analgesic action

MATERIAL AND METHOD

Plant material

Fresh leaves of *Tridax procumbens* were collected from the garden. The plant was authenticated at the Department of Botany, Govt. Vidarbha Institute of Science and humanities, Amravati by Dr. P.Y. Bhogonkar. The leaves were washed with sterile water and blotted with sterile muslin cloth and leaf juice was squeezed from all leaf parts by hand crushing. The fresh leaf juice of *Tridax procumbens* was used for oral administration.

Animals

The albino mice (*Mus musculus*) were used. The animal were housed, in cages made up of plastic or Galvanized iron, with medium size approximately 290 × 220 × 140 mm and containing bed at the bottom. The husk was changed every day for cleanliness and good health of animals. The food and drinking water supplied *ad libitum* to the animals. The temperature of the rearing room was maintained in the range of 20 to 25°C. The 12 hours of light and 12 hours of dark was maintained in the animal house. The animals were fed on commercially available pelleted (Amrut Mice Feed, Sangli) diet.

Healthy male and female albino mice (*Mus musculus*) of age three months weighing between 25 to 30 gm were used for the study. Experimentation was carried out with due permission of Institutional animal ethics committee of registered animal house of institute registration No. 1060/ac/07/CPCSEA.

Analgesic Study

Acetic acid induced writhing test:-

This test was carried out to evaluate the analgesic activity of fresh leaf juice of *Tridax procumbens* described as per the modified method of Koster *et al.*, 1959 and Santos *et al.*, 1995. For this purpose 0.25ml of 0.75% acetic acid solution (v/v 0.01 ml/10gm) was

injected intraperitoneally in each group. This causes contraction of abdominal muscles, accompanied by stretching of the hind limb, called as writhing and stretching.

- I) Control group: - received 20 µl of 0.9% saline orally.
- II) Experimental group: - received 50µl of fresh leaf juice of *Tridax procumbens*, orally.
- III) Positive control group: - received 20µl (1.25mg/ml) of Voveran dispersible which was dissolved in distilled water, orally.

After 5 minute of administration of acetic acid, the number of writhing and stretching within next 5 min. was recorded. A reduction in the writhing number as compared to the control group was considered as evidence for the presence of analgesic activity, which was expressed as percent inhibition of writhing. The writhing and stretching made after a time intervals of 30 minute were counted for 5 minute each up to 2 hours.

Data were calculated according to the following formula.

$$\% \text{ inhibition} = \frac{W_c - W_t}{W_c} \times 100$$

Wc – Mean no. of writhings (Control)

Wt – Mean no. of writhings (Treated)

RESULTS AND DISCUSSION

Acetic acid induced writhing test:

Experimental group were exhibits the acetic acid-induced writhing responses in mice, indicative of the analgesic activity of fresh leaf juice of *Tridax procumbens* significantly(p<0.001) reduced writhing responses induced by acetic acid when compared with the control group and with Voveran, fresh leaf juice of *Tridax procumbens* showed moderately significant (p<0.01).

Table -: Effect of fresh leaf juice of *Tridax procumbens* on Acetic acid induced Writhing test.

Time (min.)	Control group	Experimental group	Positive control group
5 to 10	40.66 ± 7.23	26.33 ± 3.21*	24.00 ± 2.65**
30 to 35	22.67 ± 6.11	11.00 ± 1.00*	12.00 ± 3.00*
60 to 65	13.66 ± 4.73	3.33 ± 1.15**	5.33 ± 2.08*
90 to 95	7.66 ± 0.58	2.00 ± 1.00***	3.33 ± 2.52*
120 to 125	4.00 ± 1.00	0	1.66 ± 0.58*,\$\$

(Average No. of writhes ± SD)


*=Statistically significant (p< 0.05)

**,\$\$= moderately significant (p<0.01)

***=Highly significant (p<0.001)

*: Significance compared with Control

\$: Significance compared with



Two main mechanisms of the analgesic/antinociceptive drugs are useful in pain treatment. The analgesic drug acting as inhibitors of Cyclooxygenase-2 are the peripherally acting analgesics. For the study of analgesic effect of any drug acetic acid induced writhing test (Koster *et al.*, 1959).

Acetic acid induced writhing

Acetic acid induced writhing model of visceral pain (Vyklícky, 1979) is a very sensitive test for analgesic drug development. Writhes can be described as a wave of constriction and elongation passing caudally along the abdominal wall with twisting of the trunk and extension of the hind limbs in mice. This is due to nociceptive property of acetic acid (Surender and Majumdar, 1995).

Acetic acid induced pain sensation by triggering localized inflammatory response. Such pain stimulus leads to release of free arachidonic acid from tissue phospholipids (Ahmed *et al.*, 2006). The acetic acid induced writhing response is a sensitive procedure to evaluate peripherally acting analgesics. The response is thought to be mediated by peritoneal mast cells (Ronaldo *et al.*, 2000), acid sensing ion channels (Voilley, 2004) and the prostaglandin pathways (Koster *et al.*, 1959; Hossain *et al.*, 2006). The results of acetic acid induced writhing test showed highly significant decrease in writhing due to treatment of fresh leaf juice of *Tridax procumbens*. The results were parallel to Voveran treatment. These results indicate that *Tridax procumbens* is a strong analgesic drug and is acting peripherally probably interfering with prostaglandin pathways.

Phytochemical analysis of *Tridax procumbens* showed presence of flavonoids, alkaloids, tannins, steroids (Nia *et al.*, 2003; Agrawal *et al.*, 2009; Salahdeen *et al.*, 2004; Kale *et al.*, 2008). The observed analgesic activity may be attributed to these compounds (Hasan *et al.*, 2009).

Conflict of Interest

The author declares that there is no conflict of interest.

REFERENCES

Ahmed F, Hossain MH, Rahman AA, Shahid IZ (2006) Antinociceptive and sedative effects of the bark of *Cerbera odollam*_Gaertn. *Oriental Pharmacy and Experimental Medicine*. 6,344-348.

Ali M, Ravinder E, Ranchandram R (2001) A new flavonoid from the aerial parts of *Tridax procumbens*, *Fitoterapia*. 73(3), 313 - 315.

Bensreti MM, Sewell RDE (1983) Selective effects of dopaminergic modifiers on antinociception produced by different opioid receptor agonists. *Pro .Br. Pharmacol. Soc.* 70.

Besson JM (1999) The neurobiology of pain. *Lancet*. 353, 1610-1615.

Campbell WB (1991) Lipid-derived autacoids: Eicosanoids and platelet activating factor, in: Goodman and Gilman's. Gilman A.G. Rall T.W. Neis A.S., Taylor P. (eds) 8th ed., *The pharmacological basis of therapeutics*. New York, Pergamon Press. 607-608.

Chantraine J, Laurent D, Ballivian C, Saavedra G (1998) Insecticidal activity of essential oils on *Aedes aegypti* larvae, *Phytotherapy research*. 12(5), 350 – 354.

Charak Samhita (Trans) (1949) Shree Gulab Kunverba Ayurvedic Society, Jamnagar, India.

Curtis CL, Hughes CE, Flannery CR, Little CB, Harwood JL, Caterson B (2000) n-3 Fatty acids specifically modulate catabolic factors involved in articular cartilage degradation. *Journal of Biological Chemistry*. 275 (2), 721-724.

Deshmukh AA, Gajare KA, Bele MS, Solanke MR, Deshmukh VP (2007) Evaluation of antibacterial property of fresh leaf juice of *Tridax procumbens* an in vitro study. *Journal of cell Tissue Research*. 7 (2), 1123 – 1129.

Eisner T (1990) Chemical prospecting a call for action. In: Borman F. H. Kellert S. R. editor. *Ecology, Economics and Ethics. The Broken circle*, Yale University Press, New Haven CT.

Elisabetsky E, Amodor TA, Albuquerque RR, Nunes DS, Carvalho ACT (1995). Analgesic activity of *Psychotria colorata* (Wild ex R. and S.) Triuelli Arg. *Alkaloids. Journal of Ethnopharmacol.* 48, 77-83.

Farnsworth NR (1989) Screening plants for new medicines. In: Wilson, E. O., Editor *Biodiversity, Part II*. National Academy Press, Washington, pp. 83-87.

Fasmer OB, Berge OG, Hole K (1985) Changes in nociception after lesions of descending serotonergic pathways induced with 5, 6 - dihydroxytryptamine. Different effects in the formalin and tail flick test. *Neuropharmacology*. 24, 729- 734.

Feldman M, McMahan AT (2000) Do cyclooxygenase-2 inhibitors provide benefits similar to those of traditional nonsteroidal anti-inflammatory drug, with less gastrointestinal toxicity? *Annals of Internal Medicine*. 132, 134-143.

Gadre AP, Gabhe SY (1988) Saturated and unsaturated fatty acids from *Tridax procumbens*. *Indian Journal of Pharmaceutical Sciences*. 50 (3), 168.

Gadre AP, Gabhe SY (1993) Identification of some sterols of *Tridax procumbens* by GC – MS, Indian. *Journal of Pharmaceutical Sciences*. 55 (5), 191 – 192.

- Goel AK, Murgal V (1988) A survey of medicinal plants used by the tribal's of Santal Pargana (Bihar). *Journal of economic and taxonomic botany*. 12, 329 – 335.
- Gupta M, Shaw BP, Mukerjee A (2008) Studies on Antipyretic-Analgesic and Ulcerogenic activity of polyherbal preparation in Rats and mice. *International Journal of Pharmacology*. 4(2), 88-94.
- Guyton AC, Hall JE (2006) Somatic Sensation: pain, headache and thermal sensation. In: Textbook of Medical physiology. 11th ed., Bangalore, India, Prism Book. 598-599.
- Hasan SMR, Jamila M, Majumdar MM, Akter R, Hossain MM, Mazumder MEH, Alam MA, Jahangir R, Rana MS, Arif M and Rahman S (2009) Analgesic and antioxidant activity of the Hydromethanolic Extract of *Mikania scandens* (L.) Wild. Leaves. *American Journal of Pharmacology & Toxicology*. 4(1), 1-7.
- Hirose K, Jyoyama H, Kojima Y, Eigyo M, Hatakeyama H, Asanuma F, Umehara H, Yamaguchi T (1984) Pharmacological properties of 2-[4-(2-triazolyloxy)-Phenyl]-propionic acid (480156-S), a new non-steroidal anti-inflammatory agent. *Arzneimittelforschung*. 34, 280-286.
- Hossain MM, Ali MS, Saha A, Alimuzzaman M (2006) Antinociceptive activity of whole plant extract of *Paederia foetida*. Dhaka University. *Journal of Pharmaceutical Science*. 5, 67-69.
- Ikram M (1983) *Hamdard Medicus*. 26, 16-17.
- Jain SK (1991) Ethanobotanical uses of plants, in: Dictionary of India folk medicine and ethnobotany (Jain S.K. Ed) Deep publications, New Delhi India.
- Kale MA, Shahi SR, Somani VG, Shamkuwar PB, Dhake AS, (2008) Hemostatic activity of the leaves of *Tridax procumbens* Linn. *International Journal of Green pharmacy*. 1(2), 54-55.
- Koster R, Andereson M, De Berar EJ (1959) Acetic acid for analgesic screening. *Federation Proceedings*. 18, 412-416.
- Lim KHJ, Yap KB (1999) The prescription pattern of outpatient polyclinic doctors. *Singapore Medical Journal*. 40, 416- 419.
- Malhotra SK, Moorthy S (1973) Some useful and medicinal plant of Chandrapur district (Maharashtra State) *Bull. Botanical Survey of India*. 15, 13 – 21.
- Mate GS, Naikwade NS, Chowki CSAA, Patil SB (2008) Evaluation of anti-nociceptive activity of *Cissus quadrangularis* on albino mice. *International Journal of Green pharmacy*. 2, 118-121.
- Merskey HM (1986) Pain terms. *Pain*. 3, S215-S221.
- Naseri MKGHamidi MN, Heidari A (2005) Vasorelaxatory effect of *Vitis vinifera* extract on rat aorta. *Iran Journal of Pharmaceutical Science*. 2, 93-99.
- Nia R, Paper DH, Essien EE, Oladimeji OH, Lyadi KC, Franz G, (2003) Investigation into in-vitro radical scavenging and in-vivo anti - inflammatory potential of *Tridax procumbens*. *Nigerian Journal of Physiological Sciences*. 18(1-2), 39-43.
- Olszanecki, R., Gebaska, A., Kozlovski, V.I., Gryglewski, R.J., 2002. Flavonoid and nitric oxide synthase. *Journal of Physiology and Pharmacology*. 53,571-584.
- Pandey G, Singh VK, Bhatnagar IS (1981) New records to the medical efficacy claims of certain plant recorded from Gwalior forest circle M. P. – A preliminary contribution. *Bull. Medico-ethnobotany Res*. 2,303-314.
- Patwardhan B, Hopper B (1992) Ayurvedic and Future drug development. *Journal of Alternative and complementary Medicine*. 19, 9-10.
- Phadke K (1988) *In vivo* and *in vitro* models for arthritis. *Indian Drugs*. 25 (9), 354-365.
- Ronaldo AR, Mariana LV, Sara MT, Adriana BPP, Steve P, Ferreira SH, Fernando QC (2000) Involvement of resident macrophages and mast cells in the writhing nociceptive response induced by zymosan and acetic acid in mice. *European Journal of Pharmacology*. 387, 111-118.
- Salahdeen HM, Yemitan OK, Alada ARA (2004) Effect of aqueous leaf extract of *Tridax procumbens* on blood pressure and heart rate in rats. *African Journal of Biomedical Research*.7(1), 27-29.
- Salawu OA, Chindo BA, Tijani AY, Adzu B (2008) Analgesic, anti-inflammatory, antipyretic and anti-plasmodial effects of the methanolic extract of *Crossopteryx febrifuga*. *Journal of Medicinal plant Research*. 2(8), 213-218.
- Santos ARS, Valdir CF, Rosendo AY, Joao BC (1995) Further studies on the antinociceptive action of the hydroalcoholic extracts from plants of the genus *Phyllanthus*. *Journal of Pharmacology and Experimental Therapeutics*. 183, 226 – 234.
- Saxena VK, Albert S (2005) β - Sitosterol-3-O- β -D-xylopyranoside from the flowers of *Tridax procumbens* Linn. *Journal of Chemical Science*. 117 (3), 263 – 266.
- Sebastian MK, Bhandari MM (1984) Medico ethnobotany of Mount Abu, Rajasthan, India. *Journal of Ethnopharmacol*. 12, 223-230.
- Sharma BD, Malhotra SK (1984) A contribution to ethnobotany of tribal areas in Maharashtra. *Journal of Economic and Taxonomic Botany*. 5, 533-537.
- Sharma PK, Dhyani SK, Shankar V (1979) Some useful and medicinal plant of district Dehradun and Siwalik. *J. Sci. Res. Pl. Med*. 1, 17-43.
- Simon LS (1995) Action and toxicity of non-steroidal anti-inflammatory drug. *Current Opinion in Rheumatology*. 7(3), 159-166.
- Sudhakar S, Rolla S (1985) Medical plant of upper East Godavari District (Andhra Pradesh) and need for establishment of medical farm. *Journal of Economic and Taxonomic Botany*. 7, 399- 406.
- Suleyman H, Demircan B, Karagoz Y (2007) Anti-inflammatory and side effect of cyclooxygenase inhibitors. *Pharmacological Report*. 59(3), 247-258.
- Surender S, Majumdar DK (1995) Analgesic activity of fixed oil of *Ocimum sanctum* linn. (Tulsi) and its possible

mechanism of action. *International Journal of Pharmacognosy*. 33(3), 188-192.

Terman GW, Bonica JJ (2000) Spinal mechanisms and their modulations, in: Loeser, J.D., Butler, S.H., Chapman, C.R. (eds), *Bonica's Management of pain*. Lippincott Williams and Wilkins, Philadelphia.73-152.

The Wealth of India (1985) A dictionary of Indian raw materials and industrial products, Publications and Information Directorate. Council of Scientific and Industrial Research, New Delhi. 10 (sp-w), 292-295.

Uionne RA, Gordo SM (1994) Non-steroidal anti-inflammatory drugs for acute pain control. *Dental Clinician North America*. 38, 645 - 664.

Vanu MR, Palanivelu S, Panchanatham S (2006) Immunomodulatory and anti-inflammatory effects of *Semecarpus anacardium*_Linn. Nut milk extract in experimental inflammatory conditions. *Biological and Pharmaceutical Bulletin*. 29, 693-700.

Voilley N (2004) Acid sensing Ion channels (ASICs): New targets for the analgesic effects of Non-Steroid Anti-inflammatory Drugs (NSAIDs) current Drug Targets. *Inflammation and Allergy*. 3, 71-79.

Vyklicky L (1979) Techniques for the study of pain in animals, in: Bonica, J.J., Liebeskind, J.C., Albe-Fessard, D.G., (Eds). *Advances in pain research and therapy*. Raven press. New York.

Zakaria ZA, Wen LY, Rahman MIA, Ayub AHA, Sulaiman MR, Gopalan HK (2007) Antinociceptive, Anti-inflammatory and Antipyretic properties of the aqueous extract of *Bauhinia purpurea* leaves in experimental animals. *Medical Principles and Practice*. 16, 443-449.

© 2013 -2019 | Published by IJLSCI