

## Impact Factor:

ISRA (India) = 4.971	SIS (USA) = 0.912	ICV (Poland) = 6.630
ISI (Dubai, UAE) = 0.829	PIHII (Russia) = 0.126	PIF (India) = 1.940
GIF (Australia) = 0.564	ESJI (KZ) = 8.997	IBI (India) = 4.260
JIF = 1.500	SJIF (Morocco) = 5.667	OAJI (USA) = 0.350

SOI: [1.1/TAS](#) DOI: [10.15863/TAS](#)

## International Scientific Journal Theoretical & Applied Science

p-ISSN: 2308-4944 (print) e-ISSN: 2409-0085 (online)

Year: 2020 Issue: 07 Volume: 87

Published: 23.07.2020 <http://T-Science.org>

QR – Issue



QR – Article



**Sabeeh Akbar**

Bahawal Victoria Hospital Bahawalpur  
Ex house officer

[sabeehakbar@hotmail.com](mailto:sabeehakbar@hotmail.com)

**Benazir Shafqat**

Bahawal Victoria Hospital Bahawalpur  
Ex house officer

[benazirsh@yahoo.com](mailto:benazirsh@yahoo.com)

**Bakhtawar Abbasi**

Bahawal Victoria Hospital Bahawalpur  
Ex house officer

[beckya13@hotmail.com](mailto:beckya13@hotmail.com)

## MORTALITY OF ACUTE INFERIOR WALL MYOCARDIAL INFARCTION WITH AND WITHOUT ATRIOVENTRICULAR BLOCK

**Abstract: Objective:** To determine mortality rate associated with inferior wall myocardial infarction with or without atrioventricular block.

**Design & duration:** This is a cohort study completed in duration of six months.

**Setting:** Study was conducted in Cardiology ward of Bahawal Victoria Hospital Bahawalpur

**Patients & methods:** Patients presenting to the study hospital with inferior wall myocardial infarction combined with or without atrioventricular block. Inclusion and exclusion criteria were made for selection of patients. Sample size calculated using WHO sample size calculator. Non-probability consecutive sampling technique was used. Level of significance was 5%. P-value less than 0.05 was considered statistically significant. Consent was taken from all cases for including their data in the study. Permission was also taken from ethical review board committee. Chi square test was applied. SPSS software version 20 was used for analyzing data. Results were calculated in the form of percentage, frequency, means and standard deviation.

**Results:** Total 160 cases were studied. They were divided into two groups. Patients with inferior wall MI having atrioventricular block as well were placed in group-A. Other patients with inferior wall MI without AV block were placed in group-B. Mean age was 53±7.8 years. There were 43.3% male and 57.6% female patients. Majority of cases (66.7%) were above 50 years of age. Mortality in group-A was 18.3% and in group-B 11.7%.

**Conclusion:** Inferior wall myocardial infarction complicated with atrio-ventricular block has high mortality rate as compared to MI without AV block.

**Key words:** Myocardial Infarction, Atrioventricular block, Mortality.

**Language:** English

**Citation:** Akbar, S., Shafqat, B., & Abbasi, B. (2020). Mortality of acute inferior wall myocardial infarction with and without atrioventricular block. *ISJ Theoretical & Applied Science*, 07 (87), 155-159.

**Soi:** <http://s-o-i.org/1.1/TAS-07-87-36> **Doi:** [crossref https://dx.doi.org/10.15863/TAS.2020.07.87.36](https://dx.doi.org/10.15863/TAS.2020.07.87.36)

**Scopus ASCC:** 2700.

### Introduction

Decrease in blood flow in coronary arteries causes acute coronary syndrome. Despite advanced management and developed medical field still acute

coronary syndrome is leading cause of mortality. Annually one million people die in Pakistan due to ACS. It is because of increasing prevalence of risk factors of ischemic heart disease. Thrombotic

## Impact Factor:

ISRA (India)	= 4.971	SIS (USA)	= 0.912	ICV (Poland)	= 6.630
ISI (Dubai, UAE)	= 0.829	PIHII (Russia)	= 0.126	PIF (India)	= 1.940
GIF (Australia)	= 0.564	ESJI (KZ)	= 8.997	IBI (India)	= 4.260
JIF	= 1.500	SJIF (Morocco)	= 5.667	OAJI (USA)	= 0.350

occlusion of coronary arteries results in ST-segment elevation in MI, which is main sign seen on ECG in such patients. Complications of MI are ischemia, arrhythmia and thromboembolic changes. Conduction defect in AV block along with MI is associated with high mortality rate. Usually observed conduction defects after MI are Atrioventricular nodal block of 1<sup>st</sup>, 2<sup>nd</sup> and 3<sup>rd</sup> degree. Complete AV block complicates inferior wall MI in 11-15% cases. It occurs in conditions like atrial fibrillation, cardiogenic shock and ventricular infarction. Previously mortality of MI with AV block has not been studied. This study has been conducted to evaluate prognosis of MI with AV block. This will help in early recognition of disease and to treat it immediately, hence mortality rate can be reduced.

### PATIENTS AND METHODS

This is a cohort study started in January and completed after six months in June 2020. Study was conducted in Cardiac department of Bahawal Victoria Hospital Bahawalpur. Patients presenting to the study hospital with inferior wall myocardial infarction combined with or without atrioventricular block. Inclusion and exclusion criteria were made for selection of patients. Sample size calculated using WHO sample size calculator. Non-probability consecutive sampling technique was used. Level of significance was 5%. P-value less than 0.05 was considered statistically significant. Consent was taken from all cases for including their data in the study. Permission was also taken from ethical review board committee. Chi square test was applied. SPSS software version 20 was used for analyzing data.

Results were calculated in the form of percentage, frequency, means and standard deviation.

**Sampling Technique:** Non-probability consecutive sampling

#### Inclusion Criteria

Age 20-70 years

Either sex

Patient presenting in 12 hours of signs symptoms

Inferior wall MI with AV block, and without AV block

#### Exclusion Criteria

Clinical features of cardiogenic shock


Mechanical features of MI developing in patients like ventricular septal rupture, acute MR, and ventricular wall rupture.

Patients not thrombolysed

Patients with co-morbidities like renal failure or liver failure etc.

### RESULTS


Total 160 cases were studied. They were divided into two groups. Patients with inferior wall MI having atrioventricular block as well were placed in group-A. Other patients with inferior wall MI without AV block were placed in group-B. Mean age was 53±7.8 years. There were 52(43.3%) male and 68(57.6%) female patients. Majority of cases 80(66.7%) were above 50 years of age. Mortality in group-A was 11(18.3%) and in group-B 7(11.7%). There were total 40 cases between 20-50 years of age including 22 in group-A and 18 from group-B. Out of 22 from group-A 5(22.7%) died and out of 18 from group-B 3(16.7%) died. Similarly 6(15.8%) out of 38(63.3%) cases from group-A and 4(9.5%) out of 42(70%) cases from group-B died.



(Figure-1) Group wise distribution of mortality rate

**Impact Factor:**

<b>ISRA (India)</b> = 4.971	<b>SIS (USA)</b> = 0.912	<b>ICV (Poland)</b> = 6.630
<b>ISI (Dubai, UAE)</b> = 0.829	<b>ПИИИ (Russia)</b> = 0.126	<b>PIF (India)</b> = 1.940
<b>GIF (Australia)</b> = 0.564	<b>ESJI (KZ)</b> = 8.997	<b>IBI (India)</b> = 4.260
<b>JIF</b> = 1.500	<b>SJIF (Morocco)</b> = 5.667	<b>OAJI (USA)</b> = 0.350



**(Figure-2) Age wise distribution of mortality rate**

**(Table-1) Characteristics of patients in study group. (n=120)**

Characteristics	Group-A (n=60)	Group-B (n=60)
<b>Age</b>		
<50 years	22(36.7%)	18(30%)
>50 years	38(63.3%)	42(70%)
<b>Gender</b>		
Male	28 (46.7%)	24(40%)
Female	32 (53.3%)	36(60%)
<b>Mortality</b>		
Yes	11(18.3%)	7(11.7%)
No	49(81.7%)	53(88.3%)

**DISCUSSION**

Now a day medical field has been much developed. Modern techniques and cardiac interventions have reduced mortality rate to much extent. PCI is now being practiced in developing countries as well, reducing complications related o MI. AV conduction block is a complication of MI. Mortality associated with AV after MI in hospital admitted patients in Pakistan has never been studied before. Purpose of this study is to determine mortality rate associated with MI and AV block. Total 160 cases were studied. They were divided into two groups. Patients with inferior wall MI having atrioventricular block as well were placed in group-A. Other patients with inferior wall MI without AV block were placed in group-B. Mean age was 53±7.8 years. There were 52(43.3%) male and 68(57.6%) female patients. Majority of cases 80(66.7%) were above 50 years of age. Mortality in group-A was 11(18.3%) and in

group-B 7(11.7%). There were total 40 cases between 20-50 years of age including 22 in group-A and 18 from group-B. This is a cohort study started in January and completed after six months in June 2020. Study was conducted in Cardiac department of Bahawal Victoria Hospital Bahawalpur. Patients presenting to the study hospital with inferior wall myocardial infarction combined with or without atrioventricular block. Inclusion and exclusion criteria were made for selection of patients. Sample size calculated using WHO sample size calculator. Non-probability consecutive sampling technique was used. Level of significance was 5%. P-value less than 0.05 were considered statistically significant. How AV block increases mortality rate has not been studied still. AV block are associated with large ischemic areas of heart resulting in heart failure and increasing mortality rate.

## Impact Factor:

ISRA (India) = 4.971  
ISI (Dubai, UAE) = 0.829  
GIF (Australia) = 0.564  
JIF = 1.500

SIS (USA) = 0.912  
PIIHQ (Russia) = 0.126  
ESJI (KZ) = 8.997  
SJIF (Morocco) = 5.667

ICV (Poland) = 6.630  
PIF (India) = 1.940  
IBI (India) = 4.260  
OAJI (USA) = 0.350

## CONCLUSION

Inferior wall myocardial infarction complicated with atrio-ventricular block has high mortality rate as compared to MI without AV block. Early diagnosis

and prompt treatment can prevent AV block and can reduce mortality rate.

## References:

1. Schoen, F.J. (2010). *The heart*. In Kumar V, Abbas AK, Fausto N, Aster J, editors. Robbins and Cotran pathologic basis of disease. 8th ed. (pp.529-587). Philadelphia: WB Saunders.
2. Seghieri, C., Mimmi, S., Lenzi, J., & Fantini, M.P. (2012). 30-day in-hospital mortality after acute myocardial infarction in Tuscany (Italy): an observational study using hospital discharge data. *BMC Med Res Method*, 12:170.
3. Russell, L.B., Gold, M.R., Siegel, J.E., Daniels, N., & Weinstein M.C. (1996). The role of cost effectiveness analysis in health and medicine: panel on cost-effectiveness in health and medicine. *JAMA*, 1996; 276: 1172-7.
4. Gaziano, J.M. (2005). *Global burden of cardiovascular disease*. In: Zipes DP, Libby P, Bonow RO, Braunwald E, editors. Braunwald's heart disease: a textbook of cardiovascular medicine. 7th ed. (pp.1-19). Philadelphia: Saunders.
5. Weinstein, M.C., Siegel, J.E., Gold, M.R., Kamlet, M.S., & Russell, L.B. (1996). Recommendations of the panel on cost-effectiveness in health and medicine. *JAMA*. 1996; 276:1253-8.
6. Wang, S.S., Paynter, L., Kelly, R.V., Koch, G.G., Skains, M.S., & Gettes, L.S. (2009). Electrocardiographic determination of culprit lesion site in patients with acute coronary events. *J Electrocardiol*, 42:46-51.
7. Kakouros, N., & Cokkinos, D.V. (2010). Right ventricular myocardial infarction: pathophysiology, diagnosis, and management. *Postgrad Med J*. 2010; 86:719-28.
8. Surawicz, B., Knilans, T., & Chou, S. (2008). *Electrocardiography in clinical practice: adult and pediatric*. 6th ed. Philadelphia: WB Saunders.
9. Kakouros, N., Kakouros, S., Lekakis, J., Rizors, I., & Cokkinos, D. (2011). Tissue Doppler imaging of the ENT in the acute and late phase of a first tricuspid annulus and myocardial performance index in the evaluation of right ventricular involvement in inferior myocardial infarction. *Echocardiography*. 2011; 28:311-9.
10. Hamon, M., Agostini, D., Le Page, O., & Riddell, J.W. (2008). Prognostic impact of right ventricular involvement in patients with acute myocardial infarction: Meta analysis. *Crit Care Med*. 2008; 36:2023-33.
11. Saleheen, D., & Fossard, P. (2004). CAD risk factors and acute myocardial infarction in Pakistan. *Acta Cardiol*. 2004; 59:417-24.
12. Boden, W.E., & O'rouke, R.A. (2007). CORAGE trial group. The evolving pattern of coronary artery disease in the US and Canada: Baseline characteristics of the Clinical Outcomes Utilizing Revascularization and Aggressive Drug Evaluation (COURAGE) trial. *Am J Cardiol*. 2007; 99:208-12.
13. Becker, R.C., et al. (1994). Comparison of clinical outcomes for women and men after acute myocardial infarction. *Ann Intern Med*. 1994; 120:638-45.
14. Tofler, G.H., et al. (1987). Effect of gender and race on prognosis after myocardial infarction: adverse prognosis for women, particularly black women. *J Am Coll Cardiol*. 1987; 9:473-82.
15. Honjo, K., et al. (2010). The effects of smoking and smoking cessation on mortality from cardiovascular disease amongst Japanese: pooled analysis of three large-scale cohort studies in Japan. *Tob Control*. 2010; 19: 50-7.
16. (2010). International Diabetes Federation. The diabetes atlas. 4th ed. Brussels: The Federation.
17. Ding, C.H., Teng, C.L., & Koh, C.N. (2006). Knowledge of diabetes mellitus among diabetic and non-diabetic patients. *Med J Malaysia*, 2006; 61:1134-8.
18. Barrentt-Connor, E., Ciriqui, M.H., Klauber, M.R., Hold brook, M. (1981). Diabetes and hypertension in a community of older adults. *Am J Epidemiol.*, 1981; 113: 276-84.
19. Stamler, J., et al. (1993). Diabetes other risk factors and 12-yr cardiovascular mortality for men screened in the Multiple Risk Factor Intervention Trial. *Diabetes Care*. 1993; 16:
20. Fresco, C., et al. (1996). "Prognostic value of a history of hypertension in 11,483 patients with acute myocardial infarction treated with thrombolysis. GISSI-2 Investigators. Gruppo Italiano per lo Studio della Sopravvivenza nell'Infarto Miocardico," *J Hypertens*. 1996; 14:743-50.

<b>Impact Factor:</b>	<b>ISRA (India) = 4.971</b>	<b>SIS (USA) = 0.912</b>	<b>ICV (Poland) = 6.630</b>
	<b>ISI (Dubai, UAE) = 0.829</b>	<b>ПИИИ (Russia) = 0.126</b>	<b>PIF (India) = 1.940</b>
	<b>GIF (Australia) = 0.564</b>	<b>ESJI (KZ) = 8.997</b>	<b>IBI (India) = 4.260</b>
	<b>JIF = 1.500</b>	<b>SJIF (Morocco) = 5.667</b>	<b>OAJI (USA) = 0.350</b>

---

21. (1993). An international randomized trial comparing four thrombolytic strategies for acute myocardial infarction. The GUSTO investigators. *N Engl J Med.* 1993; 329:673-82.
22. Zehender, M., Kesper, W., & Kauuder, E. (1993). Right ventricle infarction as an independent predictor of prognosis after acute inferior myocardial infarction. *N Engl J Med.* 1993; 328:981-8.
23. Khan, S., Kundi, A., & Sharieff, S. (2004). Prevalence of right ventricular myocardial infarction in patients with acute inferior wall myocardial infarction. *Int J Clin Pract.* 2004; 58:354-7.
24. Iqbal, M.A., Shah, I., Rauf, M.A., Khan, N., Khan, S.B., & Hafizullah, M. (2012). Frequency of acute right ventricular myocardial infarction in patients with acute inferior myocardial infarction. *Pak Heart J.* 2012; 45:81-5.