

DOI: 10.5281/zenodo.3685665  
UDC: 615.814.1:[616.8-009.7+616.833]



## Acupoint embedding therapy

<sup>1,2</sup>Olga Ignatov, <sup>2</sup>Oleg Pascal, <sup>1</sup>Viorel Nacu

<sup>1</sup>Laboratory of Tissue Engineering and Cell Cultures, <sup>2</sup>Department of Rehabilitation  
*Nicolae Testemitsanu* State University of Medicine and Pharmacy, Chisinau, the Republic of Moldova

Authors' ORCID iDs, academic degrees and contributions are available at the end of the article

\*Corresponding author: olga.ignatov@usmf.md

Manuscript received January 18, 2020; revised manuscript February 27, 2019; published online March 10, 2020

### Abstract

**Background:** Peripheral nerve trauma remains a major cause of motor disability, at the same time functional restoration after treatment continues to show modest results. Acupoint embedding therapy is a type of acupuncture treatment in which different biodegradable materials are inserted into specific points for long-term stimulation. It has a good analgesic effect in chronic pain, and it is considered a cure for many diseases. Different biodegradable materials have been developed and widely used. Catgut has a good biodegradability and low price, but it could cause infections and having unstable chemical properties had been limited in clinical use. Such synthetic materials as polylactic acid and polyglycolic acid present low-cost, good biodegradability and biocompatibility compared with the catgut. However, their poor hydrophilicity and cell adhesion limited their therapeutic efficacy. The ideal embedding materials are required to be safe, non-toxic, biocompatible, and to have excellent swelling and biodegradation behaviors. Acupoint embedding therapy can be a promising treatment method of peripheral nerve disorders.

**Conclusions:** Acupoint embedding therapy is an invasive treatment which can prolong point stimulation, reduces the frequencies of pain and psychological fear of patients. It seems to be a promising method of neuropathy treatment. The properties of the filaments for acupoint embedding therapy can be improved by surface modification technologies.

**Key words:** acupuncture embedding therapy, neuropathy.

### Cite this article

Ignatov O, Pascal O, Nacu V. Acupoint embedding therapy. *Mold Med J.* 2020;63(1):52-58. doi: 10.5281/zenodo.3685665.

### Introduction

Acupuncture is generally held to have originated in China, being first mentioned in documents dating from a few hundred years leading up to the Common Era. Sharpened stones and bones that date from about 6000 BC have been interpreted as instruments for acupuncture treatment [1, 2].

The first medical description of acupuncture by a European physician was by Ten Rhijne, in about 1680, who worked for the East India Company and witnessed acupuncture practice in Japan [3, 4].

In 1960s, there appeared a treatment method by implanting absorbable materials (e.g. catgut) substituting for filiform needles into the acupoints, which realized long-time needle retaining and also avoided the danger of filiform needle retaining. This method was then termed acupoint thread-embedding therapy. During recent years, by constant improvement of embedding apparatus and materials, this method has gradually evolved to be micro-invasive thread-embedding therapy [5].

**Traditional acupuncture.** Acupuncture is a medical intervention in which fine needles are applied to specific parts of the body, called acupuncture points (or acupoints) and penetrated through the muscular or other subcutaneous layers. According to the theory of traditional Chinese medicine, acupuncture modulates the flow of Qi and blood through the meridians and restores the balance of the five

organs to maintain homeostasis [6]. Each acupoint has its own specific therapeutic actions. They are presumed to be pathophysiologically associated with and possibly reflect the status of visceral organs and systemic conditions, and thus the stimulation of specific acupoints may evoke the responsiveness that controls the unbalanced internal milieu and improves body symptoms. Acupuncture stimulation is given right on the acupoint or a nearby affected area “*ashi point*” for the treatment of local symptoms, whereas distal acupuncture stimulation is applied to treat diseases in the internal organs and systemic abnormalities [7].

In manual acupuncture, an acupuncturist penetrates the skin with a metallic needle and manipulates it by rotating in one or both directions or lifting and thrusting [8]. Between 5 and 15 needles are used in a typical treatment, with the point combinations varying during a course of sessions [9]. It is known that during acupuncture practice, acupuncturists experience a special touch sensation perceived as heaviness, tenseness, or terseness, and patients perceive feelings of numbness, heaviness, soreness, and distention around the site of needle stimulation. These are called *deqi* sensations. Clinical data further indicate that patients frequently feel *deqi* sensations spreading to other parts of the body [10, 11].

In electric acupuncture, a small electric current is applied to pairs of acupuncture needles [8]. To maximize therapeutic effects, acupuncture is usually practiced first by applying

manual acupuncture to evoke *deqi* sensation and followed by electrical stimulation for 15–20 minutes [10]. Traditional acupuncture therapy has the advantages of safety, validity and non-toxic side effects, but with and unexpected short function time and frequent operation [12].

Among acuapunctures, there mainly are filiform needles, electric needles, hydro-needles (small dose point injection), fire needles, warm needles, skin needles (including plum-blossom needles), ear needles, prick blood-letting therapy and many others [13].

According to the World Health Organization, acupuncture has been shown to be an effective alternative or complementary treatment to 28 diseases, symptoms, or conditions [14].

Peripheral neuropathy is broadly defined as damage of the peripheral nervous system caused by a primary lesion or dysfunction [15]. The recovery of peripheral nerve injury is a long and slow process. This may be because of the period of time needed for neural regeneration and functional reconstruction, but treatment methods may also contribute to the delay in recovery [16, 17]. Peripheral neuropathy, particularly when it involves the large nerve fibers, is ideally suited for a localized structural needling approach. Needles placed in close proximity to a nerve stimulate the nerve fibers directly, with an electric current or a physical stimulus from manual acupuncture [18].

He et al. note that nerve injuries affect the metabolic microenvironment. Citing an example, they note that sciatic nerve injuries reduce acetylcholinesterase activity in the lumbar spinal cord microenvironment. This causes neuronal cell death thereby impeding nerve repair. The researchers note that acupuncture counteracts this effect citing that it successfully increases “acetylcholinesterase expression in spinal cord tissue after peripheral nerve injury”. As a result, this may be an important mechanism by which acupuncture promotes the healing of peripheral nerves [19].

Lu et al. revealed that acupuncture and electroacupuncture could accelerate the maturity of regenerated nerves with larger mean values of axon number, endoneurial area, blood vessel number, and blood vessel area as compared with the controls [20].

Acupuncture is an established adjuvant analgesic modality for the treatment of chronic pain, and it is considered a cure for many ailments and disorders [21]. It is thought to stimulate inhibitory nerve fibers for a short period, reducing transmission of pain signal to the brain [22]. Acupuncture treatment activates endogenous analgesic mechanisms [23], causing secretion of endorphin which is an endogenous opioid [24] and triggering release of adenosine [25], producing a rapidly effective analgesic action. Extensive research has shown that acupuncture analgesia may be initiated by stimulation of high-threshold, small-diameter nerves in the muscles [26].

Han et al. found that acupuncture can promote release of neurotransmitter such as 5-hydroxy tryptamine and in addition it generates neuropeptide through electrical stimulation of different frequencies that has significant effect to

pain reduction [27]. Park et al. speculate that acupuncture stimulation may trigger the responsiveness of sensory receptors and generate neural activity in its own specific way, which may be encoded in the cerebral cortex and autonomic neuronal center and exert its effects on regulating inflammation. Future investigations to explore whether acupuncture-specific vagal activity exists and acts on cells in target organs will be of great importance to gain insights into the mechanistic basis of acupuncture [7].

Acupuncture may elicit vegetative reflexes, thereby changing the flow of blood and enhancing functional properties of connective tissue and organs [28]. Litscher et al. showed that acupuncture may increase blood flow in the limbs [29]. Increased blood flow to the vasa nervorum and dependant capillary beds supplying the neurons [30] may therefore contribute to the immediate effect of acupuncture [28]. Over some time, these may contribute to nerve repair with measurable improvement of axons or myelin sheaths after 10 treatments. Local and central effects on vascularization may thus represent combined causes for regeneration [31, 32]. Acupuncture could regulate multiple molecules and signaling pathways that lead to excitotoxicity, oxidative stress, inflammation, and neurons death and survival and also promote neurogenesis, angiogenesis, and neuroplasticity after ischemic damage [33].

Risk factors for complications of acupuncture:

- For pregnant women, avoid needling points on the abdomen and lumbar region, and certain points known to cause strong sensations;
- Hemophilia can affect clotting factors;
- Advanced liver disease could compromise production of clotting factors;
- Patients taking blood thinners could bleed for longer periods;
- Patients with HIV infection or immunocompromised patients are at increased risk of opportunistic infections;
- Patients with diabetes are subject to poor wound healing; neuropathy can reduce sensory ability, leaving them at increased risk of undetected infection;
- Patients who have had transplants often take immune suppressants that make them prone to infections;
- High-dose steroids suppress the immune system;
- Open wounds increase risk of infection;
- Hypoglycemic, nervous, or very fatigued patients might faint.

Potential adverse events associated with acupuncture are: fainting during treatment, nausea and vomiting, increased pain, diarrhea, local skin irritation, headaches, sweating, dizziness, aggravation of symptoms, needle breakage.

Rare complications include: pneumothorax, spinal cord injury, septicemia, punctured organs, convulsions, argyria [34].

The most adverse effects associated with acupuncture are minor and serious complications are rare. Patients should be forewarned of potential common, though minor, adverse effects. Needle penetration might cause pain and bleeding. Reports of fainting remind practitioners that patients might

be better off prone than sitting during treatments. Practitioners should keep some potentially fatal effects in mind when patients report shortness of breath, pleuritic chest pain, or fever after acupuncture treatment. All acupuncture practitioners should have proper training in techniques and safety [34].

#### Acupoint embedding therapy

In traditional Chinese medicine, catgut-embedding therapy has been used for the treatment of several diseases such as musculoskeletal pain, obesity, chronic urticaria, perimenopausal syndrome, depressive neurosis and others [35]. It is special type of acupuncture that inserts medical threads into skin, subcutaneous tissue or muscles at specific points [36]. Although acupoint catgut embedding therapy is considered an invasive treatment from a western medicine perspective, it has attracted considerable attention from clinicians in China due to its easy operation and durable stimulation, and has been widely used in recent years. Suture buried in acupoints can produce persistent stimulation on the basis of selecting sutures and acupoints according to deficiency and excess, which aims to regulate *yin* and *yang*, dredge the channels and collaterals, and reinforce the anti-pathogenic *Qi* and eliminate the pathogenic *Qi*. It has dual rapid and continuous action for chronic diseases. It is the combination of a variety of therapies (acupoint blocking, acupuncture, pricking, the after effect of tissue injury, needle retention). The intensity of stimulus would be changed with the time and the special needles and sutures can produce stronger efficacy than conventional acupuncture on controlling mind (*shen*) to facilitate *Qi* flow and dredging the meridian to regulate *Qi* and blood [37].

The effects of acupoint catgut embedding therapy in Western Medicine are similar to those of manual acupuncture. It provides both physical and chemical stimulation [38, 39].

The absorbable surgical thread, a foreign protein, can induce allergic reactions and the combined effects of proteolytic enzymes and macrophage action against the absorbable surgical thread may strengthen and extend the acupoint stimulation for 15–20 days as a consequence of the mild irritation in subcutaneous tissue, inducing a more persistent and potent physiological stimulation produced by the suture at the acupoints [39]. Acupoint embedding therapy improves body's nutrient metabolism and promotes blood circulation [37].

By using the specially made disposable embedding needle, the operation of micro-invasive thread embedding can be completed swiftly and conveniently in the way of injection. The embedding needle has a handle to hold and allows rapid inserting by only a hand. The mechanical spring ensures successful embedding of the thread. The marks on the needle body make it easy to control the inserting depth. The whole operation only takes 5–10 min, as only the thread not the needle needs to be retained. The stimulation produced by the retained thread can last for 1–2 weeks, during which the patients can move freely without any influence on their life. Therefore, this method can be applied in clinic safely for patients' convenience and less pain [40].

Patient should be informed about the local pain during operation and the postoperative reaction.

After the insertion, may occur such reactions as: a) the local skin will manifest redness, swelling, hotness and pain of aseptic reaction; b) hematoma may occur due to the needling injury and thread; c) general response with a fever of about 38°C in 4–24hs, or even persistent high fever [37].

Thread-embedding acupuncture is a new subtype of acupuncture treatment developed from catgut-embedding therapy [35]. Different biodegradable materials have been developed and widely used. They are divided into natural and synthetic types according to material sources. Both of them have advantages and disadvantages [41]. For instance, natural embedding material is inexpensive and plentiful, but it also easily leads to an infection reaction due to poor biocompatibility. Hence, it has a higher applied risk due to the instable quality [42]. In contrast, synthetic embedding materials, such as polyglycolic acid [43] and polylactic acid [44], offer an excellent biodegradability and biocompatibility, and have a stable property through the whole implanting period. But their poor hydrophilicity and cell adhesion, inevitable large dimensions for producing a lasting effect largely affected their therapeutic efficacy in clinical application [45, 46].

There is a number of uncertainties, including the spinning speed, drawing temperature and stretching ratio, which affect compression behaviors in the preparation process of acupoint embedding monofilaments [47]. In addition, rigid acupoint embedding monofilaments will stimulate the subcutaneous tissue of the human body, resulting in a greater sense of pain for patients in clinical application. In contrast, soft acupoint embedding monofilaments may lead to difficulties in the embedding operation [48]. However, there is still a lack of relative study and a unified standard for a test method for monofilament compression behaviors [49].

The ideal embedding materials are required to provide properties such as safety, non-toxicity, good biocompatibility and absorbability [50, 51]. Antibacterial properties are essential in the prevention of infections [52, 53]. Also, embedding materials should provide enough mechanical properties to be implanted into the body and support peripheral nerve tissue [54]. Through good swelling behavior they are able to produce enough stimulus degree *in vivo*, yet retain a relatively small dimension *in vitro* and avoid the significant trauma [55]. Favorable hydrophilicity is also important for embedding material to adhere cells and work well in the human body [56].

#### Discussion

Catgut is a protein fiber of biological origin, that is derived from the small intestines of animals, mostly sheep or oxen [56]. Not many studies have investigated the mechanism of catgut implantation at acupoints. Li et al. explored the probable mechanism focusing on neurogenic inflammation [57]. Some studies have demonstrated that nerve stimulation could induce leukocyte activation and plasma

extravasation, which is termed neurogenic inflammation [58, 59]. The mechanisms involve the regulation of nerve conduction, signal pathways, hormone level, protein expression, oxidative stress level, and structure restoration. The treatment generally regulates some specific biochemical factors to influence particular signal pathways, eventually controlling the apoptosis and proliferation of cells and repairing the damaged structure [60]. The mechanism of acupuncture embedding regards catgut as a type of heterogeneous protein, after inserting it into acupoints, its process of softening, decomposition, liquification and absorption can effectively promote and enhance the nutrition metabolism, stress ability of bodies, vascular permeability and blood circulation [61]. Areas of catgut suture degradation contain dense accumulation of macrophages, lymphocytes, and foreign body giant cells. After complete absorption, these are replaced by a dense mass of macrophages [62]. Most studies indicate that catgut sutures are completely absorbed between 35 and 60 days [63]. However, the safety of catgut embedding has not yet been established, since it can cause various immune reactions, including allergic reactions and a subcutaneous nodule [64].

**Polydioxanone (PDO)** has been used in acupuncture embedding therapy devices in Korea. It is a synthetic monofilamentous polymer made of polyester or a polymer of polydioxanone. PDO retains 50% tensile strength after 4 weeks, takes 180 days to be absorbed, and also has low tissue reactivity [65]. In Korea, the embedding of PDO thread at certain acupoints is widely used in the clinical practice for the treatment of chronic musculoskeletal pain. Nevertheless, the current level of evidence supporting the efficacy of embedding with PDO thread for patients with chronic musculoskeletal pain is insufficient [66]. Only case studies on the application of PDO embedding therapy for the patients with chronic low back pain treatment [67], shoulder pain [68], and osteoarthritis of the knee [69] have been published. In addition, no randomized clinical trials using PDO thread embedded as a sham-controlled intervention have been reported [70].

**Polyglycolic acid (PGA)** has held great promise for various biomedical applications in human tissue due to its advantages of being easily available, low cost and having excellent biodegradability and biocompatibility [71]. PGA monofilaments prepared by melt-spinning technology for acupuncture embedding therapy materials have exhibited sufficient supply, good formation capacity and versatility in terms of surface functionalization [72, 73]. The PGA monofilament is considered to be a highly promising biodegradable material for ideal acupuncture embedding therapy material preparation [74, 75]. PGA is broken down by hydrolysis into its respective acids and alcohols. It tends to lose mechanical strength rapidly, over a period of 2–4 weeks after implantation [76]. Some shortcomings were discovered and they may increase the risk of PGA implantations, such as insufficient hydrophilicity and cell adhesion caused by the relatively smooth surfaces [45]. These features are believed to be the main contributors to the unfavorable characteristics [77].

**Polylactic acid (PLA)** is one of the highest consuming bioplastics in the world. It is an aliphatic polyester obtained from renewable sources such as corn sugar, starch, potato, and sugarcane. PLA has 37% crystallinity, elongation at break 30.7%, glass transition temperature of 53°C, and a melting temperature ranging between 170–180°C [78]. It is used in the biomedical area, and has also begun to be applied as an embedding material for curing osteoporosis [79] and pseudo-myopia [80] and other diseases. However, its long degradation time (about 2 years) and poor hydrophilicity have limited its further application in acupuncture embedding therapy [81].

**Polylactic glycolic acid (PLGA)** is polymerized by 9 polyglycolic acids (PGA) and one polylactide acid (PLA). It is simple in composition without bioproteins, and is now widely used as embedding material for its proper rigidity. When implanted in the acupuncture point of human body, it can produce a long-time stimulation which is similar to that produced by a filiform needle [82]. Histologic examination showed that the PLGA sutures were absorbed within 90 days [83], which might be the underlying mechanism of persistent analgesic effects [84]. PLGA has good biocompatibility, and no adverse reactions have been reported. Hydrophilicity, cell adhesion and degradation properties can be improved by surface modification technologies [85].

#### Surface modification technologies

Currently, there are numerous surface modification technologies, such as dip-coating, ultrasound, chemical vapor deposition, ion beam injection and surface graft polymerization, that can be applied to biomedical materials [86, 87]. Wang et al. attempted to use ultrasound treatment to modify PGA and poly(lactic-glycolic acid) (PLGA) fibers, and the results suggested that both PGA and PLGA fibers achieved better hydrophilicity and cytocompatibility, while the tensile strength of PLGA increased and that of PGA changed little [88]. Among a variety of surface modification technologies, cold plasma modification has been regarded as one of the simplest and highly effective methods to modify biomaterials. It can promote the formation of new oxidized functional groups by introducing potential activation sites on their surfaces, and modify the remaining original properties of materials to a great extent without damaging their outermost surfaces [89, 90]. Song et al. point out that the photo-degradation, thermal, and microbial biodegradable properties of the PLA films can be significantly improved by plasma modification [91].

It is reported that solution dip-coating is one of the most widely used processes in textile manufacturing and the simplest functionalization technique for material surfaces, and chitosan is a promising natural compound, which possesses a prospective future due to its advantages of non-toxicity, antibacterial properties, biocompatibility, biodegradation and swelling properties [92, 93]. Chitosan is a linear, high molecular weight heteropolysaccharide [94], consisting of N-acetyl-glucosamine and N-glucosamine units [95]. With its abundant reserve, chitosan is the second most important natural polymer globally (the first is cellulose) and has been

widely extracted from marine arthropods (prawns, crabs, shellfish, etc.) [96]. Hence, numerous chitosan coating applications on tissue engineering areas have been reported. Dubnika et al. used chitosan as an antibacterial agent to functionalize scaffolds and achieved a significant effect [97]. Niekraszewicz et al. used chitosan to modify polypropylene (PP) mesh and achieved good results in terms of mechanical and chemical properties; the biological purity was also improved [98]. Umair and colleagues reported that the PGA suture using N-halamine-based chitosan agents, its antibacterial efficacy was enhanced and could kill both *E. coli* and *S. aureus* bacteria within 15 min of contact time [99]. However, there is still a lack of detailed research about chitosan coating methods of monofilaments and existing studies prefer to target the biocompatibility, proliferation capacity, etc. Meanwhile, there are few reports on the self-swelling behaviors of embedding materials improved by chitosan coating [100].

### Conclusions

1. Acupoint embedding therapy is an invasive treatment which can prolong point stimulation, reduces the frequencies of pain and psychological fear of patients and visits to the doctors.

2. The acupoint embedding therapy seems to be a promising method of neuropathy treatment.

3. The ideal embedding materials are required to provide properties such as safety, non-toxicity, good biocompatibility, excellent swelling and biodegradation behaviors.

4. The properties of the filaments for acupoint embedding therapy can be improved by surface modification technologies.

5. It will be great if the embedded material has also an antibacterial or active factor releasing effect which can be achieved by dip coating with chitosan or other substances.

### References

- Huang KC. Acupuncture: the past and the present. New York: Vantage; 1996. 275 p.
- Ma KW. The roots and development of Chinese acupuncture: from prehistory to early 20th century. *Acupunct Med* 1992;10(Suppl):92-9.
- Baldry PE. Acupuncture, trigger points and musculoskeletal pain. Edinburgh: Churchill Livingstone; 1993.
- Bivens RE. Acupuncture, expertise and cross-cultural medicine. Manchester: Palgrave; 2000.
- Sun WS. Micro-invasive thread-embedding therapy. *Shanghai J Acupunct*. 2010;29(1):65.
- Leung L. Neurophysiological basis of acupuncture-induced analgesia: an updated review. *J Acupunct Meridian Stud*. 2012;5(6):261-270.
- Park JY, Namgung U. Electroacupuncture therapy in inflammation regulation: current perspectives. *J Inflamm Res*. 2018;11:227-237. doi: 10.2147/JIR.S141198.
- Ma W, Li Z, Lu Z, et al. Protective effects of acupuncture in cardiopulmonary bypass-induced lung injury in rats. *Inflammation*. 2017;40(4):1275-1284
- Hsu E, editor. Innovation in Chinese Medicine. Cambridge: Cambridge University Press; 2001. xv, 426 p. (Needham Research Institute Studies; 3).
- Kong J, Gollub R, Huang T, et al. Acupuncture of Qi, from qualitative history to quantitative measurement. *J Altern Complement Med*. 2007;13(10):1059-1070.
- White P, Bishop F, Hardy H, et al. Southampton needle sensation questionnaire: development and validation of a measure to gauge acupuncture needle sensation. *J Altern Complement Med*. 2008;14(4):373-379.
- Li XH, Liu YC, Gong FM, et al. Clinical study on acupuncture for treatment of juvenile moderate and mild myopia. *Chin Acupunct Moxibust*. 2003;23:147-150.
- Run-Ming Y. The origin and development of Chinese acupuncture and moxibustion. *Anc Sci Life*. 1985;4(4):224-228.
- World Health Organization. Acupuncture: review and analysis of reports on controlled clinical trials. Geneva: World Health Organization; 2002.
- Amato AA, Russell JA. Neuromuscular disorders. New York: McGraw-Hill Medical; 2008. 775 p.
- Jin WH, Wei ZR, Sun GF, Tang XJ, Wang DL. Autologous peripheral blood stem cell transplantation for treatment of peripheral nerve injury. *Chin J Tissue Eng Res*. 2013;17(1):181-185.
- Ma HB, Zhang RM. Chemically extracted acellular allogenic nerve and accompanying peripheral vein for repair of facial nerve defects. *Chin J Tissue Eng Res*. 2013;17:3325-32.
- Dimitrova A. Introducing a standardized acupuncture protocol for peripheral neuropathy: a case series. *Med Acupunct*. 2017;29(6):352-365. doi:10.1089/acu.2017.1242.
- He GH, Ruan JW, Zeng YS, Zhou X, Ding Y, Zhou GH. Improvement in acupoint selection for acupuncture of nerves surrounding the injury site: electro-acupuncture with Governor vessels with local meridian acupoints. *Neural Regen Res*. 2015 Jan;10(1):128-135. doi: 10.4103/1673-5374.150720.
- Lu MC, Ho CY, Hsu SF, et al. Effects of electrical stimulation at different frequencies on regeneration of transected peripheral nerve. *Neurorehabil Neural Repair*. 2008;22(4):367-373.
- Hollisaz MT. Use of electroacupuncture for treatment of chronic sciatic pain. *Internet J Pain Symptom Control Palliat Care*. 2006;5(1):1-4.
- Chon TY, Lee MC. Acupuncture. *Mayo Clin Proc*. 2013;88(10):1141-1146. doi: 10.1016/j.mayocp.2013.06.009.
- Inoue M, Kitakoji H, Yano T, Ishizaki N, Itoi M, Katsumi Y. Acupuncture treatment for low back pain and lower limb symptoms – the relation between acupuncture or electroacupuncture stimulation and sciatic nerve blood flow. *Evid Based Complement Alternat Med*. 2008;5(2):133-143. doi: 10.1093/ecam/nem050.
- Cheng KJ. Neuroanatomical basis of acupuncture treatment for some common illnesses. *Acupunct Med*. 2009;27(2):61-64. doi: 10.1136/aim.2009.000455.
- Goldman N, Chen M, Fujita T, et al. Adenosine A1 receptors mediate local anti-nociceptive effects of acupuncture. *Nat Neurosci*. 2010;13(7):883-888. doi: 10.1038/nn.2562.
- Kaptchuk TJ. Acupuncture: theory, efficacy, and practice. *Ann Intern Med*. 2002;136(5):374-383. doi: 10.7326/0003-4819-136-5-200203050-00010.
- Han JS. Acupuncture: neuropeptide release produced by electrical stimulation of different frequencies. *Trends Neurosci*. 2003;26(1):17-22.
- Schröder S, Liepert J, Reppis A, Gretten JH. Acupuncture treatment improves nerve conduction in peripheral neuropathy. *Eur J Neurol*. 2007 Mar;14(3):276-81.
- Litscher G, Wang L, Huber E, Nilsson G. Changed skin blood perfusion in the fingertip following acupuncture needle introduction as evaluated by laser Doppler perfusion imaging. *Lasers Med Sci*. 2002;17:19-25.
- Nathan DM, Genuth S, Lachin J, Cleary P, Crofford O, Davis M, Rand L, Siebert C. Diabetes control and complication trial group (DCCT). The effect of intense treatment of diabetes on the development and progression of long-term complication in insulin-dependent diabetes mellitus. *N Eng J Med*. 1993;329(14):977-986.
- Samsø E, Farber NE, Kampine JP, Schmeling WT. The effects of halothane on pressor and depressor responses elicited via the somato-sympathetic reflex: a potential antinociceptive action. *Anesth Analg*. 1994;79(5):971-979.
- Sato A, Schmidt RF. Somatosympathetic reflexes: afferent fibers, central pathways, discharge characteristics. *Physiol Rev*. 1973;53:916-947.
- Zhu W, Ye Y, Liu Y, et al. Mechanisms of acupuncture therapy for cerebral ischemia: an evidence-based review of clinical and animal studies on cerebral ischemia. *J Neuroimmune Pharmacol*. 2017;12(4):575-592. doi: 10.1007/s11481-017-9747-4.

34. Chung A, Bui L, Mills E. Adverse effects of acupuncture. Which are clinically significant? *Can Fam Physician*. 2003;49:985-989.
35. Huang CY, Choong MY, Li TS. Treatment of obesity by catgut embedding: an evidence-based systematic analysis. *Acupunct Med*. 2012;30(3):233-4.
36. Shin HJ, Lee DJ, Kwon K, et al. The success of thread-embedding therapy in generating hair re-growth in mice points to its possibly having a similar effect in humans. *J Pharmacopuncture*. 2015;18(4):20-5.
37. Xu Y, Cai J, Liang L, Chen G, Xu X. The application of nanotechnology in acupoint catgut embedding therapy. *Mater Sci Forum*. 2011;694:68-72.
38. Lee KH, Lee DH, Kwon KR, et al. A literary study on embedding therapy. *J Pharmacopuncture*. 2003;6:15-21.
39. Guo Y, Fang JQ, Wang RH, Kong LH, Chen RX, Lin YP, et al. *Experimental Acup-Moxibustion Science*. 3rd ed. Beijing: China Traditional Chinese Medicine Press; 2012. p. 169-171.
40. Sun W. Introduction to micro-invasive thread-embedding therapy. *J Acupunct Tuina Sci*. 2012;10(3):196-198. doi: 10.1007/s11726-012-0602-0.
41. Kondiah PJ, Choonara YE, Kondiah PPD, et al. A review of injectable polymeric hydrogel systems for application in bone tissue engineering. *Molecules (Basel, Switzerland)*. 2016;21(11). pii: E1580
42. Chifiriu MC, Grumezescu AM, Grumezescu V, et al. Biomedical applications of natural polymers for drug delivery. *Curr Org Chem*. 2014;18:152-164.
43. Lopes MS, Jardim AL, Maciel R. Poly (lactic acid) production for tissue engineering applications. *Proced Eng*. 2012;42:1402-1413.
44. Yi LF, Luo SS, Shen JB, et al. Bioinspired polylactide based on the multilayer assembly of shish-kebab structure: a strategy for achieving balanced performances. *ACS Sustain Chem Eng*. 2017;5:3063-3073.
45. Sun X, Xu C, Wu G, et al. Poly(lactic-co-glycolic acid): applications and future prospects for periodontal tissue regeneration. *Polymers*. 2017;9(6). pii: E189.
46. Liu H. Clinical comparison of poly-lactic-co-glycolic acid new catgut embedding material and catgut. *Chin J Tissue Eng Res*. 2012;16:7185-7192.
47. Capozza RC. Spinning and shaping poly-(N-acetyl-D-glucosamine). U.S. Patent 3,988,411[P]. 1976-10-26, USA, 1976.
48. Wang J, Liu W, Liang R, et al. Analytical and experimental study on flexural behavior of pultruded fibre reinforced polymer sheet piles. *J Compos Mater*. 2015;50:1-8.
49. Qi Y, Xiong W, Liu W, et al. Experimental study of the flexural and compression performance of an innovative pultruded glass-fiber-reinforced polymer-wood composite profile. *Plos One*. 2015;10:e0140893
50. Li Z, He GD, Hua JC, et al. Preparation of gamma-PGA hydrogels and swelling behaviors in salt solutions with different ionic valence numbers. *RSC Adv*. 2017;7:11085-11093.
51. Ding M, Feng H, Jin C, et al. Clinical research on microinvasive acupoint thread-embedding therapy in treating refractory prosopoplegia. *J Nanjing Univ Trad Chin Med*. 2016;32:21-24.
52. Norton C, Czuber-Dochan W, Artom M, et al. Systematic review: interventions for abdominal pain management in inflammatory bowel disease. *Aliment Pharmacol Ther*. 2017;46(2):115-125.
53. Petca A, Radu DC, Zvanca M, et al. Suture materials and technics, possible cause for C-section scar defect. *Key Eng Mater*. 2017;752:54-58.
54. Inan BB. Acupuncture stimulation analyzed from multiple aspects of western medical science. *Acupunct Electro Therapeut Res*. 2016;41:21-37.
55. Fang S, Wang M, Zheng Y, et al. Acupuncture and lifestyle modification treatment for obesity: a meta-analysis. *Am J Chin Med*. 2017;45:239-254.
56. Rajendran S, Anand S, Rigby A. Textiles for healthcare and medical applications. In: Horrocks R, Anand S, editors. *Handbook of technical textiles. Volume 2: Technical textile applications*. 2nd ed. Oxford: Elsevier; 2016. p. 135-168.
57. Li X, Liu Y, Zhang Q, Xiang N, He M, Zhong J, Chen Q, Wang X. Effect of catgut implantation at acupoints for the treatment of allergic rhinitis: a randomized, sham-controlled trial. *BMC Complement Altern Med*. 2016;16(1):454. doi: 10.1186/s12906-016-1400-x.
58. Umeno E, Nadel JA, McDonald DM. Neurogenic inflammation of the rat trachea: fate of neutrophils that adhere to venules. *J Appl Physiol*. 1990;69(6):2131-6.
59. McDonald DM. Neurogenic inflammation in the rat trachea. I. Changes in venules, leucocytes and epithelial cells. *J Neurocytol*. 1988;17(5):583-603.
60. Feng Y, Fang Y, Wang Y, Hao Y. Acupoint therapy on diabetes mellitus and its common chronic complications: a review of its mechanisms. *BioMed Res Int*. 2018;3128378. doi: 10.1155/2018/3128378.
61. Bogao B, Liugang F. Prediction and experimental research on acupoint catgut embedding for mechanism of fatigue mitigation in rat. *J Guangzhou Sport University*. 2013;33:95.
62. Lawrie P, Angus G, Reese AJ. The absorption of surgical catgut. II. The influence of size. *Br J Surg*. 1960;47:551-555.
63. Saini P, Arora M, Kuma, MR. Poly (lactic acid) blends in biomedical applications. *Adv Drug Deliv Rev*. 2016;107:47-59.
64. Chuang YT, Li TS, Lin TY, Hsu CJ. An unusual complication related to acupuncture point catgut embedding treatment of obesity. *Acupunct Med*. 2011;29(4):307-308.
65. Tajirian AL, Goldberg DJ. A review of sutures and other skin closure materials. *J Cosmet Laser Ther*. 2010;12(6):296-302.
66. Kim E, Kim HS, Jung SY, Han CH, Kim YI. Efficacy and safety of polydioxanone thread embedded at specific acupoints for non-specific chronic neck pain: a study protocol for a randomized, subject-assessor-blinded, sham-controlled pilot trial. *Trials*. 2018 Dec 6;19(1):672. doi: 10.1186/s13063-018-3058-9.
67. Jang H, Woo C, Ahn H, Kwon H. A clinical four case studies on chronic low back pain treated by needle embedding therapy. *J Korea CHUNA Manual Med Spine Nerves*. 2014;9(2):45-55.
68. Lee SM, Ji YS, Jeon JH, Kim JH, Kim YI. Effect of needle-embedding and acupuncture therapy on shoulder pain in Behcet disease patient: a case report. *J Acupunct Res*. 2013;30(4):219-24.
69. Lee JH, Yang TJ, Lee DG, Lee OJ, Wei TS. The effect of needle-embedding therapy on osteoarthritis of the knee combined with Korean medical treatment: report of five cases. *J Acupunct Res*. 2014;31(4):195-204.
70. Kwon K. The analysis on the present condition of thread-embedding therapy papers published in *Journal of Korean Medicine*. *J Korean Med Ophthalmol Otolaryngol Dermatol*. 2014;27(4):16-44.
71. Wade RJ, Burdick JA. Advances in nanofibrous scaffolds for biomedical applications: from electrospinning to self-assembly. *Nano Today*. 2014;9:722-742.
72. Tyler B, Gullotti D, Mangraviti A, et al. Polylactic acid (PLA) controlled delivery carriers for biomedical applications. *Adv Drug Deliv Rev*. 2016;107:163-175.
73. Sridhar R, Pliszka D, Luo HK, et al. Medical devices regulatory aspects: a special focus on polymeric material based devices. *Curr Pharmaceut Des*. 2015;21(42):6246-6259.
74. Ortiz M, Rosales-Ibanez R, Pozos-Guillen A. DPSC colonization of functionalized 3D textiles. *J Biomed Mater Res B Appl Biomater*. 2017;105(5):785-794.
75. Katz AR, Turner RJ. Evaluation of tensile and absorption properties of PGA sutures. *Surg Gynecol Obstet*. 1970;131(4):701-16.
76. Balaji AB, Pakalapati H, Khalid M, Walvekar R, Siddiqui H. Natural and synthetic biocompatible and biodegradable polymers. In: Navinchandra Gopal Shimpi. *Biodegradable and biocompatible polymer composites*. Oxford: Elsevier; 2018. p. 3-32.
77. Tummalapalli M, Anjum S, Kumari S, et al. Antimicrobial surgical sutures: recent developments and strategies. *Polym Rev*. 2016;56:607-630.
78. Vroman I, Tighzert L. Biodegradable polymers. *Materials*. 2009;2:307-344.
79. Degala S, Shetty S, Ramya S. Fixation of zygomatic and mandibular fractures with biodegradable plates. *Ann Maxillofacial Surg*. 2013;3:25-30.
80. Wang B, Lu XH, Zhang YQ. Excimer laser photorefractive keratectomy for myopia with thin central cornea: report of 67 cases. *Acad J First Med College PLA*. 2003;23:986.
81. Li H, Liao H, Bao C, et al. Preparation and evaluations of mangiferin-loaded PLGA scaffolds for alveolar bone repair treatment under the diabetic condition. *AAPS PharmSciTech*. 2017;18(2):529-538.

82. Sun WS, Tan ZQ. Novel biomedical material and its application value in micro-invasive thread-embedding therapy. *Shanghai J Acupunct.* 2010;29(2):131-132.
83. Craig PH, Williams JA, Davis KW, Magoun AD, Levy AJ, Bogdan-sky S, Jones JP Jr. A biologic comparison of polyglactin 910 and polyglycolic acid synthetic absorbable sutures. *Surg Gynecol Obstet.* 1975;141(1):1-10.
84. Munteanu RM, Eva L, Dobrovăț BI, et al. Longer survival of a patient with glioblastoma resected with 5-aminolevulinic acid (5-ALA)-guided surgery and foreign body reaction to polyglycolic acid (PGA) suture. *Rom J Morphol Embryol.* 2017;58(2):671-680.
85. Chu YX, Cui WQ, Xu F, et al. Acupoint embedding of polyglactin 910 sutures in patients with chronic pain due to cervical spondylotic radiculopathy: a multicenter, randomized, controlled clinical trial. *Evid Based Complement Alternat Med.* 2018;3465897. doi: 10.1155/2018/3465897.
86. Ritz U, Nusselt T, Sewing A, et al. The effect of different collagen modifications for titanium and titanium nitrite surfaces on functions of gingival fibroblasts. *Clin Oral Investig.* 2017;21:255-265.
87. Pester CW, Poelma JE, Narupai B, et al. Ambiguous anti-fouling surfaces: facile synthesis by light-mediated radical polymerization. *J Polym Sci Part A: Polym Chem.* 2016;54(2):253-262.
88. Wang B, Zhang P, Song W, et al. Modification of polyglycolic acid and poly lactic-co-glycolic acid fibers by ultrasonic treatment for enhancing hydrophilicity and cytocompatibility. *J Ind Text.* 2016;45:516-530.
89. Sanbhal N, Mao Y, Sun G, et al. Surface modification of polypropylene mesh devices with cyclodextrin via cold plasma for hernia repair: characterization and antibacterial properties. *Appl Surf Sci.* 2018;439:749-759.
90. Wang M, Favi P, Cheng X, et al. Cold atmospheric plasma (CAP) surface nanomodified 3D printed polylactic acid (PLA) scaffolds for bone regeneration. *Acta Biomater.* 2016;46:256-265.
91. Song AY, Oh YA, Roh SH, et al. Cold oxygen plasma treatments for the improvement of the physicochemical and biodegradable properties of polylactic acid films for food packaging. *J Food Sci.* 2016;81(1):e86-e96.
92. Raftery R, O'Brien FJ, Cryan SA. Chitosan for gene delivery and orthopedic tissue engineering applications. *Molecules.* 2013;18(5):5611-5647.
93. Rasti H, Parivar K, Baharara J, et al. Chitin from the Mollusc Chiton: extraction, characterization and chitosan preparation. *Iran J Pharm Res.* 2017;16:366-379.
94. Raafat D, Leib N, Wilmes M, et al. Development of *in vitro* resistance to chitosan is related to changes in cell envelope structure of *Staphylococcus aureus*. *Carbohydr Polym.* 2017;157:146-155.
95. Hps AK, Saurabh CK, Adnan A, et al. A review on chitosan-cellulose blends and nanocellulose reinforced chitosan biocomposites: properties and their applications. *Carbohydr Polym.* 2016;150:216-226.
96. Islam M, Mondal IH, Ahmed F. Study on prawn shell waste into chitosan and its derivatives as value added products for cellulosic fibres. *Res J Text Apparel.* 2017;21(2):134-145.
97. Dubnika A, Loca D, Rudovica V, et al. Functionalized silver doped hydroxyapatite scaffolds for controlled simultaneous silver ion and drug delivery. *Ceram Int.* 2017;43:3698-3705.
98. Niekraszewicz A, Kucharska M, Wawro D. Development of a manufacturing method for surgical meshes modified by chitosan. *Fibres Text East Eur.* 2007;3(62):105-109.
99. Umair MM, Jiang Z, Ullah N, et al. Development and characterisation of antibacterial suture functionalised with N-halamines. *J Ind Text.* 2016;46(1):59-74.
100. Fu S, Zhang P. A study of the chitosan coating method on polyglycolic acid and polylactic acid embedding monofilaments. *Text Res J.* 2019;89(3):270-280.

#### Authors' ORCID iDs and academic degrees

Olga Ignatov – <https://orcid.org/0000-0003-2911-3239>, MD  
Oleg Pascal – <https://orcid.org/0000-0002-4870-2293>, MD, PhD.  
Viorel Nacu – <https://orcid.org/0000-0003-2274-9912>, MD, PhD.

#### Authors' contributions

OI designed the trial, wrote the first draft of manuscript. OP interpreted the data. VN revised the manuscript critically. All the authors revised and approved the final version of the manuscript.

#### Funding

The publication of this article was possible due to the project No 20.80009.5007.20 GaN-based nanoarchitectures and three-dimensional matrices from biological materials for applications in microfluidics and tissue engineering.

This study was supported by *Nicolae Testemitsanu* State University of Medicine and Pharmacy. The authors are independent and take responsibility for the integrity of the data and accuracy of the data analysis.

#### Ethics approval and consent to participate

No approval was required for this review study.

#### Conflict of Interests

No competing interests were disclosed.