

Article

Open Access

Potential aquatic environmental risks of trifloxystrobin: Enhancement of virus susceptibility in zebrafish through initiation of autophagy

Huan Wang^{1,2,3}, Tian-Xiu Qiu^{1,2,3}, Jian-Fei Lu^{1,2,3}, Han-Wei Liu⁴, Ling Hu⁴, Lei Liu^{1,2,3,*}, Jiong Chen^{1,2,3,*}

¹ State Key Laboratory for Managing Biotic and Chemical Threats to the Quality and Safety of Agro-products, Ningbo University, Ningbo, Zhejiang 315211, China

² Laboratory of Biochemistry and Molecular Biology, School of Marine Sciences, Meishan Campus, Ningbo University, Ningbo, Zhejiang 315832, China

³ Key Laboratory of Applied Marine Biotechnology of Ministry of Education, Meishan Campus, Ningbo University, Ningbo, Zhejiang 315832, China

⁴ Ningbo Customs District Technology Center, Ningbo, Zhejiang 315832, China

ABSTRACT

Chronic pollution in aquatic ecosystems can lead to many adverse effects, including a greater susceptibility to pathogens among resident biota. Trifloxystrobin (TFS) is a strobilurin fungicide widely used in Asia to control soybean rust. However, it has the potential to enter aquatic ecosystems, where it may impair fish resistance to viral infections. To explore the potential environmental risks of TFS, we characterized the antiviral capacities of fish chronically exposed to TFS and subsequently infected with spring viraemia of carp virus (SVCV). Although TFS exhibited no significant cytotoxicity at the tested environmental concentrations during viral challenge, SVCV replication increased significantly in a time-dependent manner within epithelioma papulosum cyprini (EPC) cells and zebrafish exposed to 25 µg/L TFS. Results showed that the highest viral load was more than 100-fold that of the controls. Intracellular biochemical assays indicated

that autophagy was induced by TFS, and associated changes included an increase in autophagosomes, conversion of LC3-II, accumulation of Beclin-1, and degradation of P62 in EPC cells and zebrafish. In addition, TFS markedly decreased the expression and phosphorylation of mTOR, indicating that activation of TFS may be associated with the mTOR-mediated autophagy pathway. This study provides new insights into the mechanism of the immunosuppressive effects of TFS on non-target aquatic hosts and suggests that the existence of TFS in aquatic environments may contribute to outbreaks of viral diseases.

Keywords: Trifloxystrobin; Autophagy; SVCV; Chronic toxicity; Susceptibility

Received: 20 February 2021; Accepted: 07 May 2021; Online: 08 May 2021

Foundation items: This work was supported by the National Natural Science Foundation of China (31902410), Program of State Key Laboratory for Managing Biotic and Chemical Threats to the Quality and Safety of Agro-products (KF20200106), Natural Science Foundation of Zhejiang Province (LY21C190002), and Foundation of Ningbo City of China (202003N4120)

*Corresponding authors, E-mail: liulei2@nbu.edu.cn; jchen1975@163.com

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright ©2021 Editorial Office of Zoological Research, Kunming Institute of Zoology, Chinese Academy of Sciences

INTRODUCTION

Strobilurins are a type of agricultural fungicide developed from natural β -methoxyacrylate derivatives (Bartlett et al., 2002). Over the last 20 years, strobilurins have been increasingly applied to crop production systems as the most important component of crop disease management (Deb et al., 2010; Feng et al., 2020). Because strobilurins impart broad-spectrum, rapid, and highly efficient germicidal activities, and because these fungicides are cost effective and degrade rapidly during plant metabolism, their large-scale use has increased considerably; indeed, this class represents the most important fungicide group, accounting for 20% of the total fungicide market in 2016 (Wang et al., 2020).

Trifloxystrobin (TFS, methyl- α -methoxyimino-2-((1-(3-(trifluoromethyl) phenyl) ethylidene ((amino) oxymethyl)-benzeneacetate)), a systemic broad-spectrum foliar fungicide synthesized from the natural antibiotic strobilurin with an oximether side chain, is widely used to control fungi in greenhouses, nurseries, orchards, vegetable gardens, and paddy fields (Zhu et al., 2015b). Due to its rapid degradation in plants, soil, and water, as well as its long-lasting effects and resistance to rain, it occupies an important position among strobilurins (Luo et al., 2020). However, the widespread application of TFS has raised serious public health concerns, in particular, that the potential off-site transport of environmental residues could have non-target toxic effects on humans and wildlife (Liu et al., 2020b).

TFS exhibits relatively high water solubility and has a half-life of 1.9 to 9.8 d in soil (Wang et al., 2015). These properties promote its entry into water directly following aerial spraying or indirectly through leaching and drainage flow, after which it accumulates in water and sediment (Komárek et al., 2010; Wang et al., 2015). TFS has been frequently detected in aquatic systems, e.g., at concentrations of 0.73 $\mu\text{g/L}$ in horticultural catchments of southeastern Australia and of 170 $\mu\text{g/L}$ in Chinese paddy field water (Wightwick et al., 2012; Cao et al., 2015). Although the European Food Safety Authority (EFSA, 2017) states that TFS exhibits low acute and chronic toxicity to humans, birds, mammals, and bees, several studies have indicated that TFS can affect various biotic components of freshwater ecosystems and can be acutely toxic to some non-target freshwater organisms (Cui et al., 2017; Junges et al., 2012; Li et al., 2018; Liu et al., 2013; Zhu et al., 2015a). Thus, it is important to investigate the potential adverse effects of TFS on aquatic organisms.

Although viruses are essential intracellular agents, they can cause devastating diseases in aquatic animals because of limited effective treatments and prophylactics (Xiong et al., 2019). Given the increasing demand for fish and shellfish in contemporary societies, losses caused by viral diseases must be reduced (Balmer et al., 2017). According to the World Organisation for Animal Health (OIE), spring viraemia of carp (SVC) is a notifiable disease responsible for significant mortality in carp species, including the common carp (*Cyprinus carpio*), koi carp (*C. carpio koi*), crucian carp (*Carassius carassius*), grass carp (*Ctenopharyngodon idella*), and goldfish (*Carassius auratus*) (Ahne et al., 2002; Taylor et al., 2013). Since its first detection in Yugoslavia, the SVC virus

(SVCV) has spread to Europe, South America, North America, and Asia (Dikkeboom et al., 2004; Garver et al., 2007). At present, all aquatic organisms infected by SVCV are destroyed to eradicate the virus in affected ponds (Ahne et al., 2002), but viral outbreak remains a crucial issue to ecological security.

Depending on their physicochemical properties, pollutants may accumulate in aquatic organisms at low environmental concentrations and interact with tissues and cells, potentially leading to a series of chronic poisoning characteristics, including cell cycle arrest and endoplasmic reticulum stress. These events can, in turn, transduce messages to stress sensors related to oxidative stress and immune system damage (Ma et al., 2019; Xing et al., 2015). These physiological disorders can increase the virulence of pathogens and reduce host resistance. TFS can induce embryonic lesions, decrease hatching success, and increase oxidative stress in zebrafish embryos by increasing reactive oxygen species (ROS) and malonaldehyde (MDA) content, and inhibiting superoxide dismutase (SOD), glutathione (GSH), catalase (CAT) and antioxidant system-related mRNA genes (i.e., *Mn-SOD*, *Cu/Zn-SOD*, *CAT*, *Nrf2*, and *Bcl2*) (Li et al., 2018; Zhu et al., 2015a). These cellular components are crucial for maintaining antiviral resistance in fish against the acutely contagious and hemorrhagic SVC disease (Liu et al., 2020a; Yuan et al., 2014). Therefore, the above studies raise concern that chronic exposure to TFS at environmental concentrations may affect the resistance of aquatic organisms to pathogens.

In the present study, we characterized the defensive capacities of epithelioma papulosum cyprini (EPC) cells and zebrafish chronically exposed to TFS and subsequently challenged with SVCV infection. To provide comprehensive mechanistic information useful for evaluating the ecological risks of TFS in aquatic ecosystems, the related responses involved in the initiation of autophagy in hosts were assessed at different levels of organization, from individual organisms to gene expression.

MATERIALS AND METHODS

Fungicide formulation

TFS in analytical standard (CAS Registry Number: 141517-21-7) was purchased from Sigma-Aldrich (USA) with 100% purity. The TFS solution was mixed with dimethyl sulfoxide (DMSO; Aladdin, China) to a stock concentration of 50 mg/mL, then stored at $-20\text{ }^{\circ}\text{C}$ until further use. To obtain the final concentration and ensure that TFS was fully dissolved and mixed in the dilution water, an appropriate amount of the applied solution was sonicated for approximately 10 min.

Cell culture, virus propagation, and zebrafish husbandry

The EPC SVCV-susceptible cell line was provided by Prof. Ling-Bing Zeng (Yangtze River Fisheries Research Institute, China) and cultured in medium 199 (M199, Hyclone, USA) supplemented with 10% fetal bovine serum (FBS; Every Green, China), penicillin (100 IU/mL), and streptomycin (0.1 mg/mL) at $25\text{ }^{\circ}\text{C}$ in a 5% CO_2 incubator for 48 h.

The SVCV strain (0504) provided by Prof. Qiang Li (Key Laboratory of Mariculture, Agriculture Ministry, Dalian Ocean University, Dalian, China) was isolated from common carp and propagated in EPC cells until a terminal cytopathic effect (>75% of observed CPE) occurred (Chen et al., 2006). The virus in diseased cells was harvested and stored at -80°C until further use. After SVCV titration into cell-cultured 96-well plates for 48 h, analysis of viral titer with a 50% tissue culture infective dose (TCID_{50}) was performed according to the Reed-Muench method (Pizzi, 1950).

Zebrafish ($n=2\ 500$, total length of 1.98 ± 0.22 cm, body weight of 49.89 ± 14.10 mg, mean \pm standard deviation (SD)) were purchased from the Qing-Feng Zebrafish Breeding Center (Shanghai, China), and maintained in five static water systems consisting of six 200 L aquarium ponds at 20°C . Standard dilution water (ISO7346/3-1998) was obtained from a sterile circulating water system. Photoperiod was maintained on a constant 12:12 h (light: dark) cycle. Prior to the experiments, the fish were transferred into new containers and acclimatized at 15°C under laboratory conditions for two weeks and fed to apparent satiation three times a day (from 0800h to 2200h) with a diet of commercial dry pellets (0.1% body weight). Ten zebrafish were randomly selected to verify SVCV-free status, as described previously (Koutná et al., 2003), with no virus found in the experimental zebrafish. All procedures were performed according to the Experimental Animal Management Law of China and approved by the Animal Ethics Committee of Ningbo University (No.10300).

Antibodies

The antibody LC3B (#3868) was purchased from Cell Signaling Technology, Inc (USA). The antibodies p62 (ab109012), Beclin-1 (ab210498), mTOR (ab32028), and p-mTOR (S2448) (ab109268) were purchased from Abcam (USA). The antibody α -tubulin (AF0001 and AT819) and secondary antibodies, including horse radish peroxidase (HRP)-labeled goat anti-mouse IgG (H+L) (A0216), HRP-labeled goat anti-rabbit IgG (H+L) (A0208), Alexa Fluor 647-labeled goat anti-rabbit IgG (H+L) (A0468), and Alexa Fluor 488-labeled goat anti-mouse IgG (H+L) (A0428), were purchased from Beyotime (China).

Cytotoxicity of TFS to EPC cells

Cytotoxicity analyses were performed using Cell Counting Kit-8 (CCK-8) (Beyotime, China) based on a water-soluble tetrazolium dye (WST-8), which is more sensitive than the conventional 3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium (MTS)-based assay. Briefly, serial dilutions of TFS at concentrations of $500/2^n$ $\mu\text{g/L}$ ($n=0-6$) in triplicate were added to cells cultured on a 96-well plate in M199 supplemented with 5% FBS. At each 24 h period, cell viability following TFS exposure was tested as per the manufacturer's protocols for CCK-8. Colorimetric absorbance after the mitochondrial enzyme substrate reaction was measured at 450 nm using a microplate reader (ELX800, Gene, Hong Kong, China). As DMSO exhibits no cytotoxicity at doses of up to 0.1% (v/v) (Liu et al., 2020a), TFS-exposed signals were compared to solvent-control signals (DMSO up to 0.001%, v/v).

Virus susceptibility in EPC cells

To explore the negative influence of TFS pre-exposure on virus susceptibility, prior to SVCV infection, the EPC cells seeded on 12-well plates were exposed to TFS at 2.5 $\mu\text{g/L}$ and 25 $\mu\text{g/L}$ for 1, 3, 5, and 7 d. Afterwards, the EPC cells were washed with 0.1 mol/L phosphate-buffered saline (PBS, pH=7.4; Beyotime, China) and infected with SVCV at $10^3\times\text{TCID}_{50}/\text{mL}$ over 48 h. The expression of SVCV nucleoprotein was detected by quantitative real-time polymerase chain reaction (qRT-PCR). The solution was replaced with fresh TFS diluent every 24 h to ensure that actual concentrations deviated little from initial target concentrations.

Simultaneously, the antiviral effects of TFS were monitored as described previously (Liu et al., 2017). Briefly, EPC cells were seeded on 12-well plates to a monolayer and infected with SVCV at $10^3\times\text{TCID}_{50}/\text{mL}$ at 25°C , followed by the addition of TFS at 2.5 $\mu\text{g/L}$ and 25 $\mu\text{g/L}$. After 48 h of infection, the antiviral effects of TFS on SVCV replication in EPC cells were tested by qRT-PCR.

Cell ultrastructural observations

The sampled cells were harvested and maintained in 2.5% glutaraldehyde at 4°C overnight. The sampled cells were prepared for transmission electron microscopy as described in an earlier study (Zhang et al., 2011). Ultrathin cell sections were observed under a 1200EX transmission electron microscope (TEM) (JEOL, Japan).

Virus susceptibility in zebrafish

A total of 225 healthy zebrafish were randomly chosen and distributed into three groups (DMSO-control and TFS at 2.5 $\mu\text{g/L}$ and 25 $\mu\text{g/L}$ (75 zebrafish in each) containing ultraviolet (UV)-sterilized water). To overcome the lack of samples due to the accidental death of zebrafish, there were 25 fish per tank (20 zebrafish sampled) and three tanks per treatment. The rearing temperature of each aquarium was kept at 15°C . The experimental fish were exposed to TFS at 2.5 $\mu\text{g/L}$ and 25 $\mu\text{g/L}$ for 1, 3, 5, 7, and 14 d, followed by immersion with SVCV at $10^3\times\text{TCID}_{50}/\text{mL}$ (M199 served as the control). At each sampling time, four zebrafish in each replicate were euthanized using a washrag soaked with 40 $\mu\text{g/mL}$ MS-222 (3-aminobenzoic acid ethyl ester methanesulfonate, CAS Registry Number: 886-86-2; Sigma, USA), with samples frozen for later analysis. Subsequently, samples were thawed and homogenized with liquid nitrogen to extract total RNA for detection of SVCV nucleoprotein expression. During exposure, the solutions were replaced with fresh TFS diluent every 24 h to ensure that actual concentrations deviated little from initial target concentrations and to maintain dissolved oxygen concentrations. The same process of TFS exposure was repeated without SVCV infection. Total RNA and protein in the sampled zebrafish were isolated to determine the expression of autophagy-related genes and analyze antioxidant enzyme activity, respectively.

Total RNA extraction, viral load quantification, and qRT-PCR

The samples of EPC cells and zebrafish were immediately

frozen in liquid nitrogen for subsequent RNA isolation. Total RNA was extracted using TRIZOL reagent (Takara, Japan) following the manufacturer's instructions, and eluted in 10–40 μL nuclease-free water for storage at $-80\text{ }^{\circ}\text{C}$. Nucleic acid concentrations were quantified using a NanoDrop spectrophotometer (ThermoFisher, USA). Extracted RNA purity was determined using the A260 nm/A280 nm ratio, with absorbance at 260 and 280 nm between 1.8 and 2.0. In total, 500 ng/ μL of purified RNA was used for cDNA generation per reaction, which was reverse transcribed using HiScript Q Select RT SuperMix for qPCR (Vazyme, China). qRT-PCR was performed using ChamQ™ SYBR® qPCR Green Master Mix (Vazyme, China) in an ABI StepOnePlus™ Real-Time PCR Detection System (ThermoFisher, USA). PCR analysis was amplified as follows: 95 $^{\circ}\text{C}$ for 30 s, denaturation for 40 cycles at 95 $^{\circ}\text{C}$ for 15 s, and annealing at 60 $^{\circ}\text{C}$ for 60 s. To assess the specificity of each amplicon, melt curve analysis was also performed at the end of each thermal profile. The sequences of primers are listed in Table 1 (Bello-Perez et al., 2020; Gotesman et al., 2015; Varela et al., 2014). Each sample was run in triplicate and the β -actin gene was used as an internal control for normalization. Relative expression was determined as fold induction relative to the expression level in the positive control (without TFS exposure) (set to 1). Relative mRNA expression levels were calculated using the $2^{-\Delta\Delta\text{Ct}}$ method with the formula $\Delta\Delta\text{Ct}_i = (\text{C}_{t, \text{target gene}} - \text{C}_{t, \text{reference gene}}) - (\text{C}_{t, \text{target gene}} - \text{C}_{t, \text{reference gene}})_{\text{control}}$ (Livak & Schmittgen, 2001).

Western blot assays

Total protein in the EPC cells and zebrafish was extracted using RIPA Lysis Buffer (Beyotime, China) under ice-bath cooling according to the manufacturer's instructions. The homogenate was centrifuged at 8 000 g for 15 min at 4 $^{\circ}\text{C}$ to obtain the protein supernatant. For western blot assays, total protein (isolated using a Nuclear and Cytoplasmic Protein Extraction Kit, Beyotime, China) was separated via 12% sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) and then transferred to polyvinylidene fluoride (PVDF) membranes (0.22 μm). The blots were blocked for 2 h at room temperature with 5% bovine serum albumin (BSA) in tris-buffered saline containing 0.1% Tween 20 (TBST) and incubated with primary antibodies (1:1 000) at 4 $^{\circ}\text{C}$ overnight. After TBST washing, the blots were incubated with secondary antibodies (1:1 000) at room temperature for 1 h.

Immunofluorescence

The EPC cells were exposed to TFS for 1, 3, 5, and 7 d, as described above. The TFS-exposed cells were fixed with 4% paraformaldehyde at 4 $^{\circ}\text{C}$ overnight and permeabilized with Triton X100 (Beyotime, China) for 30 min at room temperature. Subsequently, the cells were incubated with specific primary antibodies (LC3B, 1:250; α -tubulin, 1:500) at 4 $^{\circ}\text{C}$ overnight. After incubation with Alexa Fluor 488/647-labeled secondary antibody (1:1 000) at 37 $^{\circ}\text{C}$ for 1 h, fluorescence signals were examined using an upright fluorescence microscope (Nikon, Japan).

Statistical analysis

Analysis of variance (ANOVA) and Tukey's *post hoc* tests

Table 1 Sequences of primer pairs used for analysis of gene expression by qRT-PCR

Gene		Primer sequence (from 5' to 3')
β -actin (EPC cells)	Forward	GCTATGTGGCTCTTGACTTCGA
	Reverse	CCGTCAGGCAGCTCATAGCT
SVCV nucleoprotein (N)	Forward	AACAGCGCGTCTTACATGC
	Reverse	CTAAGGCGTAAGCCATCAGC
ambra1 (zebrafish)	Forward	TCTTTCGAGAAATGGCACCT
	Reverse	CTCTCTGCGTTAGGGACAGG
wipi1 (zebrafish)	Forward	GTGAGAGGGTAGAGAACAG
	Reverse	GTAACAACGACCCAACATC
atg5 (zebrafish)	Forward	AGAGAGGCAGAACCTACTATC
	Reverse	CCTCGTGTCAAACCACATTTT
gabarap (zebrafish)	Forward	GTCTGACCTCACAGTTGGGC
	Reverse	TCCTGGTAGAGCAGTCCCAT
18S (zebrafish)	Forward	ACCACCCACAGAATCGAGAAA
	Reverse	GCCTGCGGCTTAATTTGACT

were used to compare TFS-exposed samples with DMSO-controls to determine significance (SPSS 18.0, USA). Results are presented as mean \pm SD and $P < 0.05$ was considered statistically significant.

RESULTS

Cytotoxicity of TFS to EPC cells

The potential cytotoxicity of TFS was quantitatively assessed. As shown in Figure 1, the data indicated no evident cytotoxicity in EPC cells exposed to concentrations of TFS up to 125 $\mu\text{g}/\text{L}$ for 4 d, whereas cytotoxicity was observed after exposure to 250 $\mu\text{g}/\text{L}$ for 4 d and 500 $\mu\text{g}/\text{L}$ for 1–4 d. It should be noted that TFS was not renewed in the 96-well plates during exposure. These findings suggest that TFS at concentrations (2.5 $\mu\text{g}/\text{L}$ and 25 $\mu\text{g}/\text{L}$) used in subsequent experiments did not affect EPC cell growth.

Effects of TFS on SVCV replication in EPC cells

To determine whether environmental concentrations of TFS impact viral replication, we evaluated the effects of TFS at 2.5 $\mu\text{g}/\text{L}$ and 25 $\mu\text{g}/\text{L}$ on SVCV replication. As shown in Figure 2A, TFS at 2.5 $\mu\text{g}/\text{L}$ and 25 $\mu\text{g}/\text{L}$ (without renewal during exposure) significantly up-regulated (3-fold) the expression of the SVCV nucleoprotein at 48 h post-infection (hpi), suggesting that TFS may enhance viral infection in fish cells. TFS affected SVCV infectivity without acting on the virions as there was no obvious change in SVCV replication during co-incubation of TFS and SVCV (Figure 2B). Based on these data, we analyzed the longer-term effects of TFS on SVCV replication. As shown in Figure 2C–F, up-regulation occurred at 1, 3, 5, and 7 d pre-exposure, with a nearly 40-fold increase in the highest dose group. These results suggest that TFS may have an adverse effect on the resistance of fish cells to SVCV infection.

TFS induced autophagy in EPC cells

Based on earlier studies showing that TFS induces mitophagy in human skin keratinocytes and SVCV induces autophagy




Figure 1 Cytotoxicity of TFS in EPC cells

Cytotoxicity data for both TFS and DMSO are shown. No TFS cytotoxicity was detected up to 125 µg/L over 4 d of exposure. Each value represents mean±SD, normalized to values for no treatment. Statistical significance for each study was determined between Control_{DMSO} and TFS treatments by ANOVA with Tukey's *post hoc* tests. **: $P < 0.01$; *: $P < 0.05$. $n = 4$.

necessary for viral replication (Jang et al., 2016; Liu et al., 2015), we hypothesized that TFS may regulate SVCV replication via autophagy in fish. Autophagy is characterized by the formation of double-membrane vesicles (autophagosomes) that engulf cellular cargo and degrade internal contents upon fusion with endolysosomes (Liu et al., 2015). Results obtained via TEM showed that the number of autophagic vesicles with cytosolic contents increased in the cytoplasm of TFS-exposed EPC cells, whereas vesicles were rarely observed in the control cells (Figure 3). To further investigate the effects of TFS on autophagy in EPC cells, we examined LC3 lipidation, which is regarded as a significant indicator of autophagy activation, using western blot and immunofluorescence techniques. After TFS exposure, the conversion of LC3-I to LC3-II increased significantly, thus reflecting an increase in the lipidated form compared to that in the DMSO control (Figure 4A). Furthermore, a similar effect was observed during immunocytochemical testing, in which a specific antibody against LC3 was observed in untreated and viral-infected cells. A decrease in puncta number was found in the DMSO-exposed cells compared with TFS exposure,




Figure 2 Antiviral effects of TFS *in vitro*

A: Expression of SVCV nucleoprotein (N) increased significantly in TFS-treated groups. B: TFS had no effect on SVCV infection *in vitro* by altering virus binding to host cells. TFS at 25 µg/L was preincubated with 1×10^3 TCID₅₀/mL SVCV for 15–60 min. C–F: SVCV replication was detected in TFS pre-exposed cells at 1, 3, 5, and 7 d. EPC cells were pre-exposed to 2.5 µg/L and 25 µg/L TFS for up to 7 d, followed by SVCV infection at each time point. SVCV loads were determined via qRT-PCR analysis of N gene expression. Each value represents mean±SD normalized to values for no treatment. *P*-value for each study was determined by ANOVA with Tukey's *post hoc* tests. **: $P < 0.01$; *: $P < 0.05$; ns: No significance. $n = 4$ for A–B, and $n = 8$ for C–F.




Figure 3 Ultrastructural features of TFS-exposed cells observed via transmission electron microscopy (TEM)

Red arrows point to autophagosomes.

indicating that TFS activated autophagy in EPC cells at 1, 3, 5, and 7 d (Figure 4B–E).

Autophagy-related representative protein expression in TFS-exposed cells

Autophagy induced by TFS in EPC cells was further assessed and confirmed by western blot analysis of the protein levels of several essential components. TFS altered these protein levels to some extent (Figure 5), and autophagy substrate P62/SQSTM1 decreased upon autophagy activation by TFS. Of note, TFS reduced Beclin-1 content at 1 d, but there were no obvious changes at 3, 5, or 7 d. Additionally, to investigate the mechanism by which autophagy was induced by TFS exposure, we investigated the involvement of the mTOR protein. Results showed that the expression of mTOR was inhibited by TFS, and the inhibition effect intensified at longer exposure times. As mTOR protein activation is dependent on its phosphorylation level, we also assessed pmTOR levels in EPC cells. As expected, TFS reduced mTOR phosphorylation at Ser2448 at 5 and 7 d. The change trend of mTOR-pmTOR corresponded to SVCV replication in the TFS pre-exposed cells. These results suggest that TFS facilitated autophagy in EPC cells, which increased SVCV replication.

Promotion of SVCV replication in zebrafish by TFS-induced autophagy

Based on *in vitro* pre-exposure data, we explored the ability of TFS to enhance SVCV infection *in vivo*. Zebrafish were exposed to TFS for 1, 3, 5, 7, and 14 d, followed by immersion infection with SVCV. Except for day 1, the SVCV-infected fish from the TFS-exposed groups (3 d and 5 d) exhibited significantly higher viral replication than the DMSO-exposed SVCV-infected fish (Figure 6A). Moreover, the virus increased dramatically (more than a hundred-fold) following TFS exposure for 7 and 14 d (Figure 6B, C). These results provide further evidence that TFS promotes SVCV proliferation in zebrafish.

Autophagy-related gene and protein expression levels in TFS-exposed zebrafish

To determine whether TFS-exposed zebrafish exhibited abnormal autophagy activation, TFS-induced changes in the expression of four autophagy-related genes were analyzed. As expected, the expression levels of *gabarap*, *atg5*, *wipi5*, and *ambra1* were significantly up-regulated in the TFS pre-exposed fish (no virus infection) (Figure 7A). This finding is consistent with the data from the *in vitro* study showing that autophagy-related protein levels at 25 µg/L were obviously elevated compared with those in the negative control (Figure 7B). TFS reversed the up-regulation of LC3-II/LC3-I and down-regulation of P62 in zebrafish. Furthermore, TFS exposure resulted in a decrease in mTOR levels, suggesting that TFS may enhance SVCV infection in zebrafish via excessive autophagy.

DISCUSSION

Increased food production under pesticide use has led to the detection of high levels of pesticides in environmental matrices (Collotta et al., 2013), potentially resulting in the emergence and outbreak of different diseases in fish due to interactions among pathogen virulence, host resistance, and aquatic ecosystem stability. In fact, increasing evidence points to associations between pesticide exposure and a greater predisposition of aquatic organisms to circulating pathogens, as well as an increase in disease sensitivity and mortality in fish (Dietrich et al., 2014; Morley, 2010). For example, Chinook salmon exposed to malathion and challenged with *Aeromonas salmonicida* show an increase in mortality (Dietrich et al., 2014); exposure to polycyclic aromatic hydrocarbons alters resistance in Japanese flounder (*Paralichthys olivaceus*) to viral hemorrhagic septicemia virus (Song et al., 2011); and exposure of rainbow trout (*Oncorhynchus mykiss*) to pendimethalin suppresses

immunological responses and enhances infectious hematopoietic necrosis virus infections (Dupuy et al., 2019). Therefore, the adverse effects of pesticides on fish should be continuously investigated to protect aquatic environments as well as human health (Dupuy et al., 2019). In this study, we investigated the effects of TFS on antiviral defenses in zebrafish subjected to chronic *in vivo* TFS at environmentally relevant concentrations followed by SVCV challenge. Results showed a significant increase in viral replication in TFS-exposed fish cells and zebrafish. This could be attributed to autophagic benefits to SVCV replication by removing damaged mitochondrial DNA in SVCV-infected cells (Liu et al., 2015).

Autophagy is an evolutionarily conserved membrane-trafficking process by which cells degrade old and damaged cytoplasmic components via lysosomes into basic





Figure 4 Expression of LC3B regulated by TFS in EPC cells

A: Western blot analyses of LC3B and LC3I/LC3II in EPC cells. B–E: Immunocytochemical results showing intracellular location and number of LC3B at 1, 3, 5, and 7 d, respectively.

biomolecules, thereby recycling cellular material. Autophagy therefore plays an important role in many crucial physiological processes such as cell differentiation and immunity in response to foreign substances and harmful influences (Dikic & Elazar, 2018; He & Klionsky, 2009). Autophagy is not only important for maintaining homeostasis in cells, but also in many pathogenic processes (Schmid et al., 2006; Zhang et al., 2011). The contrasting nature of autophagy has inspired much research. On the one hand, autophagy mediates both surveillance and effector functions involved in the detection and clearance of pathogens (Levine & Deretic, 2007; Shoji-Kawata & Levine, 2009) and is necessary for maintaining protein and organelle turnover. Conversely, some viruses, including SVCV, utilize the autophagy pathway to facilitate their own genomic replication and produce stronger infections (Liu et al., 2015; Orvedahl et al., 2007; Yoon et al., 2008). Numerous studies have shown that chemicals such as pesticides and metals can interfere with autophagy depending on cell type, exposure duration, and dose (Pesonen & Vähäkangas, 2019). Notably, elevated ROS levels can lead to mitochondrial dysfunction and activation of autophagy for the




Figure 5 Total amounts of Beclin-1, P62, and mTOR, and phosphorylation level of pmTOR in EPC cells, as analyzed by western blotting. Grayscale values were analyzed by Image J. Data are mean \pm SD of one representative experiment performed in triplicate. *P*-value for each study was determined by ANOVA with Tukey's *post hoc* tests. **: *P*<0.01; *: *P*<0.05.




Figure 6 SVCV replication in TFS pre-exposed zebrafish at 1–5 d (A), 7 d (B), and 14 d (C)

Each value represents mean \pm SD, normalized to values for no treatment. *P*-value for each study was determined by ANOVA with Tukey's *post hoc* tests. **: *P*<0.01; *: *P*<0.05. *n*=12 fish in each group.

elimination of dysfunctional mitochondria (Cheng et al., 2009). Interestingly, TFS targets mitochondrial respiration by blocking the flow between cytochrome *b* and cytochrome *c*1 at the ubiquinol oxidizing site (Qo) of complex III (Bartlett et al., 2002), which leads to damaged mitochondria. Evidence to date suggests that TFS can cause representative mitochondrial damage, such as altered membrane potential and oxidative stress, in zebrafish (Zhu et al., 2015a), and thus,

there is a possible connection between autophagy and TFS exposure. In this study, SVCV replication was associated with increased autophagy in TFS-exposed cells and higher numbers of autophagic vesicles were observed by TEM.

Autophagosome formation and maturation are tightly controlled and coordinated by specialized proteins (He et al., 2020), e.g., the Atg8/LC3 conjugation system and Beclin-1, which are related to autophagosome formation (Zhang et al.,




Figure 7 Alteration in autophagy-related gene and protein expression levels in TFS-exposed zebrafish

A: Expression of autophagy-related genes was changed by TFS exposure in zebrafish. Expression levels of four autophagy-related mRNA genes, i.e., *gabarap*, *atg5*, *wipi5*, and *ambra1*, were analyzed by qRT-PCR. B: Total amount of autophagy-related proteins in zebrafish were analyzed by western blotting. Grayscale values were analyzed by Image J. Experiments were performed in triplicate, and each value represents mean \pm SD. *P*-value for each study was determined by ANOVA with Tukey's *post hoc* tests. **: *P*<0.01; *: *P*<0.05. *n*=12 fish in each group.

2011), and P62 (known as sequestosome 1, SQSTM1), which is correlated with protein degradation in autophagy (Liu et al., 2015). Our western blot assays confirmed the activation of autophagy by TFS, as indicated by increased Beclin-1 and LC3I/LC3II along with decreased P62 after EPC cells and zebrafish were exposed to TFS. mTOR is one of the best-known cytoplasmic sensors and regulators of the autophagy pathway (Qin et al., 2010), and belongs to the Ser/Thr kinases, including rapamycin-sensitive mTORC1 and non-sensitive mTORC2. Specifically, mTORC1 negatively regulates autophagosome formation by regulating the phosphorylation level of the ULK1-Atg13-RB1CC1-C12ORF44/Atg101 complex (Wan et al., 2014), while mTORC2 inhibits autophagy by activating the Atg1 protein (Cheng et al., 2013). The activation of mTOR can block the expression and function of autophagy-related (ATG) proteins. Liu et al. (2015) determined that SVCV causes a significant reduction in pmTOR, which facilitates the replication of mTOR signals in fish. Our work suggests that exposure to TFS can result in the activation of autophagy in fish by inhibition of mTOR signaling, which enhances SVCV susceptibility in zebrafish.

In conclusion, we explored the aquatic environmental risks of TFS by investigating the cellular and molecular responses of zebrafish following exposure to TFS, with subsequent SVCV challenge. Our findings demonstrated non-significant cytotoxicity of TFS up to 125 µg/L in EPC cells. The significant enhancement of SVCV replication in TFS-exposed EPC cells

and zebrafish was associated with activation of autophagy. This suggests that TFS in aquatic environments may contribute to SVCV outbreaks and endanger ecological safety. At present, however, we are unable to determine the upstream regulatory mechanism of autophagy in mTOR signaling and whether there is permanent damage in fish following TFS exposure. This topic will be the focus of future research.

COMPETING INTERESTS

The authors declare that they have no competing interests.

AUTHORS' CONTRIBUTIONS

J.C. and L.L. drafted the experiments; H.W., H.W.L., L.H., and J.F.L. performed the experiments. T.X.Q. and L.L. analyzed the data and wrote the paper. All authors read and approved the final version of the manuscript.

REFERENCES

- Ahne W, Bjorklund HV, Essbauer S, Fijan N, Kurath G, Winton JR. 2002. Spring viremia of carp (SVC). *Diseases of Aquatic Organisms*, **52**(3): 261–272.
- Balmer BF, Powers RL, Zhang TH, Lee J, Vigant F, Lee B, et al. 2017. Inhibition of an aquatic rhabdovirus demonstrates promise of a broad-spectrum antiviral for use in aquaculture. *Journal of Virology*, **91**(4): e02181–16.
- Bartlett DW, Clough JM, Godwin JR, Hall AA, Hamer M, Parr-Dobrzanski B. 2002. The strobilurin fungicides. *Pest Management Science*, **58**(7):

649–662.

- Bello-Perez M, Pereira P, Coll J, Novoa B, Perez L, Falco A. 2020. Zebrafish C-reactive protein isoforms inhibit SVCV replication by blocking autophagy through interactions with cell membrane cholesterol. *Scientific Reports*, **10**(1): 566.
- Cao MC, Li SY, Wang QS, Wei P, Liu YN, Zhu GN, et al. 2015. Track of fate and primary metabolism of trifloxystrobin in rice paddy ecosystem. *Science of the Total Environment*, **518–519**: 417–423.
- Chen ZY, Liu H, Li ZQ, Wang M, Zhang QY. 2006. Detection of viral pathogen from diseased common carp (*Cyprinus carpio*) by infectious tests. *Journal of Fishery Sciences of China*, **13**(4): 617–623. (in Chinese)
- Cheng P, Ni Z, Dai X, Wang B, Ding W, Smith AR, et al. 2013. The novel BH-3 mimetic apogossypolone induces Beclin-1- and ROS-mediated autophagy in human hepatocellular carcinoma cells. *Cell Death & Disease*, **4**(2): e489.
- Cheng Y, Azad MB, Gibson SB. 2009. Superoxide is the major reactive oxygen species regulating autophagy. *Cell Death & Differentiation*, **16**(7): 1040–1052.
- Collotta M, Bertazzi PA, Bollati V. 2013. Epigenetics and pesticides. *Toxicology*, **307**: 35–41.
- Cui F, Chai TT, Liu XX, Wang CJ. 2017. Toxicity of three strobilurins (Kresoxim-methyl, pyraclostrobin, and trifloxystrobin) on *Daphnia magna*. *Environmental Toxicology and Chemistry*, **36**(1): 182–189.
- Deb D, Engel BA, Harbor J, Hahn L, Lim KJ, Zhai T. 2010. Investigating potential water quality impacts of fungicides used to combat soybean rust in Indiana. *Water, Air, and Soil Pollution*, **207**(1–4): 273–288.
- Dietrich JP, Van Gaest AL, Strickland SA, Arkoosh MR. 2014. The impact of temperature stress and pesticide exposure on mortality and disease susceptibility of endangered Pacific salmon. *Chemosphere*, **108**: 353–359.
- Dikic I, Elazar Z. 2018. Mechanism and medical implications of mammalian autophagy. *Nature Reviews Molecular Cell Biology*, **19**(6): 349–364.
- Dikkeboom AL, Radi C, Toohey-Kurth K, Marcquenski S, Engel M, Goodwin AE, et al. 2004. First report of spring viremia of carp virus (SVCV) in wild common carp in North America. *Journal of Aquatic Animal Health*, **16**(4): 169–178.
- Dupuy C, Cabon J, Louboutin L, Le Floch S, Morin T, Danion M. 2019. Cellular, humoral and molecular responses in rainbow trout (*Oncorhynchus mykiss*) exposed to a herbicide and subsequently infected with infectious hematopoietic necrosis virus. *Aquatic Toxicology*, **215**: 105282.
- European Food Safety Authority (EFSA), Arena M, Auteri D, Barnaz S, Bellisai G, Brancato A, et al. 2017. Peer review of the pesticide risk assessment of the active substance trifloxystrobin. *FESA Journal*, **15**(10): e04989.
- Feng YM, Huang YH, Zhan H, Bhatt P, Chen SH. 2020. An overview of strobilurin fungicide degradation: current status and future perspective. *Frontier in Microbiology*, **11**: 389.
- Garver KA, Dwilow AG, Richard J, Booth TF, Beniac DR, Souter BW. 2007. First detection and confirmation of spring viraemia of carp virus in common carp, *Cyprinus carpio* L., from Hamilton Harbour, Lake Ontario, Canada. *Journal of Fish Diseases*, **30**(11): 665–671.
- Gotesman M, Soliman H, Besch R, El-Matbouli M. 2015. Inhibition of spring viraemia of carp virus replication in an *Epithelioma papulosum* cyprini cell line by RNAi. *Journal of Fish Diseases*, **38**(2): 197–207.
- He BN, Wang X, Yang CL, Zhu JB, Jin YX, Fu ZW. 2020. The regulation of autophagy in the pesticide-induced toxicity: Angel or demon?. *Chemosphere*, **242**: 125138.
- He CC, Klionsky DJ. 2009. Regulation mechanisms and signaling pathways of autophagy. *Annual Review of Genetics*, **43**: 67–93.
- Jang Y, Kim JE, Jeong SH, Paik MK, Kim JS, Cho MH. 2016. Trifloxystrobin-induced mitophagy through mitochondrial damage in human skin keratinocytes. *The Journal of Toxicological Sciences*, **41**(6): 731–737.
- Junges CM, Peltzer PM, Lajmanovich RC, Attademo AM, Cabagna Zenklusen MC, Basso A. 2012. Toxicity of the fungicide trifloxystrobin on tadpoles and its effect on fish-tadpole interaction. *Chemosphere*, **87**: 1348–1354.
- Komárek M, Čadková E, Chrastrný V, Bordas F, Bollinger JC. 2010. Contamination of vineyard soils with fungicides: a review of environmental and toxicological aspects. *Environment International*, **36**(1): 138–151.
- Koutná M, Veselý T, Psikal I, Hůlová J. 2003. Identification of spring viraemia of carp virus (SVCV) by combined RT-PCR and nested PCR. *Diseases of Aquatic Organisms*, **55**(3): 229–235.
- Levine B, Deretic V. 2007. Unveiling the roles of autophagy in innate and adaptive immunity. *Nature Reviews Immunology*, **7**(10): 767–777.
- Li H, Cao FJ, Zhao F, Yang Y, Teng MM, Wang CJ, et al. 2018. Developmental toxicity, oxidative stress and immunotoxicity induced by three strobilurins (pyraclostrobin, trifloxystrobin and picoxystrobin) in zebrafish embryos. *Chemosphere*, **207**: 781–790.
- Liu L, Hu Y, Shen YF, Wang GX. 2017. Evaluation on antiviral activity of coumarin derivatives against spring viraemia of carp virus in epithelioma papulosum cyprini cells. *Antiviral Research*, **144**: 173–185.
- Liu L, Jiang C, Wu ZQ, Gong YX, Wang GX. 2013. Toxic effects of three strobilurins (trifloxystrobin, azoxystrobin and kresoxim-methyl) on mRNA expression and antioxidant enzymes in grass carp (*Ctenopharyngodon idella*) juveniles. *Ecotoxicology and Environmental Safety*, **98**: 297–302.
- Liu L, Song DW, Liu GL, Shan LP, Qiu TX, Chen J. 2020a. Hydroxycoumarin efficiently inhibits spring viraemia of carp virus infection *in vitro* and *in vivo*. *Zoological Research*, **41**(4): 395–409.
- Liu LY, Zhu BB, Wu SS, Lin L, Liu GX, Zhou Y, et al. 2015. Spring viraemia of carp virus induces autophagy for necessary viral replication. *Cellular Microbiology*, **17**(4): 595–605.
- Liu T, Liu YL, Fang K, Zhang XL, Wang XG. 2020b. Transcriptome, bioaccumulation and toxicity analyses of earthworms (*Eisenia fetida*) affected by trifloxystrobin and trifloxystrobin acid. *Environmental Pollution*, **265**: 115100.
- Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta C_t}$ method. *Methods*, **25**(4): 402–408.
- Luo XS, Qin XX, Liu ZY, Chen D, Yu WW, Zhang KK, et al. 2020. Determination, residue and risk assessment of trifloxystrobin, trifloxystrobin acid and tebuconazole in Chinese rice consumption. *Biomedical Chromatography*, **34**(1): e4694.
- Ma JG, Zhu JY, Wang WY, Ruan PP, Rajeshkumar S, Li XY. 2019. Biochemical and molecular impacts of glyphosate-based herbicide on the gills of common carp. *Environmental Pollution*, **252**: 1288–1300.
- Morley NJ. 2010. Interactive effects of infectious diseases and pollution in aquatic molluscs. *Aquatic Toxicology*, **96**(1): 27–36.
- Orvedahl A, Alexander D, Tallóczy Z, Sun QH, Wei YJ, Zhang W, et al. 2007. HSV-1 ICP34.5 confers neurovirulence by targeting the Beclin 1 autophagy protein. *Cell Host & Microbe*, **1**(1): 23–35.
- Pesonen M, Vähäkangas K. 2019. Autophagy in exposure to environmental chemicals. *Toxicology Letters*, **305**: 1–9.
- Pizzi M. 1950. Sampling variation of the fifty percent end-point, determined

- by the Reed-Muench (Behrens) method. *Human Biology*, **22**(3): 151–190.
- Qin L, Wang Z, Tao LY, Wang Y. 2010. Er stress negatively regulates AKT/TSC/mTOR pathway to enhance autophagy. *Autophagy*, **6**(2): 239–247.
- Schmid D, Dengjel J, Schoor O, Stevanovic S, Münz C. 2006. Autophagy in innate and adaptive immunity against intracellular pathogens. *Journal of Molecular Medicine*, **84**(3): 194–202.
- Shoji-Kawata S, Levine B. 2009. Autophagy, antiviral immunity, and viral countermeasures. *Biochimica et Biophysica Acta (BBA)-Molecular Cell Research*, **1793**(9): 1478–1484.
- Song JY, Nakayama K, Murakami Y, Kitamura SI. 2011. Heavy oil exposure induces high mortalities in virus carrier Japanese flounder *Paralichthys olivaceus*. *Marine Pollution Bulletin*, **63**(5–12): 362–365.
- Taylor NGH, Peeler EJ, Denham KL, Crane CN, Thrush MA, Dixon PF, et al. 2013. Spring viraemia of carp (SVC) in the UK: the road to freedom. *Preventive Veterinary Medicine*, **111**(1–2): 156–164.
- Varela M, Romero A, Dios S, Van der Vaart M, Figueras A, Meijer AH, et al. 2014. Cellular visualization of macrophage pyroptosis and interleukin-1 β release in a viral hemorrhagic infection in zebrafish larvae. *Journal of Virology*, **88**(20): 12026–12040.
- Wan G, Xie WD, Liu ZY, Xu W, Lao YZ, Huang N, et al. 2014. Hypoxia-induced *MIR155* is a potent autophagy inducer by targeting multiple players in the MTOR pathway. *Autophagy*, **10**(1): 70–79.
- Wang C, Wu JX, Zhang Y, Wang K, Zhang HY. 2015. Field dissipation of trifloxystrobin and its metabolite trifloxystrobin acid in soil and apples. *Environmental Monitoring and Assessment*, **187**(1): 4100.
- Wang K, Sun ZH, Yang LD, He L, Li XH, Wang G. 2020. Respiratory toxicity of azoxystrobin, pyraclostrobin and coumoxystrobin on *Chlorella vulgaris*. *Bulletin of Environmental Contamination and Toxicology*, **104**(6): 799–803.
- Wightwick AM, Bui AD, Zhang P, Rose G, Allinson M, Myers JH, et al. 2012. Environmental fate of fungicides in surface waters of a horticultural-production catchment in southeastern Australia. *Archives of Environmental Contamination and Toxicology*, **62**(3): 380–390.
- Xing HJ, Liu T, Zhang ZW, Wang XL, Xu SW. 2015. Acute and subchronic toxic effects of atrazine and chlorpyrifos on common carp (*Cyprinus carpio* L.): immunotoxicity assessments. *Fish & Shellfish Immunology*, **45**(2): 327–333.
- Xiong JB, Nie L, Chen J. 2019. Current understanding on the roles of gut microbiota in fish disease and immunity. *Zoological Research*, **40**(2): 70–76.
- Yoon SY, Ha YE, Choi JE, Ahn J, Lee H, Kweon HS, et al. 2008. Coxsackievirus B4 uses autophagy for replication after calpain activation in rat primary neurons. *Journal of Virology*, **82**(23): 11976–11978.
- Yuan JF, Yang Y, Nie HH, Li LJ, Gu WG, Lin L, et al. 2014. Transcriptome analysis of epithelioma papulosum cyprini cells after SVCV infection. *BMC Genomics*, **15**: 935.
- Zhang YN, Li ZC, Ge XN, Guo X, Yang HC. 2011. Autophagy promotes the replication of encephalomyocarditis virus in host cells. *Autophagy*, **7**(6): 613–628.
- Zhu B, Liu GL, Liu L, Ling F, Wang GX. 2015a. Assessment of trifloxystrobin uptake kinetics, developmental toxicity and mRNA expression in rare minnow embryos. *Chemosphere*, **120**: 447–455.
- Zhu LF, Wang HL, Liu HJ, Li W. 2015b. Effect of trifloxystrobin on hatching, survival, and gene expression of endocrine biomarkers in early life stages of medaka (*Oryzias latipes*). *Environmental Toxicology*, **30**(6): 648–655.