

Letter to the editor

Open Access

Sox9 is indispensable for testis differentiation in the red-eared slider turtle, a reptile with temperature-dependent sex determination

SOX9 (SRY-related HMG box gene 9) is an essential regulator of male sex determination and testis differentiation in many vertebrate species. However, the functional role of Sox9 in testis differentiation has not yet been identified in any reptilian species. Herein, Sox9 knockdown in the red-eared slider turtle (*Trachemys scripta*) embryos at a male-producing temperature led to complete male-to-female sex reversal, characterized by the formation of an ovary-like structure, disappearance of male marker AMH, and ectopic expression of ovarian regulator FOXL2, as well as a female distribution pattern of germ cells. Conversely, *in-ovo* overexpression of Sox9 at a female-producing temperature resulted in partial masculinization of putative female embryos, with the co-existence of AMH and FOXL2. Our study provides the first direct evidence that Sox9 is indispensable for testicular differentiation in a reptilian species, further confirming the conserved role of Sox9 in vertebrate sexual development.

Many egg-laying reptiles without sex chromosomes exhibit temperature-dependent sex determination (TSD), in which sex is established by the incubation temperature during the temperature-sensitive period (Pewphong et al., 2021). Sox9 is a member of the SRY-related HMG box (SOX) gene family and plays a conserved role in vertebrate sexual development. The expression patterns of Sox9 during embryonic gonadal development have been studied in several TSD species, which suggest the presence of Sox9 expression dimorphism between male-producing (MPT) and female-producing temperatures (FPT) (Díaz-Hernández et al., 2020; Moreno-Mendoza et al., 2001; Rhen et al., 2007; Shoemaker et al., 2007b; Spotila et al., 1998; Torres-Maldonado et al., 2001; Torres Maldonado et al., 2002; Western et al., 1999). In the red-eared slider turtle (*Trachemys scripta*), Sox9 expression in embryonic gonads is higher at the MPT than at the FPT in the

later temperature-sensitive period (Shoemaker et al., 2007a; Spotila et al., 1998). The expression of Sox9 is significantly down-regulated after exogenous estradiol treatment at the MPT, while inhibition of estrogen synthesis delays Sox9 down-regulation at the FPT in *T. scripta* (Barske & Capel, 2010; Matsumoto et al., 2013). In addition, knockdown of the male sex-determining gene *Dmrt1* in *T. scripta* results in the down-regulation of Sox9 expression, while overexpression of *Dmrt1* leads to an increase in Sox9 expression (Ge et al., 2017). These studies suggest that Sox9 may have a regulatory effect on sexual differentiation in turtles; however, its functional role has not yet been investigated in any TSD reptilian species.

In the current study, we investigated loss- and gain-of-function of Sox9 in the red-eared slider turtle using lentivirus-mediated genetic manipulation. A lentivirus carrying Sox9 short hairpin RNA (shRNA) or open reading frame (ORF) was injected into *T. scripta* embryos at stage 15, i.e., very beginning of the temperature-sensitive period (Matsumoto & Crews, 2012). Stage 25 gonads were dissected for quantitative real-time polymerase chain reaction (qRT-PCR), hematoxylin-eosin (H&E) staining, and immunofluorescence. Detailed procedures are described in the Supplementary Materials and Methods.

Immunofluorescence analysis of the control group showed that the SOX9 protein was mainly located in the nucleus of Sertoli precursor cells in MPT embryonic gonads, while no SOX9 signal was detected in the FPT embryonic gonads at stage 25 (Figure 1A). In the gonads of MPT embryos treated with LV-Sox9-shRNA, SOX9 decreased to an almost undetectable level, similar to that in the control FPT embryos (Figure 1A), thus indicating a strong inhibitory effect of LV-

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright ©2021 Editorial Office of Zoological Research, Kunming Institute of Zoology, Chinese Academy of Sciences

Received: 19 July 2021; Accepted: 18 September 2021; Online: 22 September 2021

Foundation items: This study was supported by the National Key Research and Development Program (2018YFD0900203), National Natural Science Foundation of China (31922084, 31872960), Natural Science Foundation of Zhejiang Province for Distinguished Young Scholars (LR19C190001), and Key Agricultural Project of Ningbo (2019C10018)




Figure 1 Complete sex reversal induced by Sox9 knockdown

A: Immunofluorescence of SOX9, representative images of gonad-mesonephros complexes (GMCs) and H&E staining of gonadal sections from control MPT, MPT+Sox9-RNAi, and control FPT embryos at stage 25. Gonads in GMCs are outlined by yellow dotted lines. Med: medulla, Cor: cortex. Scale bar for GMCs: 500 μ m, scale bar for immunofluorescence and H&E: 50 μ m. B: VASA and CTNNB1 immunostaining of gonadal sections from control MPT, MPT+Sox9-RNAi, and control FPT embryos at stage 25. Scale bar: 50 μ m. C: AMH and FOXL2 expression changes in response to Sox9 knockdown at stage 25. Scale bar: 50 μ m. D: Knockdown of Sox9 led to a high rate of male-to-female sex reversal in MPT embryos. Phenotype of embryonic gonads was evaluated by gonadal histological analysis and expression of sex-specific proteins (AMH and FOXL2).

Sox9-shRNA on Sox9 expression. Histological analysis showed that the MPT gonads had a dense medulla with seminiferous cords and a reduced cortex, while the FPT gonads showed a thickened outer cortex and vacuolated medulla (Figure 1A). In contrast, the Sox9 knockdown MPT gonads were strongly feminized, characterized by a highly developed cortical region and a significantly degraded medulla region, similar to that of the FPT gonads (Figure 1A). Immunofluorescence labeling showed that VASA-positive germ cells were distributed in the developed medulla of the

MPT gonads (Figure 1B). However, after Sox9 knockdown, the germ cells were mainly enriched in the outer cortex, exhibiting similar female germ cell distribution (Figure 1B). To confirm activation of the female developmental pathway in Sox9-deficient MPT embryos, we analyzed the expression of the male-specific protein AMH and ovarian development regulator FOXL2. Immunofluorescence analysis of the control group showed that AMH was strongly expressed in the Sertoli cells of the control MPT gonads but not in the control FPT gonads, while FOXL2 showed the opposite pattern and

primary expression in the cortex region of the FPT gonads (Figure 1C). After *Sox9* knockdown, the expression of AMH in most treated gonads was extremely low, while the expression of FOXL2 was robust, similar to that in normal FPT gonads (Figure 1C). Furthermore, knockdown of *Sox9* at the MPT led to a high rate (92%, 23/25) of complete male-to-female sex reversal (Figure 1D).

Sox9-overexpressing embryos were generated by an injection of a lentivirus vector carrying the *Sox9* ORF into FPT turtle eggs at stage 15. Immunofluorescence showed strong expression of the SOX9 protein in the medulla of FPT gonads after LV-*Sox9*-OE treatment (Figure 2A). Morphological analysis indicated that the FPT embryos overexpressing *Sox9* had thicker, shorter, and obviously masculinized gonads (Figure 2A). The H&E staining of gonadal sections showed typical male characteristics in some treated FPT gonads, including a degenerated cortex and well-developed medulla,

although some FPT gonads overexpressing *Sox9* still retained a developed cortex (Figure 2A). VASA staining indicated that germ cells were distributed in both the cortex and medulla after *Sox9* overexpression (Figure 2B). To confirm activation of the male developmental pathway in the *Sox9*-overexpressing FPT embryos, we analyzed the expression levels of testicular differentiation markers *Dmrt1* and *Amh* and ovarian development regulators *Cyp19a1* and *Foxl2*. The qRT-PCR results indicated that the mRNA expression levels of *Dmrt1* and *Amh* were significantly increased, while the expression levels of *Cyp19a1* and *Foxl2* were decreased in response to overexpression of *Sox9* (Figure 2D). In addition, based on immunofluorescence analysis, the ectopic expression of AMH was detected in the well-developed medulla in almost all cases, while the expression of female marker FOXL2 was markedly reduced (disappeared or limitedly retained) after overexpression of *Sox9* in the FPT




Figure 2 Partial female-to-male sex reversal induced by *Sox9* overexpression

A: Immunofluorescence of SOX9, representative images of GMCs and H&E staining of gonadal sections from control FPT, FPT+*Sox9*-OE, and control MPT embryos at stage 25. Gonads in GMCs are outlined by yellow dotted lines. Med: medulla, Cor: cortex. Scale bar for GMCs: 500 μ m, scale bar for immunofluorescence and H&E: 50 μ m. B: VASA and CTNNB1 immunostaining of gonadal sections from control FPT, FPT+*Sox9*-OE, and control MPT embryos at stage 25. Scale bar: 50 μ m. C: Responses of AMH and FOXL2 expression to *Sox9* overexpression at stage 25. Scale bar: 50 μ m. D: Expression changes in *Dmrt1*, *Amh*, *Foxl2*, and *Cyp19a1* after *Sox9* overexpression. Stage 25 gonads were dissected for qRT-PCR analysis. Results were normalized to *Gapdh*, and expression levels of MPT were defined as 1. Data were expressed as mean \pm standard deviation (SD) for three biological replicates. **: $P < 0.01$; ***: $P < 0.001$. E: Ectopic expression of *Sox9* resulted in partial female-to-male sex reversal.

gonads (Figure 2C). Gonads showing both AMH and FOXL2 expression were considered as ovotestes. In addition, 85% (34/40) of FPT embryos overexpressing *Sox9* underwent varying degrees of sex reversal, with eight individuals showing complete sex reversal and 26 individuals showing partial sex reversal (Figure 2E).

The sequences of the HMG and C-terminal reactivation domains of SOX9 are highly homologous across humans, chickens, alligators, and fish, suggesting a conserved functional role of *Sox9* in vertebrate gonadal differentiation (Da Silva et al., 1996; Vining et al., 2021). *Sox9* is essential for Sertoli cell differentiation and testis formation in mice (Chaboissier et al., 2004; Kent et al., 1996). In XY mice, the deletion or a 77% reduction of *Sox9* leads to complete male-to-female sex reversal, while 50% reduction causes partial sex reversal (Bagheri-Fam et al., 2020; Gonen et al., 2017; Lavery et al., 2011). Moreover, ectopic expression of *Sox9* in XX gonads causes male development in the absence of *Sry* (Huang et al., 1999; Vidal et al., 2001). In addition, the duplication or deletion of *Sox9* enhancers results in sex reversal in XX or XY individuals, respectively (Croft et al., 2018; Gonen et al., 2018), thus demonstrating its critical role in testis determination in mammals. *Sox9* is highly expressed in MPT gonads during sex differentiation in red-eared slider turtles (Barske & Capel, 2010; Shoemaker et al., 2007b). This, together with our results showing that knockdown or overexpression of *Sox9* can lead to partial or complete sex reversal in red-eared turtles, confirms that the conserved role of *Sox9* in testis differentiation is also present in reptilian species. Interestingly, gonad size decreased in the *Sox9*-deficient embryos, consistent with previous results reported in medaka (Nakamura et al., 2012). This may be due to the *Sox9* gene not only affecting sex determination, but also other biological processes such as cell proliferation and germ cell maintenance.

Mammalian *Sox9* is directly regulated by the master sex determining gene *Sry*, and its initial expression occurs immediately after that of *Sry* (Sekido & Lovell-Badge, 2008). *SRY* and NR5A1 (nuclear receptor subfamily 5 group A member 1, also named steroidogenic factor 1) together bind to the *Sox9* gonad-specific enhancer and cooperatively up-regulate *Sox9* expression (Bagheri-Fam et al., 2012; Sekido & Lovell-Badge, 2008). In mammals, *Sox9* is reported to activate *Amh* (anti-Müllerian hormone) during testicular development (Cutting et al., 2013; Da Silva et al., 1996; Yamashita et al., 2019). Unlike mammals, *Sox9* expression in chickens (*Gallus gallus domesticus*) and American alligators (*Alligator mississippiensis*) occurs later than that of *Amh*, indicating that *Amh* expression is not regulated by SOX9, and *Sox9* may be located downstream of the sex differentiation pathway in non-mammalian vertebrates (Hirst et al., 2018; Smith & Sinclair, 2004; Western et al., 1999). In teleosts, *Sox9* plays a role in ovary-testis transition and germ cell maintenance (Nakamura et al., 2012; Sun et al., 2013). These studies suggest that *Sox9* neofunctionalization likely occurred during vertebrate evolution, such that the specific regulatory position of SOX9 in the male sex differentiation pathway differs among vertebrates with different evolutionary statuses (Cutting et al., 2013). In the red-eared slider turtle, the sexually dimorphic expression

of *Sox9* appears later than that of *Dmrt1*, and knockdown of *Dmrt1* in MPT embryos results in down-regulation of *Sox9* (Ge et al., 2017). In fish, DMRT1 positively regulates the transcription of the *Sox9b* gene (ortholog of tetrapod *Sox9*) by directly binding to a specific cis-regulatory element within the *Sox9b* promoter (Wei et al., 2019). These results imply that *Sox9* may be involved in testis differentiation rather than sex determination in red-eared slider turtles and may be directly or indirectly regulated by *Dmrt1*.

Herein, we showed that knockdown of *Sox9* in *T. scripta* led to complete male-to-female sex reversal, while overexpression of *Sox9* caused partial female-to-male sex reversal, thus implicating the indispensable role of *Sox9* in testicular differentiation in reptiles. This study confirms the conserved role of *Sox9* in male sexual development across different vertebrate species.

SUPPLEMENTARY DATA

Supplementary data to this article can be found online.

COMPETING INTERESTS

The authors declare that they have no competing interests.

AUTHORS' CONTRIBUTIONS

C.T.G., H.B.H., and L.X. conceived and designed the study. H.B.H., L.X., W.S., Y.J.Z., and H.Y.Z. collected the samples and analyzed the data. H.B.H., L.X., and C.T.G. wrote the manuscript. All authors read and approved the final version of the manuscript.

Hang-Bo Hui^{1, #}, Ling Xiao^{1, #}, Wei Sun¹, Ying-Jie Zhou¹,
Hai-Yan Zhang¹, Chu-Tian Ge^{1, *}

¹ Institute of Animal Sex and Development, Zhejiang Wanli University, Ningbo, Zhejiang 315100, China

[#]Authors contributed equally to this work

*Corresponding author, E-mail: cge@zwu.edu.cn

REFERENCES

- Bagheri-Fam S, Combes AN, Ling CK, Wilhelm D. 2020. Heterozygous deletion of *Sox9* in mouse mimics the gonadal sex reversal phenotype associated with campomelic dysplasia in humans. *Human Molecular Genetics*, **29**(23): 3781–3792.
- Bagheri-Fam S, Sreenivasan R, Bernard P, Knowler KC, Sekido R, Lovell-Badge R, et al. 2012. *Sox9* gene regulation and the loss of the XY/XX sex-determining mechanism in the mole vole *Ellobius lutescens*. *Chromosome Research*, **20**(1): 191–199.
- Barske LA, Capel B. 2010. Estrogen represses SOX9 during sex determination in the red-eared slider turtle *Trachemys scripta*. *Developmental Biology*, **341**(1): 305–314.
- Chaboissier MC, Kobayashi A, Vidal VI P, Lützkendorf S, van de Kant HJG, Wegner M, et al. 2004. Functional analysis of *Sox8* and *Sox9* during sex determination in the mouse. *Development*, **131**(9): 1891–1901.
- Croft B, Ohnesorg T, Hewitt J, Bowles J, Quinn A, Tan J, et al. 2018. Human sex reversal is caused by duplication or deletion of core enhancers upstream of SOX9. *Nature Communications*, **9**(1): 5319.

- Cutting A, Chue J, Smith CA. 2013. Just how conserved is vertebrate sex determination?. *Developmental Dynamics*, **242**(4): 380–387.
- Da Silva SM, Hacker A, Harley V, Goodfellow P, Swain A, Lovell-Badge R. 1996. *Sox9* expression during gonadal development implies a conserved role for the gene in testis differentiation in mammals and birds. *Nature Genetics*, **14**(1): 62–68.
- Díaz-Hernández V, Dominguez-Mora P, Chino-Palomo L, Marmolejo-Valencia A, Harfush M, Merchant-Larios H. 2020. Spatiotemporal Expression of *Foxl2* and *Dmrt1* before, during, and after Sex Determination in the Sea Turtle *Lepidochelys olivacea*. *Sexual Development*, **13**(5–6): 286–296.
- Ge CT, Ye J, Zhang HY, Zhang Y, Sun W, Sang YP, et al. 2017. *Dmrt1* induces the male pathway in a turtle species with temperature-dependent sex determination. *Development*, **144**(12): 2222–2233.
- Gonen N, Futtner CR, Wood S, Garcia-Moreno SA, Salamone IM, Samson SC, et al. 2018. Sex reversal following deletion of a single distal enhancer of *Sox9*. *Science*, **360**(6396): 1469–1473.
- Gonen N, Quinn A, O'Neill HC, Koopman P, Lovell-Badge R. 2017. Normal levels of *Sox9* expression in the developing mouse testis depend on the TES/TESCO enhancer, but this does not act alone. *PLoS Genetics*, **13**(1): e1006520.
- Hirst CE, Major AT, Smith CA. 2018. Sex determination and gonadal sex differentiation in the chicken model. *The International Journal of Developmental Biology*, **62**(1–3): 153–166.
- Huang B, Wang SB, Ning Y, Lamb AN, Bartley J. 1999. Autosomal XX sex reversal caused by duplication of *SOX9*. *American Journal of Medical Genetics*, **87**(4): 349–353.
- Kent J, Wheatley SC, Andrews JE, Sinclair AH, Koopman P. 1996. A male-specific role for *SOX9* in vertebrate sex determination. *Development*, **122**(9): 2813–2822.
- Lavery R, Lardenois A, Ranc-Jianmotamedi F, Pauper E, Gregoire EP, Vigier C, et al. 2011. XY *Sox9* embryonic loss-of-function mouse mutants show complete sex reversal and produce partially fertile XY oocytes. *Developmental Biology*, **354**(1): 111–122.
- Matsumoto Y, Crews D. 2012. Molecular mechanisms of temperature-dependent sex determination in the context of ecological developmental biology. *Molecular and Cellular Endocrinology*, **354**(1–2): 103–110.
- Matsumoto Y, Yatsu R, Taylor C, Crews D. 2013. Changes in gonadal gene network by exogenous ligands in temperature-dependent sex determination. *Journal of Molecular Endocrinology*, **50**(3): 389–400.
- Moreno-Mendoza N, Harley VR, Merchant-Larios H. 2001. Temperature regulates *SOX9* expression in cultured gonads of *Lepidochelys olivacea*, a species with temperature sex determination. *Developmental Biology*, **229**(2): 319–326.
- Nakamura S, Watakabe I, Nishimura T, Toyoda A, Taniguchi Y, Tanaka M. 2012. Analysis of medaka *sox9* orthologue reveals a conserved role in germ cell maintenance. *PLoS One*, **7**(1): e29982.
- Pewphong R, Kitana J, Kitana N. 2021. Thermosensitive period for sex determination of the tropical freshwater turtle *Malayemys macrocephala*. *Integrative Zoology*, **16**(2): 160–169.
- Rhen T, Metzger K, Schroeder A, Woodward R. 2007. Expression of putative sex-determining genes during the thermosensitive period of gonad development in the snapping turtle, *Chelydra serpentina*. *Sexual Development*, **1**(4): 255–270.
- Sekido R, Lovell-Badge R. 2008. Sex determination involves synergistic action of *SRY* and *SF1* on a specific *Sox9* enhancer. *Nature*, **453**(7197): 930–934.
- Shoemaker C, Ramsey M, Queen J, Crews D. 2007a. Expression of *Sox9*, *Mis*, and *Dmrt1* in the gonad of a species with temperature-dependent sex determination. *Developmental Dynamics*, **236**(4): 1055–1063.
- Shoemaker CM, Queen J, Crews D. 2007b. Response of candidate sex-determining genes to changes in temperature reveals their involvement in the molecular network underlying temperature-dependent sex determination. *Molecular Endocrinology*, **21**(11): 2750–2763.
- Smith CA, Sinclair AH. 2004. Sex determination: insights from the chicken. *BioEssays*, **26**(2): 120–132.
- Spotila LD, Spotila JR, Hall SE. 1998. Sequence and expression analysis of *WT1* and *Sox9* in the red-eared slider turtle. *Trachemys scripta*. *Journal of Experimental Zoology*, **281**(5): 417–427.
- Sun D, Zhang Y, Wang C, Hua X, Zhang XA, Yan J. 2013. *Sox9*-related signaling controls zebrafish juvenile ovary–testis transformation. *Cell Death & Disease*, **4**(11): e930.
- Torres-Maldonado L, Moreno-Mendoza N, Landa A, Merchant-Larios H. 2001. Timing of *SOX9* downregulation and female sex determination in gonads of the sea turtle *Lepidochelys olivacea*. *Journal of Experimental Zoology*, **290**(5): 498–503.
- Torres Maldonado LC, Landa Piedra A, Moreno Mendoza N, Marmolejo Valencia A, Meza Martínez A, Merchant Larios H. 2002. Expression profiles of *Dax1*, *Dmrt1*, and *Sox9* during temperature sex determination in gonads of the sea turtle *Lepidochelys olivacea*. *General and Comparative Endocrinology*, **129**(1): 20–26.
- Vidal VPI, Chaboissier MC, de Rooij DG, Schedl A. 2001. *Sox9* induces testis development in XX transgenic mice. *Nature Genetics*, **28**(3): 216–217.
- Vining B, Ming ZH, Bagheri-Fam S, Harley V. 2021. Diverse regulation but conserved function: *SOX9* in vertebrate sex determination. *Genes*, **12**(4): 486.
- Wei L, Li XY, Li MH, Tang YH, Wei J, Wang DS. 2019. *Dmrt1* directly regulates the transcription of the testis-biased *Sox9b* gene in Nile tilapia (*Oreochromis niloticus*). *Gene*, **687**: 109–115.
- Western PS, Harry JL, Graves JAM, Sinclair AH. 1999. Temperature-dependent sex determination in the American alligator: *AMH* precedes *SOX9* expression. *Developmental Dynamics*, **216**(4–5): 411–419.
- Yamashita S, Kataoka K, Yamamoto H, Kato T, Hara S, Yamaguchi K, et al. 2019. Comparative analysis demonstrates cell type-specific conservation of *SOX9* targets between mouse and chicken. *Scientific Reports*, **9**(1): 12560.