

**O.M. Smorodska,
Yu.V. Moskalenko,
I.O. Vynnychenko,
O.I. Vynnychenko,
V.V. Kostuchenko**

**MODERN VECTOR IN TREATMENT
OF PATIENTS WITH LUNG CANCER:
TYROSINE KINASE INHIBITORS
IN EPIDERMAL GROWTH FACTOR
RECEPTOR MUTATIONS
(literature review)**

Sumy State University
Department of oncology and radiology
Rumskogo-Korsakova str., 2, Sumy, 40007, Ukraine
Сумський державний університет
кафедра онкології та радіології
(зав. – І.О. Винниченко)
вул. Римського-Корсакова, 2, Суми, 40007, Україна
e-mail: smorodska1991@gmail.com

Цитування: *Медичні перспективи*. 2021. Т. 26, № 2. С. 4-11
Cited: *Medicni perspektivi*. 2021;26(2):4-11

Key words: afatinib, gefitinib, EGFR, erlotinib, lung cancer, osimertinib
Ключові слова: EGFR, рак легень, ерлотиніб, гефітиніб, афатиніб, осимертиніб
Ключевые слова: EGFR, рак легких, эрлотиниб, гефитиниб, афатиниб, осимертиниб

Key words: afatinib, gefitinib, EGFR, erlotinib, lung cancer, osimertinib
Ключові слова: EGFR, рак легень, ерлотиніб, гефітиніб, афатиніб, осимертиніб
Ключевые слова: EGFR, рак легких, эрлотиниб, гефитиниб, афатиниб, осимертиниб

Abstract. Modern vector in treatment of patients with lung cancer: tyrosine kinase inhibitors in epidermal growth factor receptor mutations (literature review). Smorodska O.M., Moskalenko Yu.V., Vynnychenko I.O., Vynnychenko O.I., Kostuchenko V.V. Tumor molecular profiling in patients with non-small cell lung cancer (NSCLC) is used to identify driver mutations, which lead to premature carcinogenesis in more than 80% of adenocarcinoma cases, including epidermal growth factor receptor (EGFR) mutations. Identification of specific somatic aberrations allows to personalize treatment. Personalization of treatment resulted in improvement of NSCLC outcomes. The aim of our study was to consider scientific data on modern concepts of treatment of patients with non-small cell lung cancer with previously detected oncogenic mutations, especially EGFR mutation. In our study we analyzed scientific papers and data of international scientific literature on the problem of lung cancer treatment. Methods used: scientific research, analytical and generalizing. Different drugs are used in treatment of lung cancer. Choice of treatment scheme depends on type and presence of mutations. Patients with advanced non-small-cell lung cancer and detected mutation in the EGFR can be treated with tyrosine kinase inhibitors (TKIs). Nowadays three first generation drugs are recommended by FDA: afatinib, erlotinib, gefitinib. They showed good clinical benefit. Most patients with metastatic NSCLC typically show disease progression after approximately 9 to 13 months of erlotinib, gefitinib, or afatinib therapy. The first and only commercially available third-generation EGFR TKI is osimertinib - an oral drug, which selectively inhibits both EGFR-TKI and EGFR T790M resistance mutations. Nowadays scientists are in active investigation of mechanisms of acquired resistance to TKIs, but little is known yet. Clinical success can be observed in patients who were treated with TKIs. EGFR T790M is a mutation that leads to acquired resistance to EGFR TKI therapy. Its incidence is approximately 60% after disease progression on TKI drugs (erlotinib, gefitinib, or aphatinib). Third-generation EGFR TKIs demonstrate high efficacy, but acquired resistance development cannot be avoided. Mechanisms of acquired resistance to these agents are still investigated.

Реферат. Сучасний вектор у лікуванні пацієнтів, хворих на рак легень: інгібітори тирозин кінази при EGFR мутаціях (огляд літератури). Смородська О.М., Москаленко Ю.В., Винниченко І.О., Винниченко О.І., Костюченко В.В. Молекулярне дослідження пухлин у пацієнтів з недрібноклітинним раком легень (НДРЛ) використовується для визначення драйверних мутацій, що призводять до раннього канцерогенезу в більше ніж 80% випадках аденокарциноми, включаючи мутацію рецептора епідермального фактора росту (EGFR). Визначення специфічних соматичних мутацій дозволяє лікувати хворих персоналізовано, що приводить до значного покращення результатів лікування НДРЛ. Метою роботи було вивчення проблеми таргетної терапії НДРЛ шляхом аналізу відповідної наукової літератури. В якості досліджуваного матеріалу було використано дані закордонних джерел, які досліджують проблематику лікування раку легень. Науковий пошук, аналіз та

узагальнення були обрані в якості методів. Інгібітори тирозин кінрази (TKIs) використовують в якості стандартної терапії першої лінії для лікування пацієнтів з недрібноклітинним раком легень, у яких наявна мутація EGFR. До першого покоління інгібіторів тирозин кінрази відносять гефітиніб, ерлотиніб, афатиніб. Їх використання показало високі терапевтичні результати. У більшості пацієнтів із мутаціями EGFR та метастатичним недрібноклітинним раком легень приблизно через 9-13 місяців терапії ерлотинібом, гефітинібом або афатинібом спостерігається прогресія захворювання. Першим і єдиним комерційно доступним інгібітором рецепторів EGFR третього покоління є осимертиніб. Осимертиніб – оральний, незворотний EGFR-інгібітор третього покоління, який вибірково інгібує як EGFR, так і мутації стійкості до EGFR – T790M. Наразі вчені активно досліджують механізми набутої резистентності до інгібіторів тирозин кінрази, проте багато питань досі залишаються невирішеними. Перше покоління інгібіторів EGFR показало клінічний успіх серед пацієнтів, у яких наявна EGFR мутація. Наявність мутації EGFR T790M пов'язують з набутою резистентністю до терапії інгібіторами тирозин кінрази. Було показано, що в 60% випадків виявлення вказаних мутацій було зафіксовано після прогресії захворювання за умови початкової відповіді на терапію ерлотинібом, гефітинібом чи афатинібом. Інгібітори тирозин кіназ третього покоління демонструють високу ефективність та не сприяють розвитку набутої резистентності. Мало відомо про механізми розвитку резистентності, тому дослідження в цьому напрямку тривають.

Introduction: Non-small cell lung cancer (NSCLC) occupies the leading positions among deaths from cancer [5, 9, 11, 27, 35, 42, 48]. Molecular tumor profiling in patients with NSCLC is used to detect genetic mutations. Among adenocarcinomas, epidermal growth factor receptor (EGFR) pathology is the most common [7]. Over 10 years ago, this laid the basis for scientifically sound treatment of abandoned forms of NSCLC. During last years the clinical management of this disease has transformed because scientists found out a lot of milestones. This discovery allowed to make treatment personalized which resulted in improvement of treatment outcomes in such patients [50].

The aim of the study is to determine the depth of the problem of targeted treatment of NSCLC by analyzing the relevant scientific data in the literature on the problem of lung cancer treatment. Methods used: scientific research, analytical and generalizing.

One of the keys to the success of NSCLC treatment was in testing for EGFR mutations. It was found that EGFR mutations frequency differs in subgroups. Thus in the subgroup of Asia-Pacific region the frequency of EGFR mutation was the highest – 47%, while the lowest level of frequency of EGFR mutation was found in Oceania – 12%. Frequency of EGFR mutation is higher in women than in men and differs in regions. In Europe mutation occurs in 22% of women and 9% of men. In Africa – about 48% of women and 8% of men have this mutation. In North America – we can see 28% vs 19%. A history of smoking also affects the incidence of EGFR mutations. More often genetic mutation has been diagnosed in patients who had never smoked compared to those who have had this habit during lifetime [29].

The epidermal growth factor receptor (EGFR) is critical in proliferation and survival. Activating of these mutations are often seen in NSCLC [3, 14, 23, 27, 48]. EGFR mutation testing is recommended in patients with glandular or adenosquamous tumor variant, for squamous cell carcinomas it is uncommon [12, 22].

EGFR is one of the representatives in the tyrosine kinase receptor family. It is activated by binding dimerization ligand and receptor. This interaction leads to activation of several cellular signaling pathways: phosphoinositide 3 – kinase (PI3K) – AKT pathway, STAT pathway, and MAPK pathway. Activation of these pathways caused increased cell migration, proliferation, angiogenesis, survival and decreased apoptosis [13, 39]. In 2004 activating mutations of EGFR were identified [39]. A lot of activating mutations were described, but only two of them can be found in most cases. These main mutations are called classical activating mutations. The first one is exon 19 deletion. It was found in 85%. The second one is exon 21 L858R substitution (found in 90%). Presence of any of these two mutations is associated with a good clinical response to EGFR-targeted inhibitor therapies [11, 26, 36, 40, 49]. Clinical trials have shown that about 80% of patients had good results after treatment with tyrosine kinase inhibitors (TKI). Median of progression-free survival (PFS) was about 13 months [41]. It was found that treatment with TKIs improved overall survival compared with chemotherapy among patients with exon 19 deletion, while such improvement was not observed in patients with linked with disease L858R substitution [18, 22, 38, 45, 49].

After 9-13 months of tyrosine kinase inhibitor therapy, 60% of patients develop a specific T790M mutation, linked with acquired resistance to apha-tinib, erlotinib or gefitinib [6, 11, 24, 28, 47]. Resistance to treatment with targeted drugs may be primary. Most often, such patients have KRAS mutations or insertions of exon 20, as well as rearrangements of ALK or ROS1 genes [8, 9, 24, 47].

In 1948 Dr. Karnofsky reported that advanced lung cancer responds to cytotoxic chemotherapy. He also noticed that cytotoxic chemotherapy has lower outcomes in the treatment of advanced or metastatic lung cancer [30]. This idea leads to an absolutely

new treatment concept of patients with advanced non-small-cell lung cancer with a mutant epidermal growth factor receptor (EGFR). Thus, EGFR tyrosine kinase inhibitors (TKIs) were chosen as standard first-line therapy [44]. Afatinib, gefitinib and erlotinib have shown clinical benefit.

The third phase of clinical studies (IDEAL-1/-2) with gefitinib has shown possibility to use it as second-/ third- line treatment for pretreated patients with advanced non-small-cell lung. Once-daily 250 or 500 mg dose showed high antitumor activity with middle degree of toxicity (overall response rate – 9% to 19%; overall survival – 6 to 8 months) [37].

INTACT-1 and -2 study analyzes benefit of gefatinib vs platinum-doublet chemotherapy for treatment of patients with unresectable stage III/IV NSCLC. In this clinical trial chemotherapy-naïve patients were randomly divided into two groups: first group was treated with gefitinib, while second received placebo in combination with platinum-doublet chemotherapy. There were no additional benefits observed in the main indicators of survival and prognosis compared with standard chemotherapy. In 2003 based on the results of INTACT gefitinib was approved by the US FDA as monotherapy for patient with locally advanced or metastatic NSCLC after completion of the classic scheme of cytotoxic chemotherapy [35, 44].

Since 2004 erlotinib has been used to treat patients with locally advanced or/and metastatic NSCLC. It was prescribed after at least one previous chemotherapy regimen [19, 33]. For patients with sensitizing EGFR mutations, the FDA has permitted the use of this drug as first-line therapy in the case of metastatic, recurrent, or progressive tumors [17, 19]. Treatment of patients with the most common activating EGFR mutations with the first generation EGFR TKIs lead to good clinical success [4, 33]. Reliable data were obtained in a randomized phase III study (IPASS). Patients with EGFR sensitizing mutations receiving gefitinib had significantly higher survival and better quality of life compared with the group of patients treated with paclitaxel and carboplatin (standard chemotherapy) [16]. A significant improvement in disease-free survival up to 11.8 months when using erlotinib as a first-line therapy is reported in Chinese studies on phase III OPTIMAL and phase II JO22903 [15].

The aim of the updated study (CALGB 30406) was to compare treatment with erlotinib and its combination with chemotherapy in patients (predominantly white) with longstanding lung cancer. Patients receiving erlotinib as monotherapy had fewer side effects compared with chemotherapy and erlotinib group. If EGFR sensitizing mutations are detected during chemotherapy, it was recommended

to discontinue or finish planned chemotherapy and switch to target therapy. According to the NCCN guidelines based on the CALGB study, it is not recommended to combine erlotinib, gefitinib or aphantinib with chemotherapy. If the patient has a progression of the disease, but has no clinical symptoms, it is possible to continue target therapy [34].

A similar drug was afatinib. It has slightly more side effects than erlotinib or gefitinib, but has been approved by the FDA as a first-line therapy drug [2, 17, 25, 40].

It is a proven fact that the progression of the disease begins about a year after taking targeted drugs. EGFR T790M mutation occurs, which indicates the development of resistance to therapy with TKI [31, 35]. Occurrence of T790M mutation in exon 20 of the EGFR gene leads to loosing of efficacy of quinazoline-based first-generation EGFR TKI. Loosing of efficacy is observed due to absence of opportunity to bind to the ATP binding pocket at a receptor, which decreases efficacy of inhibition of signaling in future. Biology and epidemiology of the T790M resistance mutation has been studied well. For targeting at T790M mutation, third-generation EGFR TKIs were designed [1, 20]. Pyrimidines named WZ3146, WZ4002 and WZ8040 were investigated and WZ4002 was found to be the most powerful against EGFR T790M. However, the drug has not been approved by the FDA [20].

The only commercially available third-generation TKI is osimertinib. Osimertinib is a third-generation EGFR-TKI. One of its benefits is that the drug is used orally. It can selectively inhibit both sensitizing to EGFR-TKI mutations and EGFR T790M resistance mutations. But it has lower activity against wild-type. Positive results of the AURA clinical trial let osimertinib be approved worldwide for the treatment of patients with the T790M mutation, which arose during treatment with TKIs of previous generation [8, 38].

A randomized study of phase III EGFR (among patients with EGFR T790M-positive metastatic NSCLC), evaluating platinum and pemetrexed chemotherapy against osimertinib was performed. The study showed that treatment with osimertinib increased PFS compared with standard chemotherapy (10.1 vs 4.4 months), especially among patients with CNS metastases. The disease controlled rate was higher in patients treated with osimertinib compared with standart chemotherapy (93% vs 74%) [10].

It was found, that osimertinib showed good clinical benefits for treatment of patients with metastatic EGFR T790M-positive NSCLC with disease progression after or during TKI EGFR therapy. Due to this the FDA approved it. Later osimertinib was also recommended by the NCCN [34, 38].

Even after disease progression, it is recommended to continue treatment with erlotinib, gefitinib or aphantinib as this is beneficial for the patient. It is known that discontinuation of therapy leads to faster progression of the disease (appearance of symptoms, increase in tumor size and deterioration of the response during PET scan) [32].

In 2017, the NCCN group made adjustments to the treatment protocols of patients with EGFR sensitizing mutations which progressed on erlotinib, gefitinib, or aphantinib. After review, osimertinib was recommended for the treatment of patients with symptomatic brain metastases. As an alternative treatment regimen, continuation of afatinib, erlotinib or gefitinib therapy was suggested for this group of patients. When continuing therapy, it was recommended to add or modify additional therapy (e.g. local or systemic therapy). First-line systemic therapy combinations were recommended for patients who have multiple symptomatic lesions and were not sensitive to T790M. If the patient was positive for T790M, it was recommended to include osimertinib in his therapy [34].

Other mechanisms of resistance are multivarious. They include HER2 and / or MET amplification, PIK3CA and / or BRAF mutation, and small cell

lung cancer transformation. Their incidence is much lower than the T790M mutation. Extremely rare these mutations or amplifications occurred together with T790M mutation [43, 46].

CONCLUSIONS

1. Clinical success was observed in treatment with the first generation EGFR TKIs. Similar results were observed in patients with the most common EGFR activating mutations.

2. Presence of 790M mutation in EGFR leads to resistance to TKI therapy. Its incidence is approximately in 60% among patients with disease progression on TKI drugs (erlotinib, gefitinib, or aphantinib).

3. Third-generation TKIs of EGFR demonstrate high efficacy, but acquired resistance development cannot be avoided.

4. Reasons and ways of development of acquired resistance to these agents are still investigated.

Conflict of interests. The authors declare no conflict of interest.

This research was performed as part of the research topics of Department of Oncology and Radiology of Sumy State University № 0118U003570 "Efficiency of liquid biopsy and tissue biopsy in the diagnosis and treatment of malignant tumors".

REFERENCES

- Zhang Y, Chen Z, Zhang X, Xu C, Yan H, Xie Z, et al. Analysis of resistance mechanisms to abivertinib, a third-generation EGFR tyrosine kinase inhibitor, in patients with EGFR T790M-positive non-small cell lung cancer from a phase I trial. *EBioMedicine*. 2019;43:180-7. doi: <https://doi.org/10.1016/j.ebiom.2019.04.030>
- Goto K, Nishio M, Yamamoto N, Chikamori K, Hida T, Maemondo M, et al. A prospective, phase II, open-label study (JO22903) of first-line erlotinib in Japanese patients with epidermal growth factor receptor (EGFR) mutation-positive advanced non-small-cell lung cancer (NSCLC). *Lung Cancer*. 2013;82:109-14. doi: <https://doi.org/10.1016/j.lungcan.2013.07.003>
- Chen R, Chen H, Jiang B, Zhang X, Zhou Q, Tu H, et al. Bevacizumab plus chemotherapy for patients with advanced pulmonary adenocarcinoma harboring EGFR mutations. *Clinical and Translational Oncology*. 2017;20(2):243-52. doi: <https://doi.org/10.1007/s12094-017-1714-2>
- Yang JC, Sequist LV, Geater SL, Tsai CM, Mok TS, Schuler M, et al. Clinical activity of afatinib in patients with advanced non-small-cell lung cancer harbouring uncommon EGFR mutations: a combined post-hoc analysis of LUX-Lung 2, LUX-Lung 3, and LUX-Lung 6. *Lancet Oncol*. 2015;16(7):830-8. doi: [https://doi.org/10.1016/S1470-2045\(15\)00026-1](https://doi.org/10.1016/S1470-2045(15)00026-1)
- Lou N, Zhang X, Chen H, Zhou Q, Yan L, Xie Z, et al. Clinical outcomes of advanced non-small-cell lung cancer patients with EGFR mutatio, ALK rearrangement and EGFR/ALK co-alterations. *Oncotarget*. 2016;7(40):65185-95. doi: <https://doi.org/10.18632/oncotarget.11218>
- Da Cunha Santos G, Shepherd FA, Tsao MS. EGFR mutations and lung cancer. *Annu Rev Pathol*. 2011;6:49-69. doi: <https://doi.org/10.1146/annurev-pathol-011110-130206>
- Dearden S, Stevens J, Wu YL and Blowers D. Mutation incidence and coincidence in non small-cell lung cancer: meta-analyses by ethnicity and histology (mutMap). *Ann Oncol*. 2013;24:2371-6. doi: <https://doi.org/10.1093/annonc/mdt205>
- Finlay MR, Anderton M, Ashton S, Ballard P, Bethel PA, Box MR, et al. Discovery of a potent and selective EGFR inhibitor (AZD9291) of both sensitizing and T790M resistance mutations that spares the wild type form of the receptor. *J Med Chem*. 2014;57:8249-67. doi: <https://doi.org/10.1021/jm500973a>
- Tetsu O, Hangauer M, Phuchareon J, Eisele D, McCormick F. Drug Resistance to EGFR Inhibitors in Lung Cancer. *Chemotherapy*. 2016;61(5):223-35. doi: <https://doi.org/10.1159/000443368>
- Dungo RT, Keating GM. Afatinib: first global approval. *Drugs*. 2013;73:1503-15. doi: <https://doi.org/10.1007/s40265-013-0111-6>
- Yi L, Fan J, Qian R, Luo P, Zhang J. Efficacy and safety of osimertinib in treating EGFR-mutated advanced NSCLC: A meta-analysis. *International Journal of Cancer*. 2019;145(1):284-94. doi: <https://doi.org/10.1002/ijc.32097>

12. Arcila ME, Nafa K, Chaft JE, Rekhman N, Lau C, Reva BA, et al. EGFR exon 20 insertion mutations in lung adenocarcinomas: prevalence, molecular heterogeneity, and clinicopathologic characteristics. *Mol Cancer Ther*. 2013;12:220-9. doi: <https://doi.org/10.1158/1535-7163.MCT-12-0620>
13. Konduri K, Gallant J, Chae Y, Giles F, Gitlitz B, Gowen K, et al. EGFR Fusions as Novel Therapeutic Targets in Lung Cancer. *Cancer Discovery*. 2016;6(6):601-11. doi: <https://doi.org/10.1158/2159-8290.CD-16-0075>
14. Ripamonti F, Albano L, Rossini A, et al. EGFR through STAT3 modulates $\Delta N63\alpha$ expression to sustain tumor-initiating cell proliferation in squamous cell carcinomas. *J Cell Physiol*. 2013;228:871-8. doi: <https://doi.org/10.1002/jcp.24238>
15. Antonicelli A, Cafarotti S, Indini A, Galli A, Russo A, Cesario A et al. EGFR-targeted therapy for non-small cell lung cancer: focus on EGFR oncogenic mutation. *Int J Med Sci*. 2013;10:320-30. doi: <https://doi.org/10.7150/ijms.4609>
16. Shepherd FA, Pereira JR, Ciuleanu T, Tan EH, Hirsh V, Thongprasert S, et al. Erlotinib in previously treated non-small-cell lung cancer. *N Engl J Med*. 2005;353(2):123-32. doi: <https://doi.org/10.1056/NEJMoa050753>
17. Jiang Y, Zhang J, Huang J, Xu B, Li N, Cao L, et al. Erlotinib versus gefitinib for brain metastases in Asian patients with exon 19 EGFR-mutant lung adenocarcinoma: a retrospective, multicenter study. *BMC Pulmonary Medicine*. 2018;18(1). doi: <https://doi.org/10.1186/s12890-018-0734-1>
18. Yu X, Zhang X, Zhang Z, Lin Y, Wen Y, Chen Y, et al. First-generation EGFR tyrosine kinase inhibitor therapy in 106 patients with compound EGFR-mutated lung cancer: a single institution's clinical practice experience. *Cancer Communications*. 2018;38(1):51. doi: <https://doi.org/10.1186/s40880-018-0321-0>
19. Giaccone G, Herbst RS, Manegold C, Scagliotti G, Rosell R, Natale V, et al. Gefitinib in combination with gemcitabine and cisplatin in advanced non-small-cell lung cancer: a phase III trial – INTACT 1. *J Clin Oncol*. 2004;22(5):777-84. doi: <https://doi.org/10.1200/JCO.2004.08.001>
20. Mok TS, Wu YL, Thongprasert S, Yang CH, Chu DT, Saijo N, et al. Gefitinib or carboplatin-paclitaxel in pulmonary adenocarcinoma. *N Engl J Med*. 2009;361:947-57. doi: <https://doi.org/10.1056/NEJMoa0810699>
21. Chang C, Lee C, Ko J, Chang L, Lee M, Wang J, et al. Gefitinib or erlotinib in previously treated non-small-cell lung cancer patients: a cohort study in Taiwan. *Cancer Medicine*. 2017;6(7):1563-72. doi: <https://doi.org/10.1002/cam4.1121>
22. Chang I, Jiang S, Yang J, Su W, Chien L, Hsiao C, et al. Genetic Modifiers of Progression-Free Survival in Never-Smoking Lung Adenocarcinoma Patients Treated with First-Line Tyrosine Kinase Inhibitors. *American Journal of Respiratory and Critical Care Medicine*. 2017;195(5):663-73. doi: <https://doi.org/10.1164/rccm.201602-0300OC>
23. Im J, Herrmann A, Bernatchez C, Haymaker C, Molldrem J, Hong W, et al. Immune-Modulation by Epidermal Growth Factor Receptor Inhibitors: Implication on Anti-Tumor Immunity in Lung Cancer. *PLOS ONE*. 2016;11(7):e0160004. doi: <https://doi.org/10.1371/journal.pone.0160004>
24. Santoni-Rugiu E, Melchior L, Urbanska E, Jakobsen J, Stricker K, Grauslund M, et al. Intrinsic resistance to EGFR-Tyrosine Kinase Inhibitors in EGFR-Mutant Non-Small Cell Lung Cancer: Differences and Similarities with Acquired Resistance. *Cancers*. 2019;11(7):923. doi: <https://doi.org/10.3390/cancers11070923>
25. Ke E, Wu Y. Afatinib in the first-line treatment of epidermal-growth-factor-receptor mutation-positive non-small cell lung cancer: a review of the clinical evidence. *Therapeutic Advances in Respiratory Disease*. 2016;10(3):256-64. doi: <https://doi.org/10.1177/1753465816634545>
26. Kennedy BJ. The snail's pace of lung carcinoma chemotherapy cancer. *Cancer*. 1998;82(5):801-3. doi: [https://doi.org/10.1002/\(SICI\)1097-0142\(19980301\)82:5<801::AID-CNCR1>3.0.CO;2-M](https://doi.org/10.1002/(SICI)1097-0142(19980301)82:5<801::AID-CNCR1>3.0.CO;2-M)
27. Liu X, Wang P, Zhang C, Ma Z. Epidermal growth factor receptor (EGFR): A rising star in the era of precision medicine of lung cancer. *Oncotarget*. 2017;8(30):50209-20. doi: <https://doi.org/10.18632/oncotarget.16854>
28. Westover D, Zugazagoitia J, Cho B, Lovly C, Paz-Ares L. Mechanisms of acquired resistance to first- and second-generation EGFR tyrosine kinase inhibitors. *Annals of Oncology*. 2018;29:10-19. doi: <https://doi.org/10.1093/annonc/mdx703>
29. Midha A, Dearden S, McCormack R. EGFR mutation incidence in non-small-cell lung cancer of adenocarcinoma histology: a systematic review and global map by ethnicity (mutMapII). *Am J Cancer Res*. 2015;5(9):2892-911. PMID: 26609494; PMCID: PMC4633915.
30. Fukuoka M, Yano S, Giaccone G, Tamura T, Nakagawa K, Nishiwaki JYDY, et al. Multi-institutional randomized phase II trial of gefitinib for previously treated patients with advanced non-small-cell lung cancer. *J Clin Oncol*. 2003;21(12):2237-46. doi: <https://doi.org/10.1200/JCO.2003.10.038>
31. Ettinger DS, Wood D E, Aisner D L, Akerley W, Bauman J, Chirieac LR, et al. Non-Small Cell Lung Cancer, Version 5.2017. *Clinical Practice Guidelines in Oncology*. JNCCN.2017;15(4):504-534. doi: <https://doi.org/10.6004/jnccn.2017.0050>
32. Zhou W, Ercan D, Chen I, Yun CH, Li D, Chappelletti M, et al. Novel mutant-selective EGFR kinase inhibitors against EGFR T790M. *Nature*. 2000;462(7276):1070-4. doi: <https://doi.org/10.1038/nature08622>
33. Wen F, Zheng H, Zhang P, Hutton D, Li Q. OPTIMAL and ENSURE trials-based combined cost-effectiveness analysis of erlotinib versus chemotherapy for the first-line treatment of Asian patients with non-squamous non-small-cell lung cancer. *BMJ Open*. 2018;8(4):e020128. doi: <https://doi.org/10.1136/bmjopen-2017-020128>
34. Mok TS, Wu YL, Ahn MJ, Garassino MC, Kim HR, Ramalingam SS, et al. Osimertinib or platinum-



- pemetrexed in EGFR T790M-positive lung cancer. *N Engl J Med.* 2017;376:629-40.
doi: <https://doi.org/10.1056/NEJMoa1612674>
35. Pakkala S, Ramalingam S. Personalized therapy for lung cancer: Striking a moving target. *JCI Insight.* 2018;3(15):120858.
doi: <https://doi.org/10.1172/jci.insight.120858>
36. Bonomi P, Gandara D, Hirsch F, Kerr K, Obasaju C, Paz-Ares L, et al. Predictive biomarkers for response to EGFR-directed monoclonal antibodies for advanced squamous cell lung cancer. *Annals of Oncology.* 2018;29(8):1701-9.
doi: <https://doi.org/10.1093/annonc/mdy196>
37. Riely GJ, Kris MG, Zhao B, Tim A, Milton DT, Moore E, et al. Prospective assessment of discontinuation and reinitiation of erlotinib or gefitinib in patients with acquired resistance to erlotinib or gefitinib followed by the addition of everolimus. *Clin Cancer Res.* 2007;13:5150-5.
doi: <https://doi.org/10.1158/1078-0432.CCR-07-0560>
38. Leonetti A, Sharma S, Minari R, Perego P, Giovannetti E, Tiseo M. Resistance mechanisms to osimertinib in EGFR-mutated non-small cell lung cancer. *British Journal of Cancer.* 2019;121(9):725-37.
doi: <https://doi.org/10.1038/s41416-019-0573-8>
39. Paik PK, Varghese AM, Sima CS, Moreira AL, Ladanyi M, Kris MG, et al. Response to erlotinib in patients with EGFR mutant advanced non-small cell lung cancers with a squamous or squamous-like component. *Mol Cancer Ther.* 2012;11:2535-40.
doi: <https://doi.org/10.1158/1535-7163.MCT-12-0163>
40. Ricciuti B, Baglivo S, De Giglio A, Chiari R. Afatinib in the first-line treatment of patients with non-small cell lung cancer: clinical evidence and experience. *Therapeutic Advances in Respiratory Disease.* 2018;12:175346661880865.
doi: <https://doi.org/10.1177/1753466618808659>
41. Sharma SV, Bell DW, Settleman J, Haber DA. Epidermal growth factor receptor mutations in lung cancer. *Nat Rev Cancer* 2007;7:169-81.
doi: <https://doi.org/10.1038/nrc2088>
42. Siegel R, Naishadham D, Jemal A. Cancer statistics 2013. *CA Cancer J Clin* 2013;63:11-30.
doi: <https://doi.org/10.3322/caac.21166>
43. Soria J.-C, Ohe Y, Vansteenkiste J, Osimertinib in Untreated EGFR-Mutated Advanced Non-Small-Cell Lung Cancer *N Engl J Med.* 2018;378(2):113-25.
doi: <https://doi.org/10.1056/NEJMoa1713137>
44. Stewart EL, Tan SZ, Liu G. Known and putative mechanism of resistance to EGFR targeted therapies in NSCLC patients with EGFR mutations-a review. *Transl Lung Cancer Res.* 2015;4(1):67-81.
doi: <https://doi.org/10.3978/j.issn.2218-6751.2014.11.06>
45. Masters GA, Temin S, Azzoli CG, Giaccone G, Baker S, Brahmer JR, et al. Systemic therapy for Stage IV non small cell lung cancer: American Society of Clinical Oncology Clinical Practice Guideline update. *J Clin Oncol.* 2015;33:3488-515.
doi: <https://doi.org/10.1200/JCO.2015.62.1342>
46. Gou L, Li A, Yang J, Zhang X, Su J, Yan H, et al. The coexistence of MET over-expression and an EGFR T790M mutation is related to acquired resistance to EGFR tyrosine kinase inhibitors in advanced non-small cell lung cancer. *Oncotarget.* 2016;7(32):51311-9.
doi: <https://doi.org/10.18632/oncotarget.9697>
47. Tseng J, Su K, Yang T, Chen K, Hsu K, Chen H, et al. The emergence of T790M mutation in EGFR-mutant lung adenocarcinoma patients having a history of acquired resistance to EGFR-TKI: focus on rebiopsy timing and long-term existence of T790M. *Oncotarget.* 2016;7(30):48059-69.
doi: <https://doi.org/10.18632/oncotarget.10351>
48. Ochiai S, Nomoto Y, Watanabe Y, Yamashita Y, Toyomasu Y, Kawamura T, et al. The impact of epidermal growth factor receptor mutations on patterns of disease recurrence after chemoradiotherapy for locally advanced non-small cell lung cancer: a literature review and pooled analysis. *Journal of Radiation Research.* 2016;57(5):449-59.
doi: <https://doi.org/10.1093/jrr/rrw075>
49. Wu S, Shih J. Management of acquired resistance to EGFR TKI-targeted therapy in advanced non-small cell lung cancer. *Molecular Cancer.* 2018;17(1).
doi: <https://doi.org/10.1186/s12943-018-0777-1>
50. Zhang C, Leighl N, Wu Y, Zhong W. Emerging therapies for non-small cell lung cancer. *Journal of Hematology & Oncology.* 2019;12(1).
doi: <https://doi.org/10.1186/s13045-019-0731-8>

СПИСОК ЛІТЕРАТУРИ

1. Analysis of resistance mechanisms to abivertinib, a third-generation EGFR tyrosine kinase inhibitor, in patients with EGFR T790M-positive non-small cell lung cancer from a phase I trial / Y. Zhang et al. *EBioMedicine.* 2019. Vol. 43. P. 180-187.
DOI: <https://doi.org/10.1016/j.ebiom.2019.04.030>
2. A prospective, phase II, open-label study (JO22903) of first-line erlotinib in Japanese patients with epidermal growth factor receptor (EGFR) mutation-positive advanced non-small-cell lung cancer (NSCLC) / K. Goto et al. *Lung Cancer.* 2013. Vol. 82. P. 109-114.
DOI: <https://doi.org/10.1016/j.lungcan.2013.07.003>
3. Bevacizumab plus chemotherapy for patients with advanced pulmonary adenocarcinoma harboring EGFR mutations / R. Chen et al. *Clinical and Translational Oncology.* 2017. Vol. 20, No. 2. P. 243-252.
DOI: <https://doi.org/10.1007/s12094-017-1714-2>
4. Clinical activity of afatinib in patients with advanced non-small-cell lung cancer harbouring uncommon EGFR mutations: a combined post-hoc analysis of LUX-Lung 2, LUX-Lung 3, and LUX-Lung 6 / J. C. Yang et al. *Lancet Oncol.* 2015. Vol. 16, No. 7. P. 830-838.
DOI: [https://doi.org/10.1016/S1470-2045\(15\)00026-1](https://doi.org/10.1016/S1470-2045(15)00026-1)

5. Clinical outcomes of advanced non-small-cell lung cancer patients with EGFR mutation, ALK rearrangement and EGFR/ALK co-alterations / N. Lou, et al. *Oncotarget*. 2016. Vol. 7, No. 40. P. 65185-65195. DOI: <https://doi.org/10.18632/oncotarget.11218>
6. Da Cunha Santos G., Shepherd F. A., Tsao M. S. EGFR mutations and lung cancer. *Annu Rev Pathol*. 2011. Vol. 6. P. 49-69. DOI: <https://doi.org/10.1146/annurev-pathol-011110-130206>
7. Dearden S., Stevens J., Wu Y. L., Blowers D. Mutation incidence and coincidence in non small-cell lung cancer: meta-analyses by ethnicity and histology (mutMap). *Ann Oncol*. 2013. Vol. 24. P. 2371-2376. DOI: <https://doi.org/10.1093/annonc/mdt205>
8. Discovery of a potent and selective EGFR inhibitor (AZD9291) of both sensitizing and T790M resistance mutations that spares the wild type form of the receptor / M. R. Finlay et al. *J Med Chem*. 2014. Vol. 57. P. 8249-8267. DOI: <https://doi.org/10.1021/jm500973a>
9. Drug Resistance to EGFR Inhibitors in Lung Cancer / O. Tetsu et al. *Chemotherapy*. 2016. Vol. 61, No. 5. P. 223-235. DOI: <https://doi.org/10.1159/000443368>
10. Dungo RT. Keating G. M. Afatinib: first global approval. *Drugs*. 2013. Vol. 73. P. 1503-1515. DOI: <https://doi.org/10.1007/s40265-013-0111-6>
11. Efficacy and safety of osimertinib in treating EGFR-mutated advanced NSCLC: A meta-analysis / L. Yi et al. *International Journal of Cancer*. 2019. Vol. 145, No. 1. P. 284-294. DOI: <https://doi.org/10.1002/ijc.32097>
12. EGFR exon 20 insertion mutations in lung adenocarcinomas: prevalence, molecular heterogeneity, and clinicopathologic characteristics / M. E. Arcila et al. *Mol Cancer Ther*. 2013. Vol. 12. P. 220-229. DOI: <https://doi.org/10.1158/1535-7163.MCT-12-0620>
13. EGFR Fusions as Novel Therapeutic Targets in Lung Cancer / K. Konduri et al. *Cancer Discovery*. 2016. Vol. 6, No. 6. P. 601-611. DOI: <https://doi.org/10.1158/2159-8290.CD-16-0075>
14. EGFR through STAT3 modulates $\Delta N63\alpha$ expression to sustain tumor-initiating cell proliferation in squamous cell carcinomas / F. Ripamonti et al. *J Cell Physiol*. 2013. Vol. 228. P. 871-878. DOI: <https://doi.org/10.1002/jcp.24238>
15. EGFR-targeted therapy for non-small cell lung cancer: focus on EGFR oncogenic mutation / A. Antonicelli et al. *Int J Med Sci*. 2013. Vol. 10. P. 320-330. DOI: <https://doi.org/10.7150/ijms.4609>
16. Erlotinib in previously treated non-small-cell lung cancer / F. A. Shepherd et al. *N Engl J Med*. 2005. Vol. 353, No. 2. P. 123-132. DOI: <https://doi.org/10.1056/NEJMoa050753>
17. Erlotinib versus gefitinib for brain metastases in Asian patients with exon 19 EGFR-mutant lung adenocarcinoma: a retrospective, multicenter study / Y. Jiang J et al. *BMC Pulmonary Medicine*. 2018. Vol. 186 No. 1. DOI: <https://doi.org/10.1186/s12890-018-0734-1>
18. First-generation EGFR tyrosine kinase inhibitor therapy in 106 patients with compound EGFR-mutated lung cancer: a single institution's clinical practice experience / X. Yu et al. *Cancer Communications*. 2018. Vol. 38, No. 1. P. 51. DOI: <https://doi.org/10.1186/s40880-018-0321-0>
19. Gefitinib in combination with gemcitabine and cisplatin in advanced non-small-cell lung cancer: a phase III trial – INTACT 1 / G. Giaccone et al. *J Clin Oncol*. 2004. Vol. 22, No. 5. P. 777-784. DOI: <https://doi.org/10.1200/JCO.2004.08.001>
20. Gefitinib or carboplatin-paclitaxel in pulmonary adenocarcinoma / T. S. Mok et al. *N Engl J Med*. 2009. Vol. 361. P. 947-957. DOI: <https://doi.org/10.1056/NEJMoa0810699>
21. Gefitinib or erlotinib in previously treated non-small-cell lung cancer patients: a cohort study in Taiwan / C. Chang et al. *Cancer Medicine*. 2017. Vol. 6, No. 7. P. 1563-1572. DOI: <https://doi.org/10.1002/cam4.1121>
22. Genetic Modifiers of Progression-Free Survival in Never-Smoking Lung Adenocarcinoma Patients Treated with First-Line Tyrosine Kinase Inhibitors / I. Chang et al. *American Journal of Respiratory and Critical Care Medicine*. 2017. Vol. 195, No. 5. P. 663-673. DOI: <https://doi.org/10.1164/rccm.201602-0300OC>
23. Immune-Modulation by Epidermal Growth Factor Receptor Inhibitors: Implication on Anti-Tumor Immunity in Lung Cancer / J. Im et al. *PLOS ONE*. 2016. Vol. 11, No. 7. P. e0160004. DOI: <https://doi.org/10.1371/journal.pone.0160004>
24. Intrinsic resistance to EGFR-Tyrosine Kinase Inhibitors in EGFR-Mutant Non-Small Cell Lung Cancer: Differences and Similarities with Acquired Resistance / E. Santoni-Rugiu et al. *Cancers*. 2019. Vol. 11, No. 7. P. 923. DOI: <https://doi.org/10.3390/cancers11070923>
25. Ke E., Wu Y. Afatinib in the first-line treatment of epidermal-growth-factor-receptor mutation-positive non-small cell lung cancer: a review of the clinical evidence. *Therapeutic Advances in Respiratory Disease*. 2016. Vol. 10, No. 3. P. 256-264. DOI: <https://doi.org/10.1177/1753465816634545>
26. Kennedy B. J. The snail's pace of lung carcinoma chemotherapy cancer. *Cancer*. 1998. Vol. 82, No. 5. P. 801-803. DOI: [https://doi.org/10.1002/\(SICI\)1097-0142\(19980301\)82:5<801::AID-CNCR1>3.0.CO;2-M](https://doi.org/10.1002/(SICI)1097-0142(19980301)82:5<801::AID-CNCR1>3.0.CO;2-M)
27. Liu X., Wang P., Zhang C., Ma Z. Epidermal growth factor receptor (EGFR): A rising star in the era of precision medicine of lung cancer. *Oncotarget*. 2017. Vol. 8, No. 30. P. 50209-50220. DOI: <https://doi.org/10.18632/oncotarget.16854>
28. Mechanisms of acquired resistance to first- and second-generation EGFR tyrosine kinase inhibitors / D. Westover et al. *Annals of Oncology*. 2018. Vol. 29. P. 10-19. DOI: <https://doi.org/10.1093/annonc/mdx703>
29. Midha A., Dearden S., McCormack R. EGFR mutation incidence in non-small-cell lung cancer of adenocarcinoma histology: a systematic review and global map by ethnicity (mutMapII). *Am J Cancer Res*. 2015. Vol. 5, No. 9. P. 2892-2911. PMID: 26609494; PMCID: PMC4633915.
30. Multi-institutional randomized phase II trial of gefitinib for previously treated patients with advanced non-small-cell lung cancer / M. Fukuoka et al. *J Clin*

- Oncol.* 2003. Vol. 21, No. 12. P. 2237-2246. DOI: <https://doi.org/10.1200/JCO.2003.10.038>
31. Non-Small Cell Lung Cancer, Version 5.2017. Clinical Practice Guidelines in Oncology / D. S. Ettinger et al. *JNCCN*. 2017. Vol. 15, No. 4. P. 504-534. DOI: <https://doi.org/10.6004/jnccn.2017.0050>
32. Novel mutant-selective EGFR kinase inhibitors against EGFR T790M / W. Zhou et al. *Nature*. 2000. Vol. 462, No. 7276. P. 1070-1074. DOI: <https://doi.org/10.1038/nature08622>
33. OPTIMAL and ENSURE trials-based combined cost-effectiveness analysis of erlotinib versus chemotherapy for the first-line treatment of Asian patients with non-squamous non-small-cell lung cancer / F. Wen et al. *BMJ Open*. 2018. Vol. 8, No. 4. P. e020128. DOI: <https://doi.org/10.1136/bmjopen-2017-020128>
34. Osimertinib or platinum-pemetrexed in EGFR T790M-positive lung cancer / T. S. Mok et al. *N Engl J Med*. 2017. Vol. 376. P. 629-640. DOI: <https://doi.org/10.1056/NEJMoa1612674>
35. Pakkala S., Ramalingam S. Personalized therapy for lung cancer: Striking a moving target. *JCI Insight*. 2018. Vol. 3, No. 15. P. 120858. DOI: <https://doi.org/10.1172/jci.insight.120858>
36. Predictive biomarkers for response to EGFR-directed monoclonal antibodies for advanced squamous cell lung cancer / P. Bonomi et al. *Annals of Oncology*. 2018. Vol. 29, No. 8. P. 1701-1709. DOI: <https://doi.org/10.1093/annonc/mdy196>
37. Prospective assessment of discontinuation and reinitiation of erlotinib or gefitinib in patients with acquired resistance to erlotinib or gefitinib followed by the addition of everolimus / G. J. Riely et al. *Clin Cancer Res*. 2007. Vol. 13. P. 5150-5155. DOI: <https://doi.org/10.1158/1078-0432.CCR-07-0560>
38. Resistance mechanisms to osimertinib in EGFR-mutated non-small cell lung cancer / A. Leonetti et al. *British Journal of Cancer*. 2019. Vol. 121, No. 9. P. 725-737. DOI: <https://doi.org/10.1038/s41416-019-0573-8>
39. Response to erlotinib in patients with EGFR mutant advanced non-small cell lung cancers with a squamous or squamous-like component / P. K. Paik et al. *Mol Cancer Ther*. 2012. Vol. 11. P. 2535-2540. DOI: <https://doi.org/10.1158/1535-7163.MCT-12-0163>
40. Ricciuti B., Baglivo S., De Giglio A., Chiari R. Afatinib in the first-line treatment of patients with non-small cell lung cancer: clinical evidence and experience. *Therapeutic Advances in Respiratory Disease*. 2018. Vol. 12. P. 175346661880865. DOI: <https://doi.org/10.1177/1753466618808659>
41. Sharma S. V., Bell D. W., Settleman J., Haber D. A. Epidermal growth factor receptor mutations in lung cancer. *Nat Rev Cancer*. 2007. Vol. 7. P. 169-181. DOI: <https://doi.org/10.1038/nrc2088>
42. Siegel R., Naishadham D., Jemal A. Cancer statistics 2013. *CA Cancer J Clin*. 2013. Vol. 63. P. 11-30. DOI: <https://doi.org/10.3322/caac.21166>
43. Soria J.-C., Ohe Y., Vansteenkiste J. Osimertinib in Untreated EGFR-Mutated Advanced Non-Small-Cell Lung Cancer. *N Engl J Med*. 2018. Vol. 378, No. 2. P. 113-125. DOI: <https://doi.org/10.1056/NEJMoa1713137>
44. Stewart E. L., Tan S. Z., Liu G. Known and putative mechanism of resistance to EGFR targeted therapies in NSCLC patients with EGFR mutations-a review. *Transl Lung Cancer Res*. 2015. Vol. 4, No. 1. P. 67-81. DOI: <https://doi.org/10.3978/j.issn.2218-6751.2014.11.06>
45. Systemic therapy for Stage IV non small cell lung cancer: American Society of Clinical Oncology Clinical Practice Guideline update / G. A. Masters et al. *J Clin Oncol*. 2015. Vol. 33. P. 3488-3515. DOI: <https://doi.org/10.1200/JCO.2015.62.1342>
46. The coexistence of MET over-expression and an EGFR T790M mutation is related to acquired resistance to EGFR tyrosine kinase inhibitors in advanced non-small cell lung cancer / L. Gou et al. *Oncotarget*. 2016. Vol. 7, No. 32. P. 51311-51319. DOI: <https://doi.org/10.18632/oncotarget.9697>
47. The emergence of T790M mutation in EGFR-mutant lung adenocarcinoma patients having a history of acquired resistance to EGFR-TKI: focus on rebiopsy timing and long-term existence of T790M / J. Tseng et al. *Oncotarget*. 2016. Vol. 7, No. 30. P. 48059-48069. DOI: <https://doi.org/10.18632/oncotarget.10351>
48. The impact of epidermal growth factor receptor mutations on patterns of disease recurrence after chemoradiotherapy for locally advanced non-small cell lung cancer: a literature review and pooled analysis / S. Ochiai et al. *Journal of Radiation Research*. 2016. Vol. 57, No. 5. P. 449-459. DOI: <https://doi.org/10.1093/jrr/rrw075>
49. Wu S., Shih J. Management of acquired resistance to EGFR TKI-targeted therapy in advanced non-small cell lung cancer. *Molecular Cancer*. 2018. Vol. 17, No. 1. DOI: <https://doi.org/10.1186/s12943-018-0777-1>
50. Zhang C., Leigh N., Wu Y., Zhong W. Emerging therapies for non-small cell lung cancer. *Journal of Hematology & Oncology*. 2019. Vol. 12, No. 1. DOI: <https://doi.org/10.1186/s13045-019-0731-8>

The article was received
2020.02.05

