



COMPLEMENTARY OF CLASSICAL MEANS WITH RESPECT TO HERON MEAN AND THEIR SCHUR CONVEXITIES

K. M. Nagaraja¹
Ambika M Hejib
Sampathkumar. R
Venkataramana B S

Keywords:

*Classical means, Heron mean, Schur
Convexity, Schur geometric Convexity.*



A B S T R A C T

In this paper, the complementary of arithmetic mean, geometric mean, harmonic mean and contra harmonic mean with respect to Heron mean are defined. Further, by finding the partial derivatives developed the Schur convexity and Schur geometric convexity results.

© 2021 Published by Faculty of Engineeringg

1. INTRODUCTION

The well-known means in literature namely; Arithmetic mean, Geometric mean and Harmonic mean are presented by Pappus of Alexandria [Bullen, 2003]. In Pythagorean school on the basis of proportion, 10 Greek means are defined, among these means the well-known in literature are given as follows: For two positive real numbers a and b ; $A(a, b) = \frac{a+b}{2}$; $G(a, b) = \sqrt{ab}$; $H(a, b) = \frac{2ab}{a+b}$ and $C(a, b) = \frac{a^2+b^2}{a+b}$. These are respectively called Arithmetic mean (A.M), Geometric mean (G.M), Harmonic mean (H.M) and Contraharmonic mean (C.M). In 1958, C. Gini introduced Complementary means and G.Toader in 1991 proposed a generalization of complementariness and inversion (Toader, S. & Toader, G. 2005). In 1923,

Issai Schur introduced the concept of Schur convexity, which has applications in linear regression, analytic inequalities, Gamma functions, graphs and matrices, stochastic orderings, combinatorial optimization, information theoretic topics, reliability and related fields.

2. LITERATURE REVIEW

The Schur geometrical convexity and concavity of the extended and other mean values were discussed by (Chu, Y. M. et al, 2008, Shi, H. N. et al, 2010, Murali, K. & Nagaraja, K. M. 2013, 2016, Janardhana, L. et al, 2017)

Results on Schur convex and concave functions studied by (Elezovic, N. et al, 1998, Janardhana, L. et al, 2017a,

¹ Corresponding author: K M Nagaraja
Email: nagkmm@gmail.com

2017b, Lokesh, et al, 2010a, 2010b, 2011, Marshall, A. M. & Olkin, I. 1979, Nagaraja, K. M. et al, 2008, 2010, 2011, 2013, 2014a, 2014b, 2014c, 2017, Naveen Kumar, B. et al, 2014, Sampath Kumar, R. et al., 2017a, 2017b).

The Schurharmonic convexity and concavity results were discussed in (Nagaraja, K. M. et al., 2013a, 2013b). This paper was developed based on the above literature survey.

3. DEFINITIONS AND LEMMAS

Definition 1: (Toader, S. & Toader, G. 2005) A mean N is called complementary to M with respect to P (or P -complementary to M) if it verifies $P(M, N) = P$, it is denoted by $M^{(P)} = N$.

1. The complementary of geometric mean with respect to Heron mean is denoted by $G^{(He)}$ and is given by

$$G^{(He)} = \frac{1}{2} \left[2a + 2b + \sqrt{ab} - \sqrt{4(a+b)\sqrt{ab} + ab} \right].$$

2. The complementary of arithmetic mean with respect to Heron mean is denoted by $A^{(He)}$ and is given by

$$A^{(He)} = \frac{1}{4} \left[3a + 3b + 4\sqrt{ab} - \sqrt{(a+b)(5a+5b+8\sqrt{ab})} \right].$$

3. The complementary of harmonic mean with respect to Heron mean is denoted by $H^{(He)}$ and is given by

$$H^{(He)} = a + b + \sqrt{ab} - \frac{ab}{a+b} - \frac{\sqrt{2ab(a+b)^2 + 2\sqrt{ab}(a+b) - 3ab}}{a+b}$$

4. The complementary of contra harmonic mean with respect to Heron mean is denoted by $C^{(He)}$ and is given by

$$C^{(He)} = 2(a+b+\sqrt{ab}) - \frac{a^2+b^2}{a+b} - \frac{\sqrt{[(a^2+b^2)+8ab+4\sqrt{ab}(a+b)](a^2+b^2)}}{a+b}$$

Definition 2: [A. M. Marshall and I. Olkin, 1979] Let $x = (x_1, x_2, \dots, x_n)$ and $y = (y_1, y_2, \dots, y_n) \in R^n$
 (i) x is majorized by y , (in symbol $x \preceq y$). If $\sum_{i=1}^k x_i \leq \sum_{i=1}^k y_i$, $\sum_{i=1}^n x_i = \sum_{i=1}^n y_i$ where $x_1 \geq x_2 \geq \dots \geq x_n$ and $y_1 \geq y_2 \geq \dots \geq y_n$ are rearrangements of x and y in descending order. (ii) $\Omega \subseteq R^n$ is called a convex set if

$(\alpha x_1 + \beta y_1, \dots, \alpha x_n + \beta y_n)$ for every $x, y \in \Omega$, where $\alpha, \beta \in [0, 1]$ with $\alpha + \beta = 1$.

Let $\Omega \subseteq R^n$ the function $\varphi: \Omega \rightarrow R$ is said to be a Schur convex function on Ω if $x \preceq y$ on Ω implies $\varphi(x) \leq \varphi(y)$. φ is said to be a Schur concave function on Ω if and only if $-\varphi$ is Schur convex.

Definition 3: Let $x = (x_1, x_2, \dots, x_n)$ and $y = (y_1, y_2, \dots, y_n) \in R^n_+$, $\Omega \subseteq R^n$ is called geometrically convex set if for every $x, y \in \Omega$, where $\alpha, \beta \in [0, 1]$ with $\alpha + \beta = 1$.

Let $\Omega \subseteq R^n$, the function $\varphi: \Omega \rightarrow R_+$ is said to be Schur geometrically convex function on Ω if $(\ln x_1, \ln x_2, \dots, \ln x_n) \preceq (\ln y_1, \ln y_2, \dots, \ln y_n)$ on Ω implies

$\varphi(x) \leq \varphi(y)$. Then φ is said to be a Schur geometrically concave function on Ω if and only if $-\varphi$ is Schur geometrically convex.

Definition 4: $\Omega \subseteq R^n$ is called symmetric set if $x \in \Omega$ implies $Px \in \Omega$ for every $n \times n$ permutation matrix P .

The function $\varphi: \Omega \rightarrow R$ is called symmetric if for every permutation matrix P , $\varphi(Px) = \varphi(x)$ for all $x \in \Omega$.

Definition 5: Let $\Omega \subseteq R^n$ $\varphi: \Omega \rightarrow R$ be symmetric and convex function, then φ is Schur convex on Ω .

Lemma 1: Let $\Omega \subseteq R^n$ be symmetric with non-empty interior geometrically convex set and let $\varphi: \Omega \rightarrow R_+$ be continuous on Ω and differentiable in Ω^0 . If φ is symmetric on Ω and

$$(x_1 - x_2) \left(\frac{\partial \varphi}{\partial x_1} - \frac{\partial \varphi}{\partial x_2} \right) \geq 0 (\leq 0)$$

$$(\ln x_1 - \ln x_2) \left(x_1 \frac{\partial \varphi}{\partial x_1} - x_2 \frac{\partial \varphi}{\partial x_2} \right) \geq 0 (\leq 0)$$

$$(x_1 - x_2) \left(x_1^2 \frac{\partial \varphi}{\partial x_1} - x_2^2 \frac{\partial \varphi}{\partial x_2} \right) \geq 0 (\leq 0)$$

is a Schur convex (concave), Schur-geometrically convex (concave) and Schur-harmonically convex (concave) function respectively.

4. MAIN RESULTS

Theorem 1: For $a, b > 0$, the complementary of arithmetic mean with respect to Heron mean is Schur concave.

Proof: The complementary of arithmetic mean with respect to Heron mean is given by;

$$A^{He} = \frac{1}{4} \left[3a + 3b + 4\sqrt{ab} - \sqrt{(a+b)\{5a+5b+8\sqrt{ab}\}} \right]$$

The partial derivatives of A^{He} with respect to a and b are given by

$$\frac{\partial A^{He}}{\partial a} = \frac{1}{4} \left[3 + \frac{2b}{\sqrt{ab}} - \frac{\frac{4b(a+b)}{\sqrt{ab}} + 10(a+b) + 8\sqrt{ab}}{2\sqrt{5(a+b)^2 + 8\sqrt{ab}(a+b)}} \right]$$

and

$$\frac{\partial A^{He}}{\partial b} = \frac{1}{4} \left[3 + \frac{2a}{\sqrt{ab}} - \frac{\frac{4a(a+b)}{\sqrt{ab}} + 10(a+b) + 8\sqrt{ab}}{2\sqrt{5(a+b)^2 + 8\sqrt{ab}(a+b)}} \right]$$

Consider,

$$\frac{\partial A^{He}}{\partial a} - \frac{\partial A^{He}}{\partial b} = \frac{1}{2} \left[\frac{b-a}{\sqrt{ab}} - \frac{\frac{(b-a)(a+b)}{\sqrt{ab}}}{2\sqrt{5(a+b)^2 + 8\sqrt{ab}(a+b)}} \right]$$

Then,

$$(a-b) \left[\frac{\partial A^{He}}{\partial a} - \frac{\partial A^{He}}{\partial b} \right] = \frac{(a-b)^2}{2\sqrt{ab}} \left[\frac{a+b - \sqrt{5(a+b)^2 + 8\sqrt{ab}(a+b)}}{\sqrt{5(a+b)^2 + 8\sqrt{ab}(a+b)}} \right]$$

Clearly, $0 < 4(a+b)^2 + 8\sqrt{ab}(a+b)$, adding $(a+b)^2$ on both sides, then

$$a+b < \sqrt{5(a+b)^2 + 8\sqrt{ab}(a+b)}$$

Thus, $(a-b) \left[\frac{\partial A^{He}}{\partial a} - \frac{\partial A^{He}}{\partial b} \right] < 0$.

Hence the proof of the theorem 1.

Theorem 2: For $a, b > 0$, the complementary of arithmetic mean with respect to Heron mean is Schur geometrically convex.

Proof: The complementary of arithmetic mean with respect to Heron mean is given by;

$$A^{He} = \frac{1}{4} \left[3a + 3b + 4\sqrt{ab} - \sqrt{(a+b)\{5a + 5b + 8\sqrt{ab}\}} \right]$$

The partial derivatives of A^{He} with respect to a and b are given by

$$\frac{\partial A^{He}}{\partial a} = \frac{1}{4} \left[3 + \frac{2b}{\sqrt{ab}} - \frac{\frac{4b(a+b)}{\sqrt{ab}} + 10(a+b) + 8\sqrt{ab}}{2\sqrt{5(a+b)^2 + 8\sqrt{ab}(a+b)}} \right]$$

and

$$\frac{\partial A^{He}}{\partial b} = \frac{1}{4} \left[3 + \frac{2a}{\sqrt{ab}} - \frac{\frac{4a(a+b)}{\sqrt{ab}} + 10(a+b) + 8\sqrt{ab}}{2\sqrt{5(a+b)^2 + 8\sqrt{ab}(a+b)}} \right]$$

Consider,

$$a \frac{\partial A^{He}}{\partial a} - b \frac{\partial A^{He}}{\partial b} = \frac{1}{4} \left[3(a-b) - \frac{5(b-a)(a+b) + 4\sqrt{ab}(b-a)}{2\sqrt{5(a+b)^2 + 8\sqrt{ab}(a+b)}} \right]$$

Then,

$$(lna - lnb) \left[a \frac{\partial A^{He}}{\partial a} - b \frac{\partial A^{He}}{\partial b} \right] = \frac{(lna - lnb)(a-b)}{4} \left[3 - \frac{5(a+b) + 4\sqrt{ab}}{\sqrt{5(a+b)^2 + 8\sqrt{ab}(a+b)}} \right]$$

Clearly, $3\sqrt{5(a+b)^2 + 8\sqrt{ab}(a+b)} - (5(a+b) + 4\sqrt{ab}) > 0$

Thus, $(lna - lnb) \left[a \frac{\partial A^{He}}{\partial a} - b \frac{\partial A^{He}}{\partial b} \right] > 0$.

Hence the proof of the theorem 2.

Theorem 3: For $a, b > 0$, the complementary of geometric mean with respect to Heron mean is Schur convex.

Proof: The complementary of geometric mean with respect to Heron mean is given by

$$G^{He} = \frac{1}{2} \left[2a + 2b + \sqrt{ab} - \sqrt{4(a+b)\sqrt{ab} + ab} \right]$$

The partial derivative of G^{He} with respect to a and b are given by

$$\frac{\partial G^{He}}{\partial a} = \frac{1}{4} \left[4 + \frac{b}{\sqrt{ab}} - \frac{b(\sqrt{ab} + 6a + 2b)}{\sqrt{ab}\sqrt{ab + 4\sqrt{ab}(a+b)}} \right]$$

and

$$\frac{\partial G^{He}}{\partial b} = \frac{1}{4} \left[4 + \frac{a}{\sqrt{ab}} - \frac{a(\sqrt{ab} + 6b + 2a)}{\sqrt{ab}\sqrt{ab + 4\sqrt{ab}(a+b)}} \right]$$

Then,

$$(a-b) \left[\frac{\partial G^{He}}{\partial a} - \frac{\partial G^{He}}{\partial b} \right] = \frac{(a-b)^2}{4\sqrt{ab}} \left[\frac{\sqrt{ab} + 2(a+b) - \sqrt{ab + 4\sqrt{ab}(a+b)}}{\sqrt{ab + 4\sqrt{ab}(a+b)}} \right]$$

Clearly, $\sqrt{ab} + 2(a+b) - \sqrt{ab + 4\sqrt{ab}(a+b)} > 0$

Thus, $(a-b) \left[\frac{\partial G^{He}}{\partial a} - \frac{\partial G^{He}}{\partial b} \right] > 0$.

Hence the proof of the theorem 3.

Theorem 4: For $a, b > 0$, the complementary of geometric mean with respect to Heron mean is Schur geometrically convex.

Proof: The complementary of geometric mean with respect to Heron mean is given by

$$G^{He} = \frac{1}{2} \left[2a + 2b + \sqrt{ab} - \sqrt{4(a+b)\sqrt{ab} + ab} \right]$$

The partial derivative of G^{He} with respect to a and b are given by

$$\frac{\partial G^{He}}{\partial a} = \frac{1}{4} \left[4 + \frac{b}{\sqrt{ab}} - \frac{b(\sqrt{ab} + 6a + 2b)}{\sqrt{ab}\sqrt{ab + 4\sqrt{ab}(a+b)}} \right]$$

and

$$\frac{\partial G^{He}}{\partial b} = \frac{1}{4} \left[4 + \frac{a}{\sqrt{ab}} - \frac{a(\sqrt{ab} + 6b + 2a)}{\sqrt{ab}\sqrt{ab + 4\sqrt{ab}(a+b)}} \right]$$

Then, $(\ln a - \ln b) \left[a \frac{\partial G^{He}}{\partial a} - b \frac{\partial G^{He}}{\partial b} \right] =$

$$\frac{(\ln a - \ln b)(a - b)}{4} \left[4 - \frac{4\sqrt{ab}}{\sqrt{ab + 4\sqrt{ab}(a+b)}} \right]$$

Clearly, $4\sqrt{ab + 4\sqrt{ab}(a+b)} - 4\sqrt{ab} > 0$

Thus, $(\ln a - \ln b) \left[a \frac{\partial G^{He}}{\partial a} - b \frac{\partial G^{He}}{\partial b} \right] > 0$.

Hence the proof of the theorem 4.

Theorem 5: For $a, b > 0$, the complementary of contra harmonic mean with respect to Heron mean is Schur concave.

Proof: The complementary of contra harmonic mean with respect to Heron mean is given by

$$C^{He} = 2a + 2b + 2\sqrt{ab} - \frac{a^2 + b^2}{a + b}$$

$$- \frac{\sqrt{(a^2 + b^2)^2 + (a^2 + b^2)[8ab + 4\sqrt{ab}(a+b)]}}{a + b}$$

The partial derivatives of C^{He} with respect to a and b are given by

$$\frac{\partial C^{He}}{\partial a} = 2 + \frac{b}{\sqrt{ab}} - \frac{2a}{a+b} + \frac{(a^2 + b^2)}{(a+b)^2}$$

$$+ \frac{\sqrt{(a^2 + b^2)^2 + (a^2 + b^2)[8ab + 4\sqrt{ab}(a+b)]}}{(a+b)^2}$$

$$\frac{(a^2 + b^2) \left[4a + 8b + \frac{2b(a+b)}{\sqrt{ab}} + 4\sqrt{ab} \right]}{2(a+b)\sqrt{(a^2 + b^2)^2 + (a^2 + b^2)[8ab + 4\sqrt{ab}(a+b)]}}$$

$$\frac{\left[4a + 8b + \frac{2b(a+b)}{\sqrt{ab}} + 4\sqrt{ab} \right] + 16a^2b + 8a(a+b)\sqrt{ab}}{2(a+b)\sqrt{(a^2 + b^2)^2 + (a^2 + b^2)[8ab + 4\sqrt{ab}(a+b)]}}$$

and

$$\frac{\partial C^{He}}{\partial b} = 2 + \frac{a}{\sqrt{ab}} - \frac{2b}{a+b}$$

$$+ \frac{(a^2 + b^2) + \sqrt{(a^2 + b^2)^2 + (a^2 + b^2)[8ab + 4\sqrt{ab}(a+b)]}}{(a+b)^2}$$

$$- \frac{(a^2 + b^2) \left[4b + 8a + \frac{2a(a+b)}{\sqrt{ab}} + 4\sqrt{ab} \right]}{2(a+b)\sqrt{(a^2 + b^2)^2 + (a^2 + b^2)[8ab + 4\sqrt{ab}(a+b)]}}$$

$$- \frac{16ab^2 + 8b(a+b)\sqrt{ab}}{2(a+b)\sqrt{(a^2 + b^2)^2 + (a^2 + b^2)[8ab + 4\sqrt{ab}(a+b)]}}$$

Consider,

$$\frac{\partial C^{He}}{\partial a} - \frac{\partial C^{He}}{\partial b} = \frac{b-a}{\sqrt{ab}} + \frac{2(b-a)}{a+b}$$

$$+ \frac{(a^2 + b^2) \left[4(a-b) + \frac{2(a-b)(a+b)}{\sqrt{ab}} \right]}{2(a+b)\sqrt{(a^2 + b^2)^2 + (a^2 + b^2)[8ab + 4\sqrt{ab}(a+b)]}}$$

$$+ \frac{(b-a)(16ab + 8\sqrt{ab}(a+b))}{2(a+b)\sqrt{(a^2 + b^2)^2 + (a^2 + b^2)[8ab + 4\sqrt{ab}(a+b)]}}$$

Then, $(a-b) \left[\frac{\partial C^{He}}{\partial a} - \frac{\partial C^{He}}{\partial b} \right] = -(a-b)^2 \left[\frac{1}{\sqrt{ab}} + \frac{2}{a+b} \right]$

$$+ \frac{-4(a^2 + b^2) - \frac{2(a^2 + b^2)(a+b)}{\sqrt{ab}} + 16ab + 8\sqrt{ab}(a+b)}{2(a+b)\sqrt{(a^2 + b^2)^2 + (a^2 + b^2)[8ab + 4\sqrt{ab}(a+b)]}}$$

$$= \frac{-(a-b)^2(\sqrt{a} + \sqrt{b})^2}{(a+b)\sqrt{ab}}$$

$$\left[1 + \frac{4ab - a^2 - b^2}{\sqrt{(a^2 + b^2)^2 + (a^2 + b^2)[8ab + 4\sqrt{ab}(a+b)]}} \right]$$

Clearly,

$$\sqrt{(a^2 + b^2)^2 + (a^2 + b^2)[8ab + 4\sqrt{ab}(a+b)]} + 4ab - a^2 - b^2 > 0$$

Thus, $(a-b) \left[\frac{\partial C^{He}}{\partial a} - \frac{\partial C^{He}}{\partial b} \right] < 0$.

Hence the proof of the theorem 5.

Theorem 6: For $a, b > 0$, the complementary of contra harmonic mean with respect to Heron mean is Schur geometrically concave.

Proof: The complementary of contra harmonic mean with respect to Heron mean is given by

$$C^{He} = 2a + 2b + 2\sqrt{ab} - \frac{a^2 + b^2}{a + b}$$

$$- \frac{\sqrt{(a^2 + b^2)^2 + (a^2 + b^2)[8ab + 4\sqrt{ab}(a+b)]}}{a + b}$$

The partial derivatives of C^{He} with respect to a and b are given by (refer theorem 5)

Consider,

$$a \frac{\partial C^{He}}{\partial a} - b \frac{\partial C^{He}}{\partial b} = \frac{(b-a)[-a^2 - b^2]}{(a+b)^2} + \left[\frac{b-a}{(a+b)^2} \right]$$

$$\left[\frac{(a^2 + b^2)(a^2 + b^2 + 4ab)}{\sqrt{(a^2 + b^2)^2 + (a^2 + b^2)[8ab + 4\sqrt{ab}(a+b)]}} + \frac{2\sqrt{ab}(a+b)(3a^2 + 3b^2 + 8ab)}{\sqrt{(a^2 + b^2)^2 + (a^2 + b^2)[8ab + 4\sqrt{ab}(a+b)]}} \right]$$

Then,

$$(lna - lnb) \left[a \frac{\partial C^{He}}{\partial a} - b \frac{\partial C^{He}}{\partial b} \right] = \frac{(lna - lnb)(b-a)}{(a+b)^2}$$

$$\left[\frac{(a^2 + b^2)(a^2 + b^2 + 4ab)}{\sqrt{(a^2 + b^2)^2 + (a^2 + b^2)[8ab + 4\sqrt{ab}(a+b)]}} - \frac{2\sqrt{ab}(a+b)(3a^2 + 3b^2 + 8ab)}{\sqrt{(a^2 + b^2)^2 + (a^2 + b^2)[8ab + 4\sqrt{ab}(a+b)]}} - a^2 - b^2 \right]$$

Clearly, above expression is positive.

Thus, $(lna - lnb) \left[a \frac{\partial C^{He}}{\partial a} - b \frac{\partial C^{He}}{\partial b} \right] < 0$.

Hence the proof of the theorem 6.

Theorem 7: For $a, b > 0$, the complementary of harmonic mean with respect to Heron mean is Schur concave.

Proof: The complementary of harmonic mean with respect to Heron mean is given by

$$H^{He} = a + b + \sqrt{ab} - \frac{ab}{a+b}$$

$$- \frac{\sqrt{ab[2(a+b)^2 + 2(a+b)\sqrt{ab} - 3ab]}}{a+b}$$

The partial derivatives of H^{He} with respect to a and b are given by

$$\frac{\partial H^{He}}{\partial a} = 1 + \frac{b}{2\sqrt{ab}} - \frac{b}{a+b}$$

$$+ \frac{ab + \sqrt{ab[2(a+b)^2 + 2\sqrt{ab}(a+b) - 3ab]}}{(a+b)^2}$$

$$- \frac{ab \left(\frac{b(a+b)}{\sqrt{ab}} + 4(a+b) + 2\sqrt{ab} - 3b \right)}{2(a+b)\sqrt{ab[2(a+b)^2 + 2\sqrt{ab}(a+b) - 3ab]}}$$

$$- \frac{b(2(a+b)^2 + 2\sqrt{ab}(a+b) - 3ab)}{2(a+b)\sqrt{ab[2(a+b)^2 + 2\sqrt{ab}(a+b) - 3ab]}}$$

and

$$\frac{\partial H^{He}}{\partial b} = 1 + \frac{a}{2\sqrt{ab}} - \frac{a}{a+b}$$

$$+ \frac{ab + \sqrt{ab[2(a+b)^2 + 2\sqrt{ab}(a+b) - 3ab]}}{(a+b)^2}$$

$$- \frac{ab \left(\frac{a(a+b)}{\sqrt{ab}} + 4(a+b) + 2\sqrt{ab} - 3a \right)}{2(a+b)\sqrt{ab[2(a+b)^2 + 2\sqrt{ab}(a+b) - 3ab]}}$$

$$- \frac{a(2(a+b)^2 + 2\sqrt{ab}(a+b) - 3ab)}{2(a+b)\sqrt{ab[2(a+b)^2 + 2\sqrt{ab}(a+b) - 3ab]}}$$

Then,

$$(a-b) \left(\frac{\partial H^{He}}{\partial a} - \frac{\partial H^{He}}{\partial b} \right) = \left[\frac{(a-b)^2}{2\sqrt{ab}(a+b)} \right] \times$$

$$\left[2\sqrt{ab} - a - b + \frac{2(a+b)^2 + 3\sqrt{ab}(a+b) - 6ab}{\sqrt{2(a+b)^2 + 2\sqrt{ab}(a+b) - 3ab}} \right]$$

Clearly,

$$2\sqrt{ab} - a - b + \frac{2(a+b)^2 + 3\sqrt{ab}(a+b) - 6ab}{\sqrt{2(a+b)^2 + 2\sqrt{ab}(a+b) - 3ab}} > 0$$

Thus $(a-b) \left(\frac{\partial H^{He}}{\partial a} - \frac{\partial H^{He}}{\partial b} \right) < 0$.

Hence the proof of the theorem 7.

Theorem 8: For $a, b > 0$, the complementary of harmonic mean with respect to Heron mean is Schur geometrically convex.

Proof: The complementary of harmonic mean with respect to Heron mean is given by

$$H^{He} = a + b + \sqrt{ab} - \frac{ab}{a+b}$$

$$- \frac{\sqrt{ab[2(a+b)^2 + 2(a+b)\sqrt{ab} - 3ab]}}{a+b}$$

The partial derivatives of H^{He} with respect to a and b are given by (refer theorem 7)

Consider,

$$(lna - lnb) \left(a \frac{\partial H^{He}}{\partial a} - b \frac{\partial H^{He}}{\partial b} \right) = (lna - lnb)(a-b) \times$$

$$\left[\left(1 + \frac{ab + \sqrt{ab[2(a+b)^2 + 2\sqrt{ab}(a+b) - 3ab]}}{(a+b)^2} \right) - \left(\frac{2ab(a+b) + ab\sqrt{ab}}{(a+b)\sqrt{ab[2(a+b)^2 + 2\sqrt{ab}(a+b) - 3ab]}} \right) \right]$$

Clearly,

$$\left(1 + \frac{ab + \sqrt{ab[2(a+b)^2 + 2\sqrt{ab}(a+b) - 3ab]}}{(a+b)^2} \right) - \left(\frac{2ab(a+b) + ab\sqrt{ab}}{(a+b)\sqrt{ab[2(a+b)^2 + 2\sqrt{ab}(a+b) - 3ab]}} \right) > 0$$

Thus, $(\ln a - \ln b) \left(a \frac{\partial H^{He}}{\partial a} - b \frac{\partial H^{He}}{\partial b} \right) > 0$.

Hence the proof of the theorem 8.

Conclusion: The results discussed in this paper have applications in linear regression, analytic inequalities, Gamma functions, graphs and matrices, stochastic orderings, combinatorial optimization, information theoretic topics, reliability and related fields.

References:

- Bullen, P.S. (2003), *Handbook of means and their inequalities*, Kluwer Acad. Publ., Dordrecht.
- Chu, Y. M. & X. M. Zhang. (2008), The Schur geometrical Convexity of the extended mean values, *J. Convex Anal.*, 15(4), 869-890.
- Elezovic, N. & Pecaric, J. (1998). Note on Schur-Convex functions, *Rocky Mountain J. Math.*, 29, 853-856.
- Jain Zang, H.N.S. & Da-mao, L. (2010). Schur-Geometric Convexity for Difference of Means, *Applied Mathematics E-Notes*, 10, 274-284.
- Lakshmi Janardhana, R. C, Nagaraja, K. M., Sampath Kumar, R. & Murali, K. (2017). Schur convexities and concavities of Generalized Heron Means, *International Journal of Pure and Applied Mathematics*, 117 (13), 69-79.
- Lakshmi Janardhana, R. C, Nagaraja, K.M. & Lokesha, V. (2017). Schur-Geometric Convexity of Stolarsky's Extended Mean Values. *International Journal of Pure and Applied Mathematics* 114(1), 77-90.
- Lakshmi Janardhana, R. C, Nagaraja, K. M., Sampath Kumar, R. & Murali, K. (2017). Convexity of Power Type Heron Mean With Respect to Power Mean, *International Journal of Pure and Applied Mathematics*, 117 (11), 155-163.
- Lokesha, V, Naveen Kumar B, K. M. Nagaraja, Abdelmejid Bayad and M. Saraj, (2010). New Means and its Properties, *Proceedings of Jangjeon Mathematical Society*, 14(3), 243-254.
- Lokesha, V., Nagaraja, K. M., Naveen Kumar, B. & Padmanabhan, S. (2010). Extension of Homogeneous Function, *Tamsui Oxford Journal of Mathematical Sciences*, 26(4), 443-450.
- Lokesha, V., Nagaraja, K.M., Naveen Kumar, B. & Wu, Y-D. (2011). Schur convexity of Gnan mean for positive arguments, *Notes on Number theory and discrete mathematics*, 17 (4), 37-41.
- Marshall, A. M. & Olkin, I. (1979). *Inequalities: Theory of Majorization and Its Application*, New York: Academic Press.
- Murali, K. & Nagaraja, K. M. (2016). Schur Convexity of Stolarsky's Extended Mean Values, *Journal of Mathematical Inequalities*, JMI-10-59, 10(3), 725-735.
- Nagaraja, K. M, Lokesha, V. & Padmanabhan, S. (2008). A Simple Proof on Strengthening and Extension of Inequalities, *Advanced Studies in Contemporary Mathematics*, 17(1), 97- 103.
- Nagaraja, K. M. & Reddy, P.S.K. (2011). Logarithmic convexity and concavity of some Double sequences, *Scientia Magna*, 7(2), 78-81.
- Nagaraja, K. M, Lokesha, V., Xiao, Z. G. & Zhang, Z.H. (2010). The Generalized Weighted Elementary Symmetric Mean, *Ultra Scientist*, 22(2) M, 651-657.
- Nagaraja, K. M. & Sudhirkumarsahu. (2013). Schur Geometric Convexity of Gnan Mean for Two Variables, *Journal of the International Mathematical Virtual Institute*, 3, 39-59.
- Nagaraja, K. M, Reddy, P.S.K. & Sridevi, K. (2013). Schur Harmonic Convexity of Gnan Mean for Two Variables, *Journal of the International Mathematical Virtual Institute*, 3, 61-80.
- Nagaraja, K. M. & Sudhirkumarsahu. (2013). Schur Harmonic Convexity of Stolarsky extended mean values, *Scientia Magna*, 9(2), 22-34.
- Nagaraja, K. M, & Vimala, T. (2017). Convexity and Concavity of Means, *International Journal of Research – GRANTHAALAYAH*, 5 (4), 92-97.
- Nagaraja, K. M, Siva Kota Reddy, P. & Sridevi, K. (2014). Different Kinds of Schur Convexities of Power Type Alpha-Centroidal Mean, *Advances and Applications in Mathematical Sciences*, 13(2), 91-109.
- Nagaraja, K. M, Murali, K. & Lokesha, V. (2014). Schur convexity and concavity of Gnan mean, *Proceedings of Jangjeon Mathematical Society* 17(3), 355-367.
- Naveen Kumar, B., Kumar, S., Lokesha, V. & Nagaraja, K. M. (2011). Ratio of difference of means and its convexity, *International e Journal of Mathematics and Engineering*, 2(2), 932-936

Sampath Kumar, R. & Nagaraja, K. M. (2017). The Convexities of Invariant Contra Harmonic Mean with Respect to Geometric Mean, *International Journal of Pure and Applied Mathematics* 116 (22),407-412.

Sampath Kumar, R., Nagaraja, K. M. & Reddy, P. S. K. (2017). Convexity of Stolarsky's mean with respect to well-known Greek means, *International Journal of Pure and Applied Mathematics* 117 (12), 401-407

Toader, S. & Toader, G. (2005). Greek means and Arithmetic and Geometric means, *RGMA monograph*, Australia. <http://rgmia.vu.edu.au/html>.

K M Nagaraja

J.S.S. Academy of Technical Education,
Bengaluru-560060,
INDIA
nagkmn@gmail.com

Ambika M Hejib

R N S Institute of technology,
Bengaluru-560098,
INDIA
hejib.ambika@gmail.com

Sampathkumar R

R N S Institute of technology,
Bengaluru-560098,
INDIA
r.sampathkumar1967@gmail.com

Venkataramana B S

K S Institute of technology,
Bengaluru-560109,
INDIA
venkynvs@gmail.com
