

[科學短訊 Scientific Communication]

<https://doi.org/10.5281/zenodo.8376702>

First Record of *Amblyomma helvolum* Koch (Ixodidae: Amblyomminae) Parasitism on *Diploderma swinhonis* (Günther) (Agamidae: Draconinae) on the Main Island of Taiwan

ACE KEVIN S. AMARGA^{1, 2, 3}, TZONG-HAN LIN²

¹Biodiversity Program, Taiwan International Graduate Program, Biodiversity Research Center, Academia Sinica, Nangang District, Taipei 11529, Taiwan. Email: ace_amarga061@yahoo.com

²School of Life Science, National Taiwan Normal University- Gongguan Campus, Wenshan District, Taipei 11677, Taiwan

³International Union for Conservation of Nature Species Survival Commission (IUCN SSC) Parasite Specialist Group

Abstract. This article presents the first recorded instance of the reptile tick *Amblyomma helvolum* infesting Swinhoe's Japalure (*Diploderma swinhonis*) on the main island of Taiwan. Prior to this, the only published record of *A. helvolum* infestation on *D. swinhonis* was from Orchid Island.

Key words: ectoparasite, Ixodidae, new island record, Squamata.

Diploderma swinhonis (Günther) (Swinhoe's japalure), commonly known as Swinhoe's Japalure, is an endemic lizard species native to Taiwan and its adjacent islands, such as Orchid Island and Green Island (Ota 1991). Originally described in the genus *Japalura*, it was subsequently reclassified under the genus *Diploderma* (Wang et al. 2019). This diurnal species has a wide distribution, primarily found at elevations below 1,500 meters, spanning various habitat types (Ota 1991). Its diet primarily consists of small invertebrates (Norval et al. 1991). Additionally, *D. swinhonis* has been identified as a host for several parasitic invertebrate groups, including Ixodida, Nematoda, and Pentastomida (Norval et al. 2014; Amarga et al. 2023).

On July 13, 2023, an adult *D. swinhonis* was collected from Bitan (24°57'14.8" N, 121°31'59.0" E), Xindian District, New Taipei City, Taiwan. An engorged tick larva was found on the lizard's lateral torso (Fig. 1A) and was preserved in 95% ethanol. This larva was identified as *Amblyomma helvolum* Koch (Ixodida: Ixodidae: Amblyomminae) based on Kawk et al. (2023) (Fig. 1B). The species is widely distributed in the Oriental and Australasian regions and has been recorded in multiple countries, including Indonesia, Laos, Malaysia, the Philippines, Singapore, Taiwan, Timor Leste, and Vietnam (Robbins 2005; Petney et al. 2019; Amarga et al. 2022; Oda et al. 2022; Kwak et al. 2023). It primarily parasitizes Squamata but has been found on mammals and testudines in rare cases (Guglielmone et al. 2020; Oda et al. 2022). The Taiwanese records on reptiles were summarized in Amarga et al. (2023). The first documentation of *A. helvolum* parasitizing *D. swinhonis* was in 2021 on Orchid Island, southeast of Taiwan (Amarga et al. 2023). In addition, we hypothesize that *A. helvolum* may also infest other *Diploderma* species that coexist with *D. swinhonis* such as *D. brevipes* (Gressitt) and *D. polygonatum* Hallowell.




Figure 1. *Amblyomma helvolum* larva on *Diploderma swinhonis*: (A) attachment site on the lower torso (yellow arrow); (B) dorsal habitus of the specimen.

稿件收到 Received: 7 August 2023

稿件接受 Accepted: 19 September 2023

稿件出版 Published: 25 September 2023

Acknowledgements

The authors would like to thank Dr. Hui-Yun Tseng (National Taiwan University) for providing laboratory space.

References

- Amarga, A. K. S., Lin, J.-W., Tseng, H.-Y., Robbins, R.G., Kwak, M. L. & Lin, S.-M. 2023. New host records of the reptile tick *Amblyomma helvolum* Koch (Ixodida: Ixodidae: Amblyomminae) in Taiwan. *Systematic & Applied Acarology* 28(5): 931–943.
- Amarga, A. K. S., Gestíada, G. A. & Cuevas, C. B. 2022. New locality record for the reptile tick *Amblyomma helvolum* Koch, 1844 (Acari: Ixodidae) on the Bicol Peninsula, Luzon Island, Philippines. *Thailand Natural History Museum Journal* 14(2): 87–91.
- Guglielmone, A. A., Petney, T. N. & Robbins, R. G. 2020. Ixodidae (Acari: Ixodoidea): descriptions and redescriptions of all known species from 1758 to December 31, 2019. *Zootaxa* 4871(1): 1–322.
- Kwak, M.L., Jones, M. D., Harman, M. E. A., Smith, S. N., D'souza, A., Knierim, T., Barnes, C. H., Waengsothorn, S., Amarga, A. K. S., Kuo, C.-C. & Nakao, R. 2023. The East Indies reptile tick *Amblyomma helvolum* Koch, 1844 (Acari: Ixodidae), taxonomy, biology and new host records, including the first record of human infestation. *Tick and Tick-borne Diseases* 14(6): 102224.
- Oda, F. H., Martins, T. F., Labruna, M. B., O'Shea, M. & Kaiser, H. 2022. Ticks (Acari: Ixodidae) of three Timor-Leste reptiles: first country of *Amblyomma helvolum*, with new interactions and an updated list of host species. *Ticks and Tick-borne Diseases* 14(2): 102060.
- Ota, H. 1991. Taxonomic redefinition of *Japalura swinhonis* Günther (Agamidae: Squamata), with a description of a new subspecies of *J. polygonata* from Taiwan. *Herpetologica* 47: 280–294.
- Norval, G., Huang, S.-C., Mao, J.-J., Goldberg, S. R. & Slater, K. 2012. Additional notes on the diet of *Japalura swinhonis* (Agamidae) from southwestern Taiwan, with comments about its dietary overlap with the sympatric *Anolis sagrei* (Polychrotidae). *Basic and Applied Herpetology* 26: 87–97.
- Norval, G., Goldberg, S. R., Bursey, C. R., Mao, J.-J. & Slater, K. 2014. Internal parasites of lizards from Taiwan. *Herpetological Conservation and Biology* 9(3): 484–494.
- Petney, T. N., Saijuntha, W., Boulanger, N., Chitimia-Dobler, L., Pfeffer, M., Eamudomkarn, C., Andrews, R. H., Ahamad, M., Putthasorn, N., Muders, S. V., Petney, D.A., Robbins, R. G. 2019. Ticks (Argasidae, Ixodidae) and tick-borne diseases of continental Southeast Asia. *Zootaxa* 4558(1): 1–89.
- Robbins, R.G. 2005. The ticks (Acari: Ixodida: Argasidae, Ixodidae) of Taiwan: A synonymic checklist. *Proceedings of the Entomological Society of Washington* 107(2): 245–253.
- Wang, K., Che, J., Lin, S.-M., Deepak, V., Aniruddha, D.-R., Jiang, K., Jin, J., Chen, H. & Siler, C.D. 2019. Multilocus phylogeny and revised classification for mountain dragons of the genus *Japalura* s.l. (Reptilia: Agamidae: Draconinae) from Asia. *Zoological Journal of the Linnean Society* 185(1): 246–267.

臺灣本島首次記錄南蛇花蜱（硬蜱科：花蜱亞科）寄生於斯文豪氏攀蜥（飛蜥科：飛蜥亞科）

艾斯^{1、2、3}、林宗翰²

¹ 中央研究院生物多樣性中心 生物多樣性臺灣國際研究生博士學位學程 11529 臺北市南港區

E-mail: ace_amarga061@yahoo.com

² 國立臺灣師範大學生命科學專業學院 11677 臺北市文山區

³ 國際自然保護聯盟物種存續委員 (IUCN SSC) 寄生蟲專家組

摘要：本文報導臺灣本島的南蛇花蜱 (*Amblyomma helvolum*) 寄生於斯文豪氏攀蜥 (*Diploderma swinhonis*) 的首次紀錄。在本文發表之前，臺灣唯一的南蛇花蜱寄生於斯文霍氏攀蜥的紀錄來自蘭嶼。

關鍵字：體表寄生生物、硬蜱科、島嶼新紀錄、有鱗目