

Profitability analysis of cut flower cultivation in Bangladesh: Constraints and opportunities

Md. Imrul Kaysar¹, Md. Shofiqul Islam², Mohammad Shamsul Hoq³ Sultana Parvin Mukta⁴ and A K M Golam Kausar⁵

Received 6 August 2023, Revised 15 December 2023, Accepted 25 December 2023, Published online 31 December 2023

ABSTRACT

Bangladesh has immense potential for cut flower production and export to the world market. Although the export earnings from cut flower cultivation are gradually increasing in our country, production and profitability are hindered by some constraints. The present study was undertaken in Jashore (Jhikargachha) and Dhaka (Savar) districts to examine the financial profitability, constraints, and opportunities of cut-flower cultivation in Bangladesh. A total of 120 cut flower cultivating farmers were selected for interview in 2019, taking 60 sample farmers from each location. Multi-stage random sampling method was followed to collect primary data. For profitability analysis, two cut flowers, namely gladiolus and rose, were selected for the study. The result indicates that per hectare total cost, net return, and benefit-cost ratio (BCR) were Tk. 687439, Tk. 261509 and 1.38, respectively, for gladiolus cultivation. On the other hand, net present value (NPV), internal rate of return (IRR), and BCR of rose cultivation were Tk. 2325762, 146%, and 1.46, respectively, indicating that rose production is highly profitable to the farmers of the study areas. Despite such potentialities, cut flower cultivation was constrained by the requirement of high initial investment, lack of modern varieties, lack of credit facility, lack of storage facility, attack by pests & diseases, and uncertain market price. However, there is a huge scope to increase cut flower production by removing constraints in Bangladesh that can boost up livelihoods, incomes, and living conditions of rural people.

Keywords: Cut flower, Gladiolus, Rose, BCR, IRR

Cite this article as: Kaysar, M.I., Islam, M.S., Hoq, M.S., Mukta, S.P. and Kausar, A.K.M.G. 2023. Profitability analysis of cut flower cultivation in Bangladesh: Constraints and opportunities. Int. J. Agril. Res. Innov. Tech. 13(2): 41-48. https://doi.org/10.3329/ijarit.v13i2.70853

Introduction

The fertile land and favorable climate conditions are suitable for flower production in Bangladesh. Flower production was initially begun in the late seventies in our country by some innovative farmers on a small-scale, but rapid commercial production was initiated in the mid-eighties, especially in the Jashore district (Jahan, 2009). Commercially grown most common flower species in Bangladesh are rose (Rosa indica), tuberose (Polianthes tuberosa), marigold (Tagetes *erecta*, *T. patula*) and gladiolus (*Gladiolus* spp.). Currently, around 10000 hectares of land are under cut flower cultivation, and approximately 5000 promising farmers are cultivating flowers and foliage as their sole livelihood (Ahmed et al., 2021; Jahan, 2009). More than 0.15 million people in Bangladesh are engaged in the floriculture business directly or

indirectly (Rakibuzzaman et al., 2018). Flowers are usually used in various social functions such as weddings, worshipping, interior decoration, religious functions, birthday parties, welcoming friends, or relatives as a symbol of beauty.

Bangladesh has a competitive advantage in flower production due to the availability of nursery plants and seeds, easy production technology, competitive prices, and export potentialities (Ahmed et al., 2021; Mou, 2012). The current size of the flower market is around Tk. 10 billion and flower is being grown across the country (Laboni et al., 2019). Bangladesh exports various kinds of flowers and floral products to different developed countries like Italy, the United States, Portugal, Arabia, Singapore, Japan, Germany, and France etc. (Laboni et al., 2019).

¹Pulses Research Centre and Regional Agricultural Research Station, Bangladesh Agricultural Research Institute (BARI), Ishwardi-6620, Pabna, Bangladesh

²Agricultural Economics Division, Bangladesh Sugarcrop Research Institute (BSRI), Ishwardi-6620, Pabna, Bangladesh

³Agricultural Economics Division, Bangladesh Agricultural Research Institute (BARI), Gazipur, Bangladesh

⁴Bangladesh Institute of Research and Training on Applied Nutrition (BIRTAN), Narayanganj, Bangladesh

⁵Postgraduate Student, Resource Economics Group, Humboldt University of Berlin, 10117 Germany

^{*}Corresponding author's email: shafiqbau07@gmail.com (Md. Shofiqul Islam)

The trade of flowers was worth approximately €15bn (£10.6bn) globally in 2015 (Ahmed et al., 2021). Bangladesh also exported cut flowers and foliage worth USD 0.8 million in FY 2021-22 (EPM, 2023). Despite high potentialities, production and export are constrained by some barriers such as the lack of storage facilities, the inappropriate temperature in the airplane (10°C), the lack of modern seedlings, the lack of credit facilities, and uncertain market prices, etc. (Jahan, 2009; Laboni et al., 2019).

Floriculture appeared as a winsome business because many farmers converted to floriculture from other crops for higher benefits (Mou, 2012; Rakibuzzaman et al., 2018). As a result, it not only generates employment opportunities for many, especially poor women but also enlarges the scope to export. However, due to various production problems and lack of government support, cut flower producers face losses, which can narrow the potentiality of flower exports (Chowdhury and Khan, 2015). So, realizing the

above importance a study was undertaken in two different locations of Bangladesh to examine the financial profitability and constraints of cut-flower production in Bangladesh to derive some policy recommendations supporting cut-flower production.

Materials and Methods

Two cut flowers, namely rose and gladiolus were selected for analyzing the profitability of cut flower cultivation. Based on area coverage, two Upazila from two districts, namely Jhikargachha Upazila from Jashore district and Savar Upazila from Dhaka district were selected for the study. A multistage random method was followed to collect data. A total of 120 farmers were selected for interview, taking 60 farmers from each location. Secondary data have been collected from the Bangladesh Bureau of Statistics (BBS), Hortex Foundation, Export Promotion Bureau (EPB), national & international journals, articles, and earlier research reports.





Fig. 1. Map of the study areas.

A mostly tabular method of analysis was followed to achieve the objectives of the study. The profit function and project analysis were used in the study. The profit equation was used to estimate the profitability of gladiolus flower cultivation (Islam *et al.*, 2016; Miah *et al.*, 2019). The net return of gladiolus cultivation was estimated as follows:

$$\pi_{ijk} = P_{ijk} Q_{ijk} - \left(TVC_{ijk} + TFC_{ijk}\right) \dots \dots \dots \dots (1)$$

Where, π = Net return from ith flower per hectare; P_{ijk} = Per unit price of ith flower (Tk. unit-i); Q_{ijk} = Quantity of ith flower (unit ha-i); TVC_{ijk} = Total variable cost of ith crops (Tk ha-i); TFC_{ijk} = Total fixed cost of ith crops (Tk ha-i); i (1...2)= number of flower; j (1..2)= number of location and k (1.....120) = number of farmers.

Data were categorized according to the year for the rose. The age of the rose garden was classified as 1st year, 2nd year, 3rd year, 4th year, 5th year, 6th-9th year, and 10th-12th year. In presenting the

results, tabular methods of analysis were used using descriptive statistics. The NPV, BCR and IRR have been calculated in the case of rose cultivation with the help of the following formula (Kaysar *et al.*, 2017; Kaysar *et al.*, 2019):

Where, B_t = Total benefit (Tk/ha) in t^{th} year; C_t = Total cost (Tk/ha) in t^{th} year; t = Number of year and i = interest (discount) rate.

$$IRR = L + \frac{NPV \text{ at } L}{NPV \text{ at } L - NPV \text{ at } H} \times (H - L) \dots (4)$$

Where L represents a lower discount rate and H represents a higher discount rate.

Results and Discussion

Production practices of cut flower cultivation

The most appropriate planting time of gladiolus and rose was October to December and mid-February to April in the study areas, respectively (Table 1). Farmers cultivated exotic varieties of gladiolus like Pink, Kolmi, Beguni, Sada, and Halod varieties. In case of roses, the farmers cultivated Lingkon and Mirinda varieties for grafting on the rootstock. Both rose and gladiolus farmers had little knowledge about modern

flower varieties. The plant-to-plant distance for gladiolus and rose cultivation was found to be 6 inches and 13-15 inches, respectively, while line-to-line spacing was found to be 6-9 inches and 13-18 inches, respectively (Table 1). The average no. of seedlings required for gladiolus and rose cultivation was 800-1000 and 130-136 decimal⁻¹, respectively, in the study areas. The average no. of weeding ranges 6-10 times in all areas. The irrigation requirement was higher for rose cultivation compared to gladiolus cultivation (Table 1).

Table 1. Production practices of cut flower cultivation by the farmers.

| Parameter | Gladio | olus | Rose | | |
|----------------------------|-------------------------------------|-----------------------------------|---------------------------|---------------|--|
| | Jashore (Jhikargachha) | Dhaka (Savar) | Jashore (Jhikargachha) | Dhaka (Savar) | |
| Variety used | Pink, Kolmi, Beguni, Sada, Halod | Pink, Kolmi, Sada, Halod | Lingkon | Mirinda | |
| Planting time | October-December | October-January Mid Feb-Mid April | | March-April | |
| Spacing | | | | | |
| Plant to plant (inch) | 6" | 6" | 13"-14" | 13"-15" | |
| Line to line (inch) | 6" | 9" | 15"-18" | 15"-18" | |
| No. of seedlings decimal-1 | 800-1000 | 800-1000 | 123-130 | 130-136 | |
| Age of seedling (months) | 9-10 | 9-10 | 2.5-4.0 | 2.5-6.0 | |
| No. of weeding year-1 | 6-7 | 7-8 | 12 | 10 | |
| No. of irrigation year-1 | 8-9 | 8-9 | 20 | 18 | |

Profitability analysis of gladiolus cultivation

The result indicates that, on average, 393 mandays ha⁻¹ of human labour were applied by gladiolus farmers, but it varied from district to district (Table 2). Farmers used 212485 corms as a seed per hectare in all areas. The gladiolus farmers applied per hectare 6612.5 kg cow-dung and 23 kg oilcake as an organic fertilizer in all the selected areas. The farmers of Dhaka district used more cow-dung (7382 kg ha⁻¹) relative to Jashore (5843 kg ha⁻¹) district. The farmers applied Urea,

TSP, DAP, MOP, and Gypsum fertilizers at the rate of 267 kg ha⁻¹, 806 kg ha⁻¹, 97 kg ha⁻¹, 151 kg ha⁻¹, and 124 kg ha⁻¹, respectively, but mostly in an unbalanced way (Table 2). Proper agronomic management practices like those that balanced fertilizer application can enhance rose and gladiolus yields in the study areas. There are lot of evidence that balanced fertilizer application practices can improve both crop yields and farming profitability (Islam *et al.*, 2022a; Islam *et al.*, 2022b).

Table 2. Input use pattern (ha-1) of gladiolus cultivation.

| Particulars | Jashore (Jhikargachha) | Dhaka (Savar) | All area |
|---------------------------|------------------------|---------------|----------|
| Human Labour (man-days) | 372 | 416 | 394 |
| Own labour | 164 | 193 | 179 |
| Hired labour | 208 | 223 | 216 |
| Seed (Corm) | 220467 | 203933 | 212485 |
| Organic fertilizer (kg) | | | |
| Cow-dung | 5843 | 7382 | 6612.5 |
| Oilcake | 45 | | 23 |
| Inorganic fertilizer (kg) | | | |
| Urea | 333 | 196 | 267 |
| TSP | 793 | 820 | 806 |
| DAP | 156 | 33 | 97 |
| MOP | 178 | 122 | 151 |
| Gypsum | 137 | 109 | 124 |

However, the results show that the total production cost of gladiolus was Tk. 687439 ha⁻¹ in which the TVC was Tk. 670598 ha⁻¹ and the TFC was Tk. 16841 ha⁻¹ (Table 3). The total production cost of Savar farmers (Tk. 716064 ha⁻¹) was found to be higher relative to the farmers of the Jashore district (Tk. 659609 ha⁻¹) due to they

used comparatively lower inputs than farmers of the Dhaka district. Among different cost items, the results implied that seed cost was the highest (Tk. 460126 ha⁻¹), followed by human labour (Tk. 107592 ha⁻¹), fertilizer (Tk. 55577 ha⁻¹) and irrigation (Tk. 20592 ha⁻¹) (Table 3).

Table 3. Per hectare cost of gladiolus cultivation (Tk. ha-1).

| Particulars | Jashore (Jhikargachha) | Dhaka (Savar) | All Area |
|-------------------------------|------------------------|---------------|----------|
| A. Total variable cost (TVC) | 644639 | 697352 | 670598 |
| Land preparation | 12187 | 10577 | 11410 |
| Human labour | 92215 | 124067 | 107592 |
| Own labour | 40542 | 57258 | 48612 |
| Hired labour | 51674 | 66808 | 58980 |
| Seed | 456372 | 464147 | 460126 |
| Organic fertilizer | | | |
| Cow-dung | 14876 | 30002 | 22178 |
| Oilcake | 1449 | | 750 |
| Inorganic fertilizer | | | |
| Urea | 6287 3573 | | 4977 |
| TSP | 19392 | 21096 | 20215 |
| DAP | 4988 | 1127 | 3124 |
| MOP | 3025 | 3025 2223 | |
| Gypsum | 1295 | 1065 | 1627 |
| Irrigation | 17867 | 23512 | 20592 |
| Insecticide/Pesticide | 9879 | 10770 | 10309 |
| Interest in operating capital | 4806 | 5191 | 4992 |
| B. Total fixed cost (TFC) | | | |
| Land use cost | 14970 | 18712 | 16841 |
| Total Cost | 659609 | 716064 | 687439 |

two ways: as a flower (stick) and as a seed (corm). The yield of the gladiolus flower was around 179350 stick ha-1, whereas, in the case of corm yield, it was found at 146335 corm ha-1. On average, the gross return, gross margin, and net return of gladiolus cultivation was Tk. 948948 ha-

The returns came from gladiolus cultivation in 1, Tk. 278350 ha-1 and Tk. 261509 ha-1, respectively, whereas the average BCR was estimated at 1.38 in all selected areas. The highest BCR (1.45) was found in the Jashore district (Table 4).

Table 4. Per hectare yield, gross return, and net return of gladiolus cultivation.

| Particular | unit | Jashore (Jhikargachha) | Dhaka (Savar) | All area |
|---------------------------|------------------------|---------------------------|---------------|----------|
| Yield as a flower (stick) | no. ha-1 | 187397 | 171304 | 179350 |
| Yield as seed (corm) | no. ha-1 | 149918 | 142753 | 146335 |
| Average sale price flower | Tk. stick-1 | 3.5 | 4.0 | 3.8 |
| Average sale price corm | Tk. corm ⁻¹ | 2.0 | 1.8 | 1.9 |
| Gross return | Tk. ha-1 | 955726 | 942170 | 948948 |
| Total variable cost | Tk. ha-1 | 644639 | 697352 | 670598 |
| Gross margin | Tk. ha-1 | 311087 | 244818 | 278350 |
| Total cost | Tk. ha-1 | 659609 | 716064 | 687439 |
| Net return | Tk. ha-1 | 296117 | 226105 | 261509 |
| Benefit-cost ratio (BCR) | | 1.45 | 1.32 | 1.38 |

Profitability analysis of rose cultivation

The results show that, on average 1165 man-days ha-1 human labour was required for rose cultivation (Table 5). Farmers used an average of 6264.1 kg ha⁻¹ cow-dung. The highest 7567 kg ha⁻¹ cow-dung was used during the 1st year of gardening, whereas the lowest 5293 kg ha-1 was

used in the 3rd year of gardening. On average, farmers used 775 kg ha-1 of oilcake in rose cultivation. Similar to gladiolus, the rose farmers applied 319 kg ha⁻¹, 936 kg ha⁻¹, 51 kg ha⁻¹, 177 kg ha-1, and 172 kg ha-1 of Urea, TSP, DAP, MOP, and Gypsum, respectively, without following scientific recommendations (Table 5).

Table 5. Input use pattern (ha⁻¹) of rose cultivation.

| Particular | Particular | | | | | | | |
|---------------------------|-----------------|-----------------|-------------|-----------------|-------------------|----------------------------------|------------------------------------|----------|
| | 1 st | 2 nd | $3^{ m rd}$ | 4 th | 5^{th} | 6 th -9 th | 10 th -12 th | All year |
| Human Labour (man-days) | 1191 | 1142 | 1093 | 1188 | 1117 | 1252 | 1172 | 1165 |
| Own labour (man-days) | 496 | 515 | 533 | 466 | 411 | 486 | 444 | 479 |
| Hired labour (man-days) | 695 | 627 | 560 | 722 | 706 | 766 | 728 | 686 |
| Seedling (cutting) (no.) | 31481 | | | | | | | |
| Organic fertilizer (kg) | | | | | | | | |
| Cow-dung | 7567 | 6430 | 5293 | 7070 | 5939 | 5800 | 5750 | 6264.1 |
| Oilcake | 856 | 881 | 907 | 904 | 748 | | 1130 | 775 |
| Inorganic fertilizer (kg) | | | | | | | | |
| Urea | 386 | 409 | 431 | 216 | 297 | 124 | 368 | 319 |
| TSP | 1156 | 1016 | 875 | 1041 | 922 | 823 | 717 | 936 |
| DAP | 20 | 69 | 118 | | | 150 | | 51 |
| MoP | 170 | 158 | 146 | 115 | 300 | 124 | 225 | 177 |
| Gypsum | 200 | 240 | 281 | | 300 | 185 | | 172 |

Different cost items like human labour, seedlings, organic fertilizer, inorganic fertilizer, pesticides, irrigation, bamboo sticks, rope, net, etc., were included to estimate the total production cost of rose cultivation (Table 6). Land use cost was added as a fixed cost. The seedling cost was required only for the first year, and it was highest (Tk. 377786 ha⁻¹) among different cost items.

Among various cost items, human labour cost was the highest cost in the subsequent years. The average per hectare total cost was Tk. 634975 in all study areas. The highest cost (Tk. 980986 ha⁻¹) was observed in the 1st year garden due to the high cost of seedlings (Table 6).

Table 6. Per hectare total cost of rose cultivation (Tk. ha⁻¹) in the study areas.

| Particular | | | | Ye | ar | | | |
|------------------------|-----------------|-----------------|-------------|-----------------|-------------|----------------------------------|------------------------------------|----------|
| | 1 st | 2 nd | $3^{ m rd}$ | 4 th | $5^{ m th}$ | 6 th -9 th | 10 th -12 th | All year |
| A. Total variable cost | 913486 | 504395 | 502480 | 522149 | 497674 | 506847 | 525294 | 567475 |
| Land preparation | 7875 | | | | | | | 7875 |
| Human labour | 327498 | 313968 | 300438 | 326700 | 307175 | 344300 | 322163 | 320320 |
| Own labour | 136483 | 141529 | 146575 | 128150 | 113025 | 133650 | 122100 | 131644 |
| Hired labour | 191015 | 172439 | 153863 | 198550 | 194150 | 210650 | 200063 | 188676 |
| Seed | 377776 | | | | | | | |
| Organic fertilizer | | | | | | | | |
| Cow-dung | 46053 | 41144 | 36234 | 29656 | 31951 | 36960 | 46200 | 38314 |
| Oilcake | 27398 | 28203 | 29008 | 28928 | 23936 | | 36160 | 24805 |
| Inorganic fertilizer | | | | | | | | |
| Urea | 8110 | 8581 | 9051 | 4536 | 6237 | 2480 | 7718 | 6673 |
| TSP | 30064 | 26407 | 22750 | 27053 | 23972 | 21398 | 18651 | 24328 |
| DAP | 626 | 2193 | 3760 | | | 4800 | | 1626 |
| Mop | 2890 | 2682 | 2474 | 1955 | 5100 | 1984 | 3825 | 2987 |
| Gypsum | 2004 | 2405 | 2805 | | 3000 | 1665 | | 1697 |
| Irrigation | 24781 | 35886 | 46991 | 48105 | 46424 | 44460 | 37250 | 40557 |
| Insecticide/Pesticide | 36887 | 42928 | 48970 | 55216 | 49879 | 48800 | 53329 | 48001 |
| Bamboo stick | 17796 | | | | | | | 2542 |
| Net and rope | 3729 | | | | | | | 533 |
| Interest in operating | 27405 | 15132 | 15074 | 15664 | 14930 | 15205 | 15759 | 17024 |
| capital | | | | | | | | |
| B. Total fixed cost | | | | | | | | |
| Land use cost | 67500 | 67500 | 67500 | 67500 | 67500 | 67500 | 67500 | 67500 |
| C. Total Cost (A+B) | 980986 | 571895 | 569980 | 589649 | 565174 | 574347 | 592794 | 634975 |

From Table 7, it is apparent that the highest yield (1133316 stem ha⁻¹) was found in the 2nd year garden, which started to decline from the 3rd to the subsequent year. Gross return was estimated as yield multiplied by the price of a rose stem. The price varied from area to area and season to season. The average gross return was obtained at

Tk. 863479 ha⁻¹ in which the highest gross return was found in 2nd year (Tk. 1246648 ha⁻¹) and the lowest in 1st year (Tk. 554066 ha⁻¹). The average net return was found (Tk. 314852 ha⁻¹) similarly highest in the 5th year (Tk. 674752 ha⁻¹). The negative net return was found in 1st year (Tk. 426921 ha⁻¹) due to high initial investment.

Table 7. Per hectare yield and return of rose cultivation in the study areas.

| Particular | Year | | | | | | | |
|---------------------|-----------------|-----------------|-------------|-----------------|-------------|----------------------------------|------------------------------------|----------|
| | 1 st | 2 nd | $3^{ m rd}$ | 4 th | $5^{ m th}$ | 6 th -9 th | 10 th -12 th | All year |
| Yield (stem ha-1) | 503696 | 1133316 | 1101835 | 944430 | 912949 | 787025 | 661101 | 863479 |
| Gross return | 554066 | 1246648 | 1212019 | 1038873 | 1004244 | 865728 | 727211 | 949827 |
| Total variable cost | 913486 | 504395 | 502480 | 522149 | 497674 | 506847 | 525294 | 567475 |
| Gross margin | -359421 | 742252 | 709539 | 516724 | 506570 | 358881 | 201917 | 382352 |
| Total cost | 980986 | 571895 | 569980 | 589649 | 565174 | 574347 | 592794 | 634975 |
| Net return | -426921 | 674752 | 642039 | 449224 | 439070 | 291381 | 134417 | 314852 |

In the study, the best discount factor was considered at 6.5% (See appendix Table A1 and A2). The discounted gross cost and benefit of present worth at a 6.5% rate of interest was Tk. 5094902 ha⁻¹ and Tk.7420663 ha⁻¹, respectively. The estimated average net present worth of rose cultivation at a 6.5% discount rate was Tk. 2325762 ha-1 in the selected locations. On the other hand, the estimated BCR was 1.46 at a 6.5% rate of interest, which indicates that rose farmers are getting higher profits compared to gladiolus farmers. In the rose cultivation project, the IRR 146%, which was greater than the opportunity cost of capital. Therefore, it is acceptable and highly profitable (See Appendix Table A1 and A2). Similar types of results are also reported by Jahan (2009) while examining the financial analysis of cut flower production. Cultivation of flowers can provide almost 3-5 times and 1.5-2.0 times higher profits compared to rice and vegetables, respectively, in Bangladesh (Mou, 2012).

Constraints of cut flower cultivation

The results show that about 86% of farmers reported that cut-flower cultivation required a

huge amount of initial investment, which might be 6-7 times higher than cereal crops (Table 8). Another 80% suffered a lack of improved variety of seed/seedlings. They used different imported exotic varieties of seed due to the unavailability of improved verity. The pest and diseases attack (75%), lack of credit facility (75%), lack of training facilities (75%), damage during rainy and foggy weather (62%), and lack of high technologyoriented floriculture (67%) were the major production problems reported by the farmers. The farmers also faced some marketing problems, such as the absence of cold storage facilities at the production point (85%), uncertain market price (80%), and seasonal demand (79%) (Table 8). Effective extension services like training, demonstration, etc. can improve farmers' scientific knowledge regarding flower production dramatically, which ultimately can boost flower production (Islam et al., 2023; Jahan, 2009). In addition, Govt. of Bangladesh should take the initiative to develop storage facilities in the cut flower growing areas so that farmers can store unsold flowers in cold storage to keep the flowers alive (Mou, 2012).

Table 8. Farmers' responses about constraints of cut flower cultivation.

| Constraint | Jashore (Jhikargachha) | Dhaka (Savar) | All Area (%) |
|---|---------------------------|------------------|-----------------|
| Production constraints (%) | | | |
| High initial investment requirement | 84 | 88 | 86 |
| Absence of improved variety | 82 | 78 | 80 |
| Lack of credit facility | 76 | 74 | 75 |
| Pest and disease attack | 76 | 74 | 75 |
| Damage during rainy and foggy weather | 64 | 60 | 62 |
| Lack of high technology-oriented floriculture | 62 | 72 | 67 |
| Lack of training facility | 76 | 74 | 75 |
| Marketing constraints (%) | | | |
| Uncertain market price | 82 | 78 | 80 |
| Seasonal demand | 74 | 84 | 79 |
| Absence of cold storage | 82 | 88 | 85 |
| Limited buyers | 66 | 62 | 64 |
| Dominance of market intermediaries | 74 | 84 | 79 |
| Social and other constraints (%) | | | |
| Thief and animal problems | 62 | 72 | 67 |
| Spoilage | 66 | 62 | 64 |

Conclusion and Policy Recommendation

There is a high market demand for increasing diversified use of various flowers in Bangladesh. It is clear from our study that cut flower production is more profitable than other cereal crops, though it requires high investment initially. The profit margin is comparatively high cultivation relative to rose gladiolus cultivation. Many new farmers are getting involved in flower cultivation, and it is being promoted gradually in our country. Lack of storage facilities, lack of modern varieties, lack of credit & training facilities, and uncertain market prices are hindering cut flower production. Following policy recommendations should be taken by the Govt. of Bangladesh to improve cut flower cultivation covering the above constraints.

- i. Cold storage facilities should be developed at the production point, as cut flowers are highly perishable.
- ii. Flower-growing farmers need to be informed about modern flower production technologies like fertilizer, HYV seed, irrigation water, the new variety of flowers, flower caps, packaging materials, etc., through training programs and demonstration.
- iii. Market information should be easily accessible to the farmers so that farmers can be aware of uncertain market prices. To ensure better prices, a contract growing system should be encouraged for the farmers.
- iv. Cut flower cultivation should be supported by the Govt. of Bangladesh to coordinate the public-private partnership (PPP).
- v. Institutional credit facilities should be made easily available for cut flower farmers.

References

- Ahmed, M.S., Bagum, T., Mondal, T. and Das, D. 2021. Use of improved post-harvest practices by the flower farmers: A study on some selected cut flowers. *Int. J. Agric. Educ. Ext.* 7(1): 424-431.
- Chowdhury, R. and Khan, F. 2015. Cut flower export from Bangladesh: Prospects, challenges, and propositions. *Manarat Int. Uni. Stud.* 5(1): 11-21.
- EPB. 2023. Monthly summary sheet of export performance (goods) for FY 2022-2023. Export Promotion Bureau, Bangladesh. Retrieved from http://www.epb.gov.bd
- Islam, M.S., Alam, M.J., Bell, R.W., Boyd, D., Hutchison, J. and Miah, M.M. 2023. Fertilizer use gaps of women-headed households under diverse rice-based cropping patterns: Survey-based evidence

from the Eastern Gangetic Plain, South Asia. *Heliyon*. 9(3): e14139.

https://doi.org/10.1016/j.heliyon.2023.e14139

Islam, M.S., Bell, R.W., Miah, M.A.M. and Alam, M.J. 2022a. Farmers' fertilizer use gaps relative to government recommendations in the saline coastal zone of the Ganges Delta. *Agron. Sustain. Dev.* 42: 59.

https://doi.org/10.1007/s13593-022-00797-1

Islam, M.S., Bell, R.W., Miah, M.A.M. and Alam, M.J. 2022b. Unbalanced fertilizer use in the Eastern Gangetic Plain: The influence of Government recommendations, fertilizer type, farm size and cropping patterns. *PLoS ONE*. 17(7): e0272146.

https://doi.org/10.1371/journal.pone.0272146

Islam M.S., Khatun, S., Kamruzzaman, M., Kaysar, M.I. and Islam, S. 2016. Economics of sugarcane cultivation in some selected char lands of Bangladesh. *Int. J. Bus. Manag. Soc. Res.* 2(2): 132-139.

https://doi.org/10.18801/ijbmsr.020216.15 Jahan, H. 2009. Production, post-harvest

handling and marketing of cut-flowers in Bangladesh: An Agribusiness Study. *SAARC J. Agric.* 7(2): 1-14.

Kaysar, M.I., Hoq, M.S., Islam, M.W., Islam, M.S. and Islam, M.T. 2019. Profitability Analysis of Papaya Cultivation in Some Selected Areas of Bangladesh. *Bangladesh J. Agril. Res.* 44(1): 115-126.

https://doi.org/10.3329/bjar.v44i1.40936

Kaysar, M.I., Hoq, M.S., Mia, M.S., Islam, M.S. and Islam, M.M. 2017. An economic analysis of Jara and Colombo lemon production in Bangladesh. *J. Bangladesh Agric. Univ.* 15(2): 289-296.

https://doi.org/10.3329/jbau.v15i2.35078

- Laboni, S.A., Promy, J.A. and Abdullah, S.Z. 2019. Export potentiality of flower industry: a case study on Bangladeshi flower industry. *North American Acad. Res.* 2(11): 251-271
- Miah, M.A.M., Rouf, M.A., Islam, M.S., Alam, M.J., Rashid, M.H., Boyd, D., Bell, R.W., Haque, M.E., Satter, M.A., Rahman, M.W., Jahan, H. and Hutchenson, J. 2019. Assessment of gaps in current fertilizer use by farmers and scientific recommendation in selected areas of Bangladesh. Krishi Gobeshona Foundation (KGF), BARC, Farmgate, Dhaka. 128p.
- Mou, N.H. 2012. Profitability of flower production and marketing system of Bangladesh. *Bangladesh J. Agril. Res.* 37(1): 77-95. https://doi.org/10.3329/bjar.v37i1.11179
- Rakibuzzaman, M., Rahul, S.K., Jahan, M.R., Ifaz, M.I. and Uddin, A.J. 2018. Flower industry in Bangladesh. *Int. J. Bus. Soc. Sci. Res.* 7(1): 50-56.

Appendix

A1. Financial analysis of rose cultivation

| Year | Gross cost | Goss benefit | Discount factor at 6.5% | Present worth of cost at 6.5% | Present worth of benefit at 6.5% | | | |
|-------|---------------|--------------|----------------------------|-------------------------------|----------------------------------|--|--|--|
| 1 | 980986 | 554066 | 0.939 | 921114 | 520249 | | | |
| 2 | 571895 | 1246648 | 0.882 | 504217 | 1099118 | | | |
| 3 | 569980 | 1212019 | 0.828 | 471857 | 1003368 | | | |
| 4 | 589649 | 1038873 | 0.777 | 458348 | 807540 | | | |
| 5 | 565174 | 1004244 | 0.730 | 412510 | 732978 | | | |
| 6 | 574347 | 865728 | 0.685 | 393620 | 593313 | | | |
| 7 | 574347 | 865728 | 0.644 | 369596 | 557101 | | | |
| 8 | 574347 | 865728 | 0.604 | 347038 | 523100 | | | |
| 9 | 574347 | 865728 | 0.567 | 325858 | 491173 | | | |
| 10 | 592794 | 727211 | 0.533 | 315797 | 387404 | | | |
| 11 | 592794 | 727211 | 0.500 | 296523 | 363760 | | | |
| 12 | 592794 | 727211 | 0.470 | 278425 | 341559 | | | |
| Total | | | | 5094902 | 7420664 | | | |
| | BCR=1.46 | | | | | | | |

A2. Financial analysis of rose cultivation

| Year | Incremental benefit | Lower discount at 145% | Higher discount at 150% | NPV at 145% | NPV at 150% |
|-------|------------------------|------------------------|-------------------------|----------------|----------------|
| 1 | -426921 | 0.408 | 0.400 | -174253 | -170768 |
| 2 | 674752 | 0.167 | 0.160 | 112412 | 107960 |
| 3 | 642039 | 0.068 | 0.064 | 43658 | 41090 |
| 4 | 449224 | 0.028 | 0.026 | 12468 | 11500 |
| 5 | 439070 | 0.011 | 0.010 | 4974 | 4496 |
| 6 | 291381 | 0.005 | 0.004 | 1347 | 1193 |
| 7 | 291381 | 0.002 | 0.002 | 550 | 477 |
| 8 | 291381 | 0.001 | 0.001 | 224 | 191 |
| 9 | 291381 | 0.000 | 0.000 | 92 | 76 |
| 10 | 134417 | 0.000 | 0.000 | 17 | 14 |
| 11 | 134417 | 0.000 | 0.000 | 7 | 6 |
| 12 | 134417 | 0.000 | 0.000 | 3 | 2 |
| Total | | | | 1499 | -3761 |
| | | IRR= | 146% | | |